REVIEWS

THE STRUCTURE AND FUNCTION OF GLUTAMATE RECEPTOR **ION CHANNELS**

Dean R. Madden

As in the case of many ligand-gated ion channels, the biochemical and electrophysiological properties of the ionotropic glutamate receptors have been studied extensively. Nevertheless, we still do not understand the molecular mechanisms that harness the free energy of agonist binding, first to drive channel opening, and then to allow the channel to close (desensitize) even though agonist remains bound. Recent crystallographic analyses of the ligand-binding domains of these receptors have identified conformational changes associated with agonist binding, yielding a working hypothesis of channel function. This opens the way to determining how the domains and subunits are assembled into an oligomeric channel, how the domains are connected, how the channel is formed, and where it is located relative to the ligand-binding domains, all of which govern the processes of channel activation and desensitization.

ION CHANNEL STRUCTURE



Ion gradients are widely used in transmembrane signalling. These gradients are established by the active transport of ions from low to high electrochemical potentials across the membrane, driven by coupling transport to an energetically favourable process, such as ATP hydrolysis or respiration. Once established, such a gradient is relatively stable, as the lipid bilayer is essentially impermeant to ions. Furthermore, the partitioning of ions is typically not electroneutral, so that in addition to the concentration gradient for each ionic species, an overall electrostatic potential is built up across the membrane. Electrochemical gradients represent sources of potential energy that can be tapped for a variety of purposes. In neurons, the encoding, transmission and integration of information relies primarily on the carefully choreographed control of transmembrane electrostatic potential by a host of ion channels. Fast synaptic communication between nerve cells is one such process, mediated by ion channels that are located in the postsynaptic membrane and activated by neurotransmitters that are released from the presynaptic cell.

The development of patch-clamp techniques that are sensitive to the opening and closing of even a single ion channel¹ has permitted the detailed characterization of the electrophysiological behaviour of many ligandgated ion channels, including those responsible for neurotransmission². Typically, ligand-gated ion channels are closed in the resting state, but open in response to the binding of agonist, allowing selected ions to flow down their electrochemical gradients through an internal pore. After activation, the channels undergo spontaneous desensitization, closing even in the continuing presence of agonist. In a synapse, such channels mediate local depolarization or hyperpolarization of the plasma membrane, depending on the channel's ion selectivity and the nature of the corresponding transmembrane gradients. The change in membrane potential represents a signal that can be further processed by the receiving cell, and the magnitude and duration of the signal depends on several factors, including the time course of neurotransmitter concentration in the cleft, and the kinetics and extent of channel activation and desensitization.

Ion Channel Structure Research Group, Max Planck Institute for Medical Research, Jahnstrasse 29, 69120 Heidelberg, Germany, and Department of Biochemistry, Dartmouth Medical School, Hanover, New Hampshire 03755, USA. e-mail: dean.madden@ dartmouth edu DOI: 10.1038/nrn725

Box 1 | Cloned mammalian glutamate receptors

Ionotropic glutamate receptors (iGluRs) with high affinity for the agonist AMPA (α -amino-5-methyl-3-hydroxy-4-isoxazole propionic acid) are homo- or hetero-oligomers composed of subunits known as GluRA–GluRD (or GluR1–GluR4)^{114–116}. They are responsible for fast excitatory synaptic signalling in the brain. In addition, recent evidence indicates that they are involved in the activity-dependent modulation of synaptic strength¹¹⁷, a key component of the synaptic plasticity that is thought to underpin learning and memory.

Kainate receptors are homo- or hetero-oligomers of the subunits GluR5–GluR7, KA1 and KA2 (REFS 118–122). Like the AMPA receptors, they can mediate excitatory synaptic signals, but they are also involved in modulating the presynaptic release of neurotransmitter, and therefore in regulating the strength of synaptic connections. The kainate receptors are also potential therapeutic targets in the treatment of epilepsy and pain¹²³.

A third family of glutamate receptors is activated by NMDA (N-methyl-D-aspartate). These receptors are formed as heteromers that contain both NR1 (REE 124) and NR2 (NR2A–NR2D) $^{125-128}$ subunits, and in some cases NR3 subunits (NR3A and NR3B) $^{129-131}$. NMDA receptors require co-agonism by glycine 132 or D-serine 133 , which bind to the NR1 subunits, and glutamate, which binds to the NR2 subunits. Another unique feature of NMDA receptors is that extracellular Mg $^{2+}$ blocks these channels at resting membrane potentials; this block is relieved by depolarization of the membrane, allowing NMDA receptors to act as integrators of synaptic activity 134,135 .

So, all three families of glutamate receptor contain multiple subunits that can co-assemble, but only within families $^{136-138}$, to produce many receptor combinations. In addition, alternative splicing and RNA editing generate further variability 11,139 . As these modifications can affect both the extent and kinetics of the channel's electrophysiological response, the iGluR family shows an enormous molecular and functional diversity that is likely to be physiologically important.

In addition to these iGluR subunits, two orphan mammalian subunits have been cloned $(\delta 1 \text{ and } \delta 2)^{140}$; although no ion channel functionality has yet been shown for these subunits, a mutation in one of them is responsible for the neurological defects associated with the lurcher phenotype in mice^{75,141}. Finally, iGluR homologues have also been cloned from a prokaryote⁴⁴ and from various invertebrates^{142,143} and plants¹⁴⁴. The GluR family is therefore likely to be evolutionarily old¹⁴⁵.

ALLOSTERIC

A term originally used to describe enzymes that have two or more receptor sites, one of which (the active site) binds the principal substrate, whereas the other(s) bind(s) effector molecules that can influence its biological activity. More generally, it is used to describe the indirect coupling of distinct sites within a protein, mediated by conformational changes.

P-LOOP SEQUENCE
A conserved structural motif found in many different ion channels that constitutes part of the channel pore.

PDZ DOMAIN

A peptide-binding domain that is important for the organization of membrane proteins, particularly at cell-cell junctions, including synapses. They can bind to the carboxyl termini of proteins or can form dimers with other PDZ domains. PDZ domains are named after the proteins in which these sequence motifs were originally identified (PSD95, Discs large, zona occludens 1).

How agonist binding triggers the temporally regulated, multistep sequence of functional transitions in ligand-gated ion channels is one of the fundamental questions in ion channel molecular biology. This question can be broken down according to the functional steps that are associated with agonist binding, channel activation, ion permeation and channel desensitization. As the site of agonist binding is usually distant from the ion pore, activation of the channel must be mediated by ALLOSTERIC coupling of conformational changes at the binding site to corresponding changes at the channel's gate.

Recent experimental breakthroughs have provided new information on several of these processes. For glutamate receptor ion channels, crystallographic analyses of the ligand-binding domain in various functional conformations have revealed several activationdependent conformational states and a dimerization interface that is implicated in receptor desensitization³⁻⁵. In parallel, crystallographic and electron-microscopic data have been obtained for a prokaryotic K+ channel^{6,7}, the Torpedo acetylcholine receptor (nAChR)8 and an acetylcholine-binding protein9, with general and specific implications for the working of ion channels. This review aims to integrate structural insights with new functional and biochemical data, and to describe advances in our understanding of the mechanism of glutamate receptor ion channel function that are emerging as a result.

iGluR: a modular receptor

Ligand-gated ion channels are generally formed as homo- or hetero-oligomeric assemblies of integral membrane protein subunits. On the basis of pharmacological and sequence-homology criteria, they can be divided into several families; one such family comprises glutamate-gated channels, which are found in the preand postsynaptic cell membranes of the central nervous system. These glutamate receptor ion channels (iGluRs) are responsible for most fast excitatory signalling in the brain, and are thought to contribute to the synaptic plasticity that has been implicated in our ability to learn and form memories. In keeping with the physiological importance of iGluRs, their dysfunction is implicated in a range of neuropathologies, including epilepsy, stroke damage and the perception of pain¹⁰. Pharmacologically distinct subfamilies of iGluR have been identified, characterized by their affinities for the synthetic agonists AMPA (α-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid), NMDA (N-methyl-p-aspartate) and kainate11 (BOX 1).

An important advantage of the iGluRs for studies of ion channel function is their modular construction. This allows a reductionist approach to the questions outlined above. As illustrated in FIG. 1 (side view), a typical iGluR subunit consists of an amino-terminal domain (NTD), a ligand-binding domain (S1S2), three transmembrane domains and a re-entrant pore loop, and a carboxy-terminal domain¹². The NTD shows sequence homology to the leucine/isoleucine/valine-binding protein LIVBP, one of the bacterial periplasmic binding proteins (PBPs)13. In the NR2 subunits (BOX 1), it forms or contributes to the binding site of many NMDA receptor modulators^{14–25}, and influences receptor desensitization^{26,27}. For the non-NMDA receptors, it contributes to receptor assembly^{28–30}. This domain is absent from the related kainate-binding proteins of frog, chick and goldfish brains^{31–33}. The S1S2 ligandbinding domain is formed by two sequences that share sequence and structural homology with the glutaminebinding protein QBP^{3,34,35}. The re-entrant pore loop is homologous to the P-LOOP SEQUENCES of other channels36,37, and is located between the first and second transmembrane domains, forming a pore domain similar to those found in K+ channels. In iGluRs, as in K+ channels, the P-loop sequence lines the channel and determines many of its electrophysiological properties³⁸⁻⁴¹. In addition, neuronal iGluRs contain a third transmembrane domain and a cytoplasmic carboxyterminal domain that interacts with PDZ-DOMAIN proteins of the postsynaptic density, which control the targeting and localization of these receptors⁴².

The modular design indicates that the iGluR might have been assembled from components, in line with the 'genes-in-pieces' hypothesis⁴³. Support for this idea came from the discovery of a prokaryotic glutamate receptor ion channel (GluR0) — a potential evolutionary 'missing link'⁴⁴. As GluR0 lacks an NTD and has only two transmembrane domains, it could have been formed by the fusion of a PBP with a primitive K⁺ channel⁴⁴. Like the K⁺ channel KcsA, the structure of




Figure 1 | The modular nature of iGluR subunits. The amino terminus of ionotropic glutamate receptor (iGluR) subunits is extracellular. An amino-terminal domain (NTD) is followed by the S1 half-domain, two transmembrane domains with an intervening re-entrant P loop, the S2 halfdomain and a third transmembrane domain. The carboxyl terminus is located in the cytoplasm, where it can interact with proteins of the postsynaptic density. The S1 and S2 half-domains form the iGluR ligand-binding domain, which is homologous to the bacterial glutamine-binding protein QBP. The structure of this domain has been determined crystallographically, and is shown in enlarged form as a ribbon diagram. It consists of two lobes (lobe I, blue; lobe II, red), separated by a ligand-binding cleft 4.

which has been determined by X-ray crystallography (see Protein Data Bank (PDB) entry 1BL8)6, GluR0 forms a glutamate-gated channel that is selective for potassium ions⁴⁴. As a result of this structural and functional homology of the pore-forming domain, the extensive structural and biophysical studies of K+ channel ion permeation can be mined for insights into iGluR function.

\$1\$2: a molecular piston

A fortunate consequence of the modularity of the iGluR construction is that its ligand-binding domain can be expressed in isolation from the rest of the subunit^{45–49}; isolated constructs from AMPA receptors have been shown to bind ligand with both high affinity and high specific activity^{45,46,50}. It is this characteristic of the protein that has provided the greatest insights into potential mechanisms of iGluR channel gating, as the conversion of the free energy of agonist binding into mechanical work takes place entirely within the context of a selfcontained domain that can be studied in isolation. This is consistent with the observation that gating of non-NMDA glutamate receptors seems to involve the independent association of four molecules of agonist, one with each of four receptor subunits⁵¹. By contrast, the agonist-binding site of nAChR is located at the interfaces between adjacent subunits in the oligomer^{9,52}.

The existence of the S1S2 domain was first postulated on the basis of the above-mentioned homology to a bacterial PBP that is specific for glutamine^{13,34}. X-ray crystallography has shown the PBPs to be bilobate molecules that bind their ligands using a 'Venus-flytrap' mechanism⁵³: in the absence of ligand, the cleft between the lobes is open, but in the presence of ligand, the cleft is closed, trapping the ligand within it⁵⁴. The cleft closure can be described as a rigid-body motion of one lobe relative to the other, typically involving a rotation of \sim 35–55° about an axis through the hinge^{54–56}. However, various PBP-homologous proteins have been identified, some of which show much smaller domain closures in response to ligand binding. These include the purine and Lac repressor proteins, which close by 17–24° and 6°, respectively^{57–59}. The smaller cleft closure seems to be dictated by the repressors' need to oligomerize⁶⁰, a constraint that would also be expected to apply to the iGluR⁶¹.

Since 1998, crystallographic insights have been obtained into the structure of the S1S2 domain core of the AMPA receptor GluR2. The overall structure of the domain core closely resembles that of OBP3, with two lobes packed together to form a ligand-binding cleft on one side of the molecule (FIG. 1). The transmembrane domains of the receptor are tethered to the core domain by peptides that are attached to the carboxyl terminus of S1, and the amino and carboxyl termini of S2. These attachment sites are clustered on the back and at the bottom end of the molecule, as shown in FIG. 1. Parallel small-angle X-ray scattering measurements of the molecular envelope of a larger S1S2 construct showed that the peptide tethers extend beyond the end of the core domain, forming an even more elongated particle⁶¹. These peptides are therefore distant from both the agonist-binding cleft and the hinge domain on the back of the molecule.

Further insights have now been provided by the determination of crystallographic structures of the S1S2 core, alone and in complexes with a number of antagonists and agonists4. One key finding is that the extent of cleft closure more or less tracks the extent of channel activation and desensitization (FIG. 2). In response to binding of the agonists glutamate (see PDB entry 1FTJ) and AMPA (PDB 1FTM), the S1S2 core undergoes a cleft closure of ~20° relative to its ligand-free (apo) conformation (PDB 1FTO)⁴ (FIG. 2). This closure is smaller than that seen for the bacterial PBP, and more in keeping with the conformational changes that are seen in oligomeric repressor proteins⁶¹. Most interestingly, the cleft closure in the presence of the agonist kainate (PDB 1FTK and 1FW0) was only ~12°, even smaller than that seen with glutamate or AMPA (FIG. 2). Kainate is a 'partial agonist' of AMPA receptors, inducing both a smaller extent of activation and a smaller degree of desensitization than AMPA or glutamate⁶². Finally, the antagonist 6,7-dinitroquinoxaline-2,3-dione (DNQX; PDB 1FTL) stabilizes the open conformation of the cleft. In addition to the extent of cleft closure, the direction of rotation varies between the different S1S2-agonist complexes4. Such behaviour has been observed previously in examining other PBP homologues⁶⁰, and might provide a second mechanism by which different agonists can induce different channel responses.




Figure 2 | **Cleft closure mirrors extent of activation.** Upper panel: schematic current traces indicating the response of non-NMDA (N-methyl-p-aspartate) ionotropic glutamate receptors to the antagonist 6,7-dinitroquinoxaline-2,3-dione (DNQX; left), the partial agonist kainate (middle) and the full agonist glutamate (right) 62 . Lower panel: corresponding structures of the S1S2 domain. DNQX binding to the S1S2 domain stabilizes the open, apo conformation of the domain, which is associated with the closed state of the channel (left). Binding of kainate (middle) induces a ~12° cleft closure in the S1S2 domain, and leads to smaller peak currents and lower levels of channel desensitization than does glutamate binding (right), which causes a ~20° cleft closure in the S1S2 domain 4 .

The closing of the cleft

The structures of the S1S2 core also provide a framework for understanding the stepwise interaction of agonist with the iGluR binding site (FIG. 3). STOPPED-FLOW FLUORESCENCE STUDIES of wild-type and mutant S1S2 domains show that agonists first dock to one side of the binding cleft, interacting with two side chains (Tyr451 and Glu403 in GluR4)63 that form a ridge-like structure at the edge of the open binding site4. This docking interaction is modelled in FIG. 3 (middle) on the basis of the structure of the ligand-free domain and the interactions of glutamate with lobe I in the co-crystal structure. In a subsequent, slower step, S1S2 undergoes a conformational change that locks the agonist into the closed cleft (FIG. 3, right). This locking step presumably corresponds to the cleft closure detected crystallographically, and involves a glutamate side chain (Glu706 in GluR4) that forms a hydrogen bond with the agonist α -amino group^{3,63}. This side chain repositions itself within the cleft as a result of ligand binding4. The hydrogen bond is part of a three-way network that links side chains on each lobe of the protein (Glu706 and Thr481) together with the agonist, and it might be responsible for the high stability of the resulting complex: the backward isomerization rate constant is only $\sim 10 \text{ s}^{-1}$ (REF. 63).

What is the mechanism of cleft closure? It seems to involve a pre-existing open closed equilibrium for the ligand-binding domain. In the absence of agonist, the balance of the equilibrium favours the open state; the balance is shifted in favour of the closed state in the presence of agonist, which interacts more strongly with this conformation (FIG. 4). Consistent with such an equilibrium, the structure of the ligand-free form of the GluR2

S1S2 core has an open binding cleft (FIG. 4, upper left)⁴, whereas that of the GluR0 S1S2 domain has a closed conformation (FIG. 4, upper right; see PDB 1IIW)⁵. All agonist–iGluR complexes studied so far have been closed (FIG. 4, lower right). However, crystallographic studies of the related ligand-binding domain of the metabotropic glutamate receptor mGluR1 have revealed a mixture of open and closed conformations in the presence of glutamate (FIG. 4, lower left and lower right; see PDB 1EWK)⁶⁴.

For the PBP, the existence of the equilibria shown in FIG. 4 is well established by biochemical analyses^{65–67}. In addition to the expected open, empty, and closed, ligandbound structures, crystallographic studies have shown PBP in both the closed, empty state⁶⁸ and the open, ligand-bound state^{69,70}. For the PBP, the open conformation is not universally favoured in the apo state. Although both the phosphate- and arabinose-binding proteins adopt a predominantly open conformation in the absence of ligand^{71,72}, the cleft of the dicarboxylatebinding protein DctP is predominantly closed in the absence of ligand, opens to allow ligand to enter, and then recloses⁷³. The kinetics of GluR4 S1S2 ligand binding are qualitatively similar to those of the phosphatebinding protein⁶³. As iGluR agonists do not have access to the binding site in the closed state^{4,5}, this indicates that the eukaryotic iGluR domains preferentially adopt an open conformation in the absence of ligand (FIG. 4, upper left).

Assuming a simple allosteric coupling between the states of the ligand-binding domain and the channel gate, observations of spontaneous cleft closure correlate well with evidence of spontaneous activation of prokaryotic, NMDA and certain mutant non-NMDA glutamate receptors. In particular, although spontaneous activation is not observed for GluR2, it is observed for the prokaryotic GluR0 channel and for NMDA receptors, presumably reflecting the conformational equilibrium observed for the isolated binding cleft^{5,74}. Furthermore, on introducing the Lurcher Mutation⁷⁵ into both AMPA and kainate receptor subunits, spontaneous activation can be detected in the form of membrane current leakage in the absence of agonist⁷⁶⁻⁷⁸. The closed-channel state of the wild-type eukaryotic receptors seems to be sufficiently stable to prevent spontaneous opening (see below); however, when destabilized by mutations, the conformational equilibrium becomes apparent. Spontaneous activation has also been reported for nAChR mutants⁷⁹. So, it is quite possible that entire receptor subunits exist in a conformational equilibrium that is differentially stabilized in the absence or presence of agonist (simple allostery), as has been proposed for iGluRs and other ion channels^{80,81}. However, even though there seems to be an allosteric switch between the well-defined end states, the linear freeenergy relationships of transition-state energies in nAChR mutants indicate that the switching process might involve a 'conformational wave' that propagates from the binding site to the channel gate⁸².

Whereas simple allostery seems to operate in iGluRs and other ligand-gated ion channel subunits, the question

STOPPED-FLOW FLUORESCENCE STUDIES

The properties of a fluorophore are dependent on its chemical environment, and can therefore be influenced by molecular interactions or conformational changes. In stopped-flow experiments, interacting molecules are rapidly mixed in a flow cell, and the temporal evolution of their fluorescence signals is monitored to determine the kinetics of the interaction.

LURCHER MUTATION A spontaneous gain-of-function mutation in the $\delta 2$ glutamate receptor gene of mice that causes neuronal cell death, leading to loss of motor control in heterozygotes and to early postnatal mortality in homozygotes




Figure 3 | Ligand binding to the S1S2 domain. From left to right, glutamate docks in the open ligand-binding cleft on lobe I (blue), near to side chains Tyr451 and Glu403, shown as modelled (middle). A subsequent, kinetically slower cleft closure allows the formation of a three-way hydrogen-bond network (right) between the glutamate α -amino moiety and side chains Thr481 and Glu706 from both sides of the cleft^{3,63}

remains as to whether these channels are gated by a concerted Monod-Wyman-Changeux (MWC) mechanism, such as that used to describe haemoglobin cooperativity⁸³. For the nAChR, several lines of evidence indicate a concerted mechanism⁷⁹. The ability of an iGluR point mutation to convert an antagonist into an agonist⁸⁴ indicates that these receptors might also have a concerted gating mechanism⁷⁹. However, reports that binding of agonist to individual iGluR subunits can lead to incremental gating of the channel⁵¹ would argue that concerted transitions are not always required. Using non-desensitizing GluR3/GluR6 chimaeras or cyclothiazide to block desensitization, and antagonist presaturation to slow the kinetics of agonist association, a three-step opening of the iGluR pore could be detected. The dwell times in the different states could be well described if it was assumed that the small-conductance state required the binding of two agonist molecules, and the medium- and large-conductance states required one additional agonist molecule each, with independent binding to each site⁵¹. Conductance sublevels have also been observed for various K⁺ channels; although the resulting models differ in detail from each other and from that proposed for the iGluR, these observations generally support the functional homology between iGluR and K+ channels85,86. The altered ion selectivity of the K⁺ channel substates supports the hypothesis that they represent partially gated channels (rather than showing rapid switching between fully open and closed states)87.

Pharmacological specificity

One of the curious observations provided by the first iGluR S1S2 structure was that the ligand-binding site showed an overall negative charge, despite the fact that its physiological agonist and many high-affinity synthetic ligands are themselves negatively charged³. A similar coulombic mismatch has been observed between the phosphate-binding protein and its ligand, and is apparently resolved in that case by the precise distribution of both hydrogen-bonding partners and dipolar interactions within the binding site^{72,88}.

Our understanding of the pharmacological specificity of the glutamate receptors has been greatly advanced by high-resolution structural analysis of the ligand-binding domain of GluR2 in a complex with several different ligands4. On the basis of these studies, it seems that complementarity of the agonist-binding pocket for various ligands is mediated by a series of 'subsites', not all of which need to be occupied for highaffinity binding of a given agonist; water molecules are sometimes found in unused subsites4. This strategy presumably accounts for the ability of the iGluR binding site to bind various pharmacological agents89. Unlike those of the PBP, the subsites responsible for interaction with the agonist side-chain moieties are located such that the ligand lies across, rather than along, the domain's hinge⁶³. In keeping with its status as a primordial glutamate receptor, the orientation of agonist in the GluR0 binding site corresponds to that found in the PBP⁵. Ultimately, a full understanding of the differential selectivity of the AMPA, NR1, NR2 and kainate receptor subunits will probably require comparative structural analysis of the corresponding binding-site complexes.




Figure 4 | Conformational transitions in the S1S2 domain. In the absence of ligand (top), the S1S2 domain of the glutamate receptor (GluR) probably exists in an open⇔closed equilibrium that is skewed towards the open form. Ligand can bind only to the open cleft (apo bound equilibrium; left-hand side). In the presence of agonist (bottom), the closed conformation is stabilized relative to the open conformation, shifting the open⇔closed equilibrium towards the right. Glu, glutamate.




Figure 5 | **Assembly of the iGluR. a** | In most crystal structures of ionotropic glutamate receptors (iGluRs), a dimer is formed between two S1S2 domains 4 . Dimerization is mediated by interactions on the back of lobe I (blue), and involves side chains that have been implicated in channel desensitization. The cleft closure associated with agonist binding moves the ends of lobes II (red) $\sim\!5$ Å apart. **b** | The 'dimer-of-dimers' model of iGluR assembly 30 . Monomers associate most strongly through interactions between their amino-terminal domains (NTDs) (star in middle figure). Dimers undergo a secondary dimerization, mediated by interactions in the S2 and/or transmembrane domains (stars in right-hand figure). The crystallographically observed S1S2 dimer probably corresponds to this secondary dimerization interaction.

In addition to structural studies, INFRARED SPECTROSCOPY offers an alternative method for detailed analysis of protein-ligand interactions. In combination with crystallographic structure, the vibrational spectra also allow the characterization of the ligand electronic bonding configurations stabilized by interaction with the binding site — data that is inaccessible to structural analysis alone. In the case of the iGluR, they show that the ligand charge distribution adapts to optimize binding; nevertheless, electronegative ligand moieties interact with electronegative functional groups in the binding site, and therefore may not be bound with maximum affinity90,91. Extended to include binding-site mutants, this approach offers the prospect of detailed structureactivity information that might prove useful in the design of new agonists and antagonists.

INFRARED SPECTROSCOPY
Many interatomic molecular vibrations show energy transitions that lead to the absorption of photons at characteristic frequencies in the infrared range. The energy transitions, and therefore the associated infrared absorption spectra, are highly sensitive to the stereochemical environment of the functional group(s) involved.

Putting the pieces together

Despite our schematic understanding of the sequence of building blocks that constitutes a glutamate receptor subunit (FIG. 1), we know neither the spatial disposition of these domains within a subunit nor the packing of the subunits required to form a functional ion channel. However, it is this molecular architecture that provides the context within which conformational changes induced in S1S2 by agonist binding can be communicated to the channel gate.

To understand how the conformational changes in the ligand-binding domain are harnessed to produce physiological signals, it will be necessary to determine the relative orientations of the domains within an oligomeric channel. On the basis of electrophysiological, biochemical and hydrodynamic analyses, iGluRs are thought to be tetramers^{28,30,51,92–94}. One set of subunit–subunit interactions seems to be mediated by the NTD^{29,30}. In cross-linking experiments, these domains have been shown to dimerize when expressed as soluble constructs²⁸.

The GluR4 S1S2 domain cannot be cross-linked in dimeric form²⁸. Moreover, dimers are not observed in solution-scattering studies at protein concentrations as high as 0.75 mM (REF. 61). However, the GluR2 S1S2 domain forms crystallographic or non-crystallographic dimers in many different crystal forms⁴ (FIG. 5a). The crystal environment provides access to far higher concentrations than can be achieved in solution: the effective concentration of a ~30-kD protein, such as S1S2, exceeds 20 mM, assuming standard values for solvent content (50%) and protein density (1.33 g ml⁻¹). The most straightforward interpretation of these results is that the dimerization of eukaryotic S1S2 domains is much weaker than that of the NTD. The physiological relevance of the weak S1S2 dimerization is supported by the fact that the dimer is observed in many different crystal forms, that the interface contains numerous amino-acid residues that are implicated in the modulation of iGluR desensitization95, and that it provides a plausible model for iGluR activation by the S1S2 cleft closure^{4,5} (see below). Interestingly, in the case of the GluR0 S1S2 domain, dimers are observed not only crystallographically, but also in solution by analytical centrifugation, with an equilibrium dissociation constant of $0.8 \mu M$ (REF. 5).

Recently, using epitope-tagged AMPA receptor chimaeras, Ayalon and Stern-Bach³⁰ provided evidence of a sequential assembly of tetrameric iGluR channels as dimers of dimers (FIG. 5). In this model, an initial subunit dimerization is mediated primarily by interactions between compatible NTDs (FIG. 5b, step 1: 'dimer of monomers'). However, assembly of functional receptors (FIG. 5b, step 2: 'dimer of dimers') requires that these dimers undergo a second dimerization, this time requiring compatibility between the S2 and transmembrane domains of the subunits. This secondary dimerization is not observed in the absence of NTD compatibility³⁰, indicating that it might be weaker than the primary dimerization. Given the domains involved, the crystallographically observed dimerization presumably corresponds to part of the secondary interaction. A dimer-of-dimers model of iGluR assembly is consistent with images of GluR2 homomers determined by electron microscopy, with dimensions of $11 \times 14 \times 17$ nm: a fourfold symmetric assembly would have been expected to have at least two approximately equal dimensions⁹⁴. A dimerof-dimers assembly is not unprecedented for ion channels: it has also been proposed for the cyclicnucleotide-gated channels96.

ELECTRON PARAMAGNETIC RESONANCE SPECTROSCOPY When an atom with an unpaired electron is placed in a magnetic field, the spin of the unpaired electron can align, either in the same direction as the field or in the opposite direction. Electron paramagnetic resonance (EPR) spectroscopy is used to measure the absorption of microwave radiation that accompanies the transition between these two states.

EC.

The concentration of agonist that evokes a half-maximal response.

INHIBITION CONSTANT A measure of affinity, determined by the displacement of a labelled reporter ligand.

Pull to open?

Models of iGluR activation and desensitization have been developed that account for the observations described above^{4,5}. In these models (FIG. 6), cleft closure in a given subunit pulls the transmembrane domains of that subunit away from the pore axis (FIG. 6, side view, middle panel), opening the channel gate. Subsequent slippage between subunits allows the gate to close again, while the subunits remain in the agonist-bound conformation (FIG. 6, side view, right panel).

This model derives from the following considerations^{4,5}. To a first approximation, the crystallographically observed S1S2 dimers are formed 'back to back', with the agonist-binding clefts of each monomer facing out from the twofold symmetry axis (FIG. 5a). Furthermore, the dimerization interaction is mediated exclusively by residues that are located in the upper lobe (FIG. 1, lobe I). As a result, cleft closure in the two domains would increase the separation between the two bottom lobes, which are connected to the transmembrane

domains, by ~5 Å (FIG. 1 and FIG. 6). As the domains are positioned such that cleft closure can also have a horizontal component (FIG. 5a), a model can be constructed in which the transmembrane domains are also twisted in a manner analogous to the rigid-body motions detected in the KcsA transmembrane domain by ELECTRON PARAMAGNETIC RESONANCE SPECTROSCOPY⁹⁷. Finally, side chains implicated by mutagenesis in the kinetics and/or extent of desensitization are found at the dimer interface. As it is known that agonist remains bound to iGluRs in the desensitized state98,99, it is likely that the domain remains in the closed-cleft conformation, indicating that interdomain rather than intradomain rearrangements are required.

However, the mechanism of desensitization must also account for the fact that the desensitized state shows much higher affinity for agonist than does the resting state, as indicated by the difference between EC₅₀ and Inhibition Constant values11, and by the characteristics of receptor desensitization in the presence of AMPA




Figure 6 | A model for iGluR activation and desensitization^{4,5}. Upper row: ionotropic glutamate receptor (iGluR) channel physiology. In response to the application of glutamate to mouse embryonic hippocampal neurons, non-NMDA (V-methyl-p-aspartate) iGluRs open and then rapidly desensitize to a steady-state level146. Middle row: schematic top view of channels in the resting, open and desensitized states. Binding of glutamate (Glu) triggers a conformational change that opens the ion pore (red lightning bolt). Although agonist remains bound, a further, most likely interdomain conformational change allows the channel to close again, Lower row: schematic side view of two of the four iGluR subunits in a channel. S1S2 cleft closure is coupled to a conformational change in the transmembrane domains that pulls open the ion pore (arrows, middle figure). Repacking of the subunits (arrows, right-hand figure) recloses the pore, even though S1S2 and the transmembrane domains remain in the closed-cleft conformation.

and kainate 98. Furthermore, the high ligand-binding affinity of S1S2 constructs more closely matches that of the desensitized than of the resting state $^{4,45,100}.$ How can these observations be reconciled with the proposal that the crystallographically observed agonist-bound state of S1S2 reflects its activated rather than its desensitized state? The low affinity of the resting state for agonist presumably reflects the conformational coupling of S1S2 cleft closure to the energetically unfavourable process of opening the ion channel — allostery in its original sense of interaction between remote binding sites⁸¹. If channel opening were sufficiently unfavourable, this could effectively eliminate the problem of spontaneous activation described above; mutations such as lurcher would act by destabilizing the closed channel relative to the open channel. Given such a situation, it is possible that the transition from the activated to the desensitized state might allow slight conformational changes within S1S2, leading to the formation of further energetically favourable interactions4. Alternatively, additional conformational changes might not be required within S1S2. Instead, desensitization might simply involve effective decoupling of cleft closure from the energetically unfavourable open state of the channel. This could occur either by intersubunit rearrangements or by rearrangements between the domains within a subunit; for example, as mediated by the connecting peptides. In any case, decoupling of S1S2 from the channel is unlikely to be absolute, as agonist remains trapped in the binding site during open-channel block99.

Pass the salt

An important question associated with the model shown in FIG. 6 is how conformational changes in the transmembrane domains can lead to ion permeation in the iGluR. Despite their apparent evolutionary and overall structural relationship, the iGluR and K⁺ channel ion pores differ in their selectivity profiles.

For the prototypical bacterial K+ channel, KcsA, crystallographic and electrophysiological analyses have provided a very detailed understanding of the mechanism of ion conduction^{6,7,101}. Potassium ions are dehydrated at the entrance of the channel by protein functional groups that are arranged to mimic the hydration shell, and the precision of the geometry required to achieve dehydration accounts for the observed high selectivity for potassium ions^{6,7}. Dehydrated ions are hastened through the channel by repulsive coulombic interactions between multiple ions bound in the permeation pathway^{6,101}.

This mechanism seems to be exploited by voltage-gated eukaryotic K+ channels¹02, and might also apply to the prokaryotic GluR0 channel, which shares the ion selectivity characteristics of the K+ channels⁴4. Nevertheless, it is unlikely to apply in detail to the eukaryotic glutamate receptors, which are permeable not only to both K+ and Na+, but, depending on subunit composition, to divalent cations as well¹0. Many other ligandgated ion channels also show broader ion selectivity, including the nAChR.

What channel opening mechanism is likely to account for the broader ion selectivity and incremental

gating of the iGluR? The 4.6-Å-resolution structure of the closed form of the Torpedo nAChR channel reveals a hole with a radius of \sim 3.5 Å along the channel axis at the position of the gate, which is only slightly smaller than the radius of hydrated sodium and potassium ions; this constriction seems to be surrounded by hydrophobic leucine side chains8. Furthermore, electron-microscopic analysis of nAChR freeze-trapped in the open state revealed rotation of the kinked helices lining the pore, leading to a widening of the narrowest part of the permeation pathway¹⁰³. It therefore seems likely that, in this case, the channel opens wide enough to allow ions to pass through together with their hydration shells. Consistent with this idea, the open-channel dimensions of the nAChR pore have been estimated as 6.4-8.4 Å (REFS 104-106).

Corresponding measurements give values in excess of 7 Å for recombinant non-NMDA receptors¹⁰⁷, and ~5.5 Å for NMDA receptors¹⁰⁸. So, it is likely that the broad cation selectivity of the iGluRs results from their ability to pass hydrated or partially hydrated ions through a pore that is significantly wider than that of the KcsA channel. Despite these differences, analysis of the iGluR pore-loop structure by scanning cysteine mutagenesis confirms its similarity to the K+ channel pore loop^{40,109}. Indeed, Ca²⁺ channels, which are also homologous to the K⁺ channels, seem to have large pores (~6 Å in diameter) that are highly selective for Ca2+, but also conduct monovalent and divalent cations in its absence¹¹⁰. Furthermore, K⁺ channel mutants have been identified that reduce or eliminate selectivity among monovalent cations (although not divalent cations), indicating that relatively minor sequence changes are sufficient to relax the highly selective interactions in the permeation pathway^{111,112}. Given the underlying structural similarities, it will be interesting to determine the mechanisms by which homologous P loops mediate the selective and geometrically precise dehydration of ions in some channels, but act as relatively non-selective molecular sluice gates in other channels.

Peptides leading to the channel

Another interesting structural question concerns the nature of the symmetry of the iGluR transmembrane domains. Most P-loop-containing channels are fourfold or pseudo-fourfold symmetrical assemblies. However, as described above, the iGluRs are thought to be assembled as dimers of dimers³⁰. If the transmembrane domains preserve a true or approximate fourfold symmetry, then the peptides connecting them to the extracellular domains will be required to mediate a transition from two- to fourfold symmetry. Alternatively, it might be that the transmembrane domains show a breakdown of fourfold symmetry, and that this explains the ion selectivity differences described above. In any case, the dimer-of-dimers assembly also raises questions about the independence and equivalence of agonist binding to each of the four iGluR subunits, suggested by the analysis of conductance substates⁵¹.

A further question concerns the role of the peptides that connect the PBP-homologous S1S2 core to the

transmembrane domains. Ranging in length from about 5 to 20 amino acids, these peptides would be responsible for coupling the motions between the ends of lobe II to the channel gate, and they have been implicated in determining the desensitization characteristics of the iGluR^{26,27,113}. What is not clear is whether the peptides act as passive 'tie rods', transmitting the bulk motion directly to all three transmembrane domains of a subunit, or whether they undergo independent conformational changes, in which case differential forces would be applied to the three transmembrane domains of a given subunit.

Concluding remarks

The modular design of the iGluR has made possible the most extensive structural analysis yet of the interaction of ligands with an ion channel binding site. Together with detailed biophysical analyses of the ligand-binding interaction, and the homology of the S1S2 domain core to the well-characterized PBP family, these data provide unparalleled insights into the conformational changes that convert the free energy of agonist binding into mechanical work that can be harnessed to drive channel activation.

In parallel, our understanding of glutamate receptor function has benefited both from new information on the assembly of the channels, and from the explosion of

structural information on the homologous bacterial K⁺ channel KcsA. This has led to the first schematic models of the steps that lead from agonist binding to channel activation and subsequent desensitization.

However, as outlined above, several intriguing questions remain. Given the likely functional differences in ion selectivity, it will be interesting to establish in molecular detail how the P-loop sequence mediates two fundamentally different gating mechanisms; that is, how the relatively rigid and highly selective ion pore of KcsA becomes a less selective, mechanically gated pore in the iGluR. Another important issue concerns the orientations of all the domains in an assembled tetrameric glutamate receptor channel relative to one another in various functional states. These molecular-packing interactions will determine how the closure of the four S1S2 domains is translated into an activation signal, and the nature of the postulated slippage required for desensitization. The details of these structural contacts might also reveal how the NTD and connecting peptides modulate the kinetics of the channel response. Except for the S1S2 dimerization, none of these interactions has been characterized in molecular detail. As structural information on larger domains or intact iGluRs begins to emerge, we can look forward to the development of a framework for understanding the biophysical basis of these versatile molecular machines.

- Hamill, O. P., Marty, A., Neher, E., Sakmann, B. & Sigworth, F. J. Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. Pflügers Arch. 391, 85-100 (1981)
- Sakmann, B. & Neher, E. (eds) Single Channel Recording (Plenum, New York, 1983).
- Armstrong, N., Sun, Y., Chen, G.-Q. & Gouaux, E. Structure of a glutamate-receptor ligand-binding core in complex with kainate. Nature 395, 913-917 (1998).

The first atomic-resolution structure of a glutamate receptor ligand-binding domain.

- Armstrong, N. A. & Gouaux, F. Mechanisms for activation and antagonism of an AMPA-sensitive glutamate receptor: crystal structures of the GluR2 ligand binding core. Neuron **28**, 165–181 (2000).
 - Crystallographic structures of the iGluR ligandbinding domain in the apo state and in a complex with agonists and antagonists revealed cleft closure on agonist binding. The extent of closure correlates with the extent of channel activation by a given agonist, providing the first model of iGluR activation
- Mayer, M. L., Olson, R. & Gouaux, E. Mechanisms for ligand binding to GluR0 ion channels: crystal structures of the alutamate and serine complexes and a closed appostate J. Mol. Biol. **311**, 815–836 (2001).
 - This study provides the first evidence of dimerization of the ligand-binding domain in solution. The crystallographic structure of the ligand-free prokaryotic iGluR ligand-binding domain revealed a closed cleft, indicating an open-closed equilibrium that is probably reflected in spontaneous activity of the channel.
- Doyle, D. A. et al. The structure of the potassium channel: molecular basis of K+ conduction and selectivity. Science
 - The first crystallographic structure of an ion channel, revealing mechanisms by which ion selectivity is combined with high flux. Zhou, Y., Morais-Cabral, J. H., Kaufman, A. & MacKinnon, R.
- Chemistry of ion coordination and hydration revealed by a K+ channel-Fab complex at 2.0 Å resolution. Nature 414, 43-48 (2001).
- Unwin, N. The Croonian lecture 2000. Nicotinic acetylcholine receptor and the structural basis of fast synaptic transmission. Phil. Trans. R. Soc. Lond. B 355, 1813-1829 (2000).

- Brejc, K. et al. Crystal structure of an ACh-binding protein reveals the ligand-binding domain of nicotinic receptors Nature 411, 269-276 (2001).
 - The crystal structure of an acetylcholine-binding protein pentamer reveals the location of ligandbinding sites at the subunit interfaces.
- Dingledine, R., Borges, K., Bowie, D. & Traynelis, S. F. The glutamate receptor ion channels. Pharmacol. Rev. 51, 7-61 (1999).
- Hollmann, M. & Heinemann, S. Cloned glutamate receptors. Annu. Rev. Neurosci. **17**, 31–108 (1994).
- Hollmann, M., Maron, C. & Heinemann, S. N-glycosylation site tagging suggests a three transmembrane domai topology for the glutamate receptor GluR1. *Neuron* **13**, 1331–1343 (1994).
- O'Hara, P. J. et al. The ligand-binding domain of metabotropic glutamate receptors is related to bacterial periplasmic binding proteins. Neuron 11, 41-52 (1993).
- Choi, Y. B. & Lipton, S. A. Identification and mechanism of action of two histidine residues underlying high-affinity Zn2 inhibition of the NMDA receptor. Neuron 23, 171-180 (1999)
- Fayyazuddin, A., Villarroel, A., Le Goff, A., Lerma, J. & Nevton, J. Four residues of the extracellular N-terminal domain of the NR2A subunit control high-affinity Zn2+ binding to NMDA receptors. Neuron 25, 683-694 (2000)
- Low, C. M., Zheng, F., Lyuboslavsky, P. & Traynelis, S. F. Molecular determinants of coordinated proton and zinc inhibition of N-methyl-p-aspartate NR1/NR2A receptors Proc. Natl Acad. Sci. USA 97, 11062-11067 (2000).
- Paoletti, P. et al. Molecular organization of a zinc binding N-terminal modulatory domain in a NMDA receptor subunit. Neuron 28, 911-925 (2000).
- Gallagher, M. J., Huang, H., Pritchett, D. B. & Lynch, D. R. Interactions between ifenprodil and the NR2B subunit of the N-methyl-p-aspartate receptor. J. Biol. Chem. 271, 9603-9611 (1996).
- Gallagher, M. J., Huang, H., Grant, E. R. & Lynch, D. R. The NR2B-specific interactions of polyamines and protons with the N-methyl-p-aspartate receptor. J. Biol. Chem. 272, 24971-24979 (1997)
- Brimecombe, J. C., Gallagher, M. J., Lynch, D. R. & Aizenman, E. An NR2B point mutation affecting haloperidol and CP101.606 sensitivity of single recombinant N-methylp-aspartate receptors. J. Pharmacol. Exp. Ther. 286, 627-634 (1998)

- 21. Gallagher, M. J., Huang, H. & Lynch, D. R. Modulation of the N-methyl-D-aspartate receptor by haloperidol: NR2Bspecific interactions. J. Neurochem. 70, 2120-2128 (1
- Shim, S. S., Grant, E. R., Singh, S., Gallagher, M. J. & Lynch, D. R. Actions of butyrophenones and other antipsychotic agents at NMDA receptors: relationship with clinical effects and structural considerations. Neurochem. Int. 34, 167-175 (1999).
- Lynch, D. R. et al. Pharmacological characterization of interactions of RO 25-6981 with the NR2B (£2) subunit. Eur. J. Pharmacol. 416, 185-195 (2001).
- Masuko, T. et al. A regulatory domain (R1-R2) in the amino terminus of the N-methyl-p-aspartate receptor: effects of spermine, protons, and ifenprodil, and structural similarity to bacterial leucine/isoleucine/valine binding protein. Mol. Pharmacol. 55, 957-969 (1999).
- Zheng, F. et al. Allosteric interaction between the amino terminal domain and the ligand binding domain of NR2A Nature Neurosci. 4, 894-901 (2001).
- Krupp, J. J., Vissel, B., Heinemann, S. F. & Westbrook, G. L. N-terminal domains in the NR2 subunit control desensitization of NMDA receptors. *Neuron* **20**, 317–327 (1998).
- Villarroel, A., Regalado, M. P. & Lerma, J. Glycineindependent NMDA receptor desensitization — localization of structural determinants. Neuron 20, 329-339 (1998).
- Kuusinen, A., Abele, R., Madden, D. R. & Keinänen, K. Oligomerization and ligand-binding properties of the ectodomain of the AMPA receptor subunit GluRD. J. Biol. Chem. 274, 28937-28943 (1999).
- Leuschner, W. D. & Hoch, W. Subtype-specific assembly of $\alpha\text{-amino-3-hydroxy-5-methyl-4-isoxazole}$ propionic acid receptor subunits is mediated by their N-terminal domains. J. Biol. Chem. **274**, 16907–16916 (1999).
- Avalon, G. & Stern-Bach, Y. Functional assembly of AMPA and kainate receptors is mediated by several discret protein-protein interactions, Neuron 13, 103-113 (2001). A detailed analysis of the intersubunit interactions that govern iGluR assembly indicates that the receptors are assembled as dimers of dimers
- Wada, K. et al. Sequence and expression of a frog brain complementary DNA encoding a kainate-binding protein. *Nature* **342**, 684–689 (1989).
- Gregor, P., Mano, I., Maoz, I., McKeown, M. & Teichberg, V. I. Molecular structure of the chick cerebellar kainate-binding subunit of a putative glutamate receptor. Nature 342,

- Wo, Z. G. & Oswald, R. E. Transmembrane topology of two kainate receptor subunits revealed by N-glycosylation. Proc. Natl Acad. Sci. USA 91, 7154–7158 (1994).
- Nakanishi, N., Schneider, N. A. & Axel, R. A family of glutamate receptor genes: evidence for the formation of heteromultimeric receptors with distinct channel properties. *Neuron* 5, 569–581 (1990).
- Hsiao, C. D., Sun, Y. J., Rose, J. & Wang, B. C. The crystal structure of glutamine-binding protein from Escherichia coli. J. Mol. Biol. 262, 225–242 (1996)
- Mol. Biol. 262, 225–242 (1996).
 Wo, Z. G. & Oswald, R. E. Unraveling the modular design of glutamate-gated ion channels. *Trends Neurosci.* 18, 181–168 (1995)
- Wood, M. W., VanDongen, H. M. & VanDongen, A. M. Structural conservation of ion conduction pathways in K channels and glutamate receptors. *Proc. Natl Acad. Sci.* USA 92, 4882–4886 (1995).
- Verdoorn, T. A., Burnashev, N., Monyer, H., Seeburg, P. H. & Sakmann, B. Structural determinants of ion flow through recombinant glutamate receptor channels. Science 252, 1715–1718 (1991).
- Burnashev, N. et al. Control by asparagine residues of calcium permeability and magnesium blockade in the NMDA receptor. Science 257, 1415–1419 (1992).
- Kuner, T., Wollmuth, L. P., Karlin, A., Seeburg, P. H. & Sakmann, B. Structure of the NMDA receptor channel M2 segment inferred from the accessibility of substituted cysteines. *Neuron* 17, 343–352 (1996).
 Panchenko, V. A., Glasser, C. P. & Mayer, M. L. Structural
- Panchenko, V. A., Glasser, C. R. & Mayer, M. L. Structural similarities between glutamate receptor channels and K⁺ channels examined by scanning mutagenesis. *J. Gen. Physiol.* 117, 345–359 (2001).
- Bolton, M. M., Blanpied, T. A. & Ehlers, M. D. Localization and stabilization of ionotropic glutamate receptors at synapses. Cell. Mol. Life Sci. 57, 1517–1525 (2000).
- Gilbert, W. Why genes in pieces? *Nature* 271, 501 (1978).
 Chen, G.-Q., Cui, C., Mayer, M. L. & Gouaux, E. Functional
- Chen, G.-Q., Cui, C., Mayer, M. L. & Gouaux, E. Functional characterization of a potassium-selective prokaryotic glutamate receptor. *Nature* 402, 817–821 (1999).
 The identification of a prokaryotic iGluR provides molecular evidence that the iGluRs are derived from the fusion of a PBP with a primitive K* channel.
 Kuusinen, A., Arvola, M. & Keinänen, K. Molecular
- Kuusinen, A., Arvola, M. & Keinänen, K. Molecular dissection of the agonist binding site of an AMPA receptor. EMBO J. 14, 6327–6332 (1995).

The first demonstration of the ability to express functional iGluR ligand-binding domains as independent soluble proteins.

- Chen, G.-Q. & Gouaux, E. Overexpression of a glutamate receptor (GluR2) ligand binding domain in *Escherichia coli*: application of a novel protein folding screen. *Proc. Natl Acad. Sci. USA* 94, 13431–13436 (1997).
- Keinänen, K., Jouppila, A. & Kuusinen, A. Characterization of the kainate-binding domain of the glutamate receptor GluR-6 subunit. *Biochem. J.* 330, 1461–1467 (1998).
- Ivanovic, A., Reiländer, H., Laube, B. & Kuhse, J. Expression and initial characterization of a soluble glycine binding domain of the N-methyl-p-aspartate receptor NR1 subunit. *J. Biol. Chem.* 273, 19933–19937 (1998).
 Miyazaki, J., Nakanishi, S. & Jingami, H. Expression and
- Miyazaki, J., Nakanishi, S. & Jingami, H. Expression and characterization of a glycine-binding fragment of the N-methyl-a-aspartate receptor subunit NR1. Biochem. J. 340, 687–692 (1999).
- Abele, R., Lampinen, M., Keinänen, K. & Madden, D. R. Disulfide bonding and cysteine accessibility in the α-amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptor subunit GluRD: implications for redox modulation of glutamate receptors. J. Biol. Chem. 273, 25132–25138 (1998).
- Rosenmund, C., Stern-Bach, Y. & Stevens, C. F.
 The tetrameric structure of a glutamate receptor channel. Science 280, 1596–1599 (1998).
 - The analysis of gating transitions in iGluR under nondesensitizing conditions revealed incremental opening of the conducting pore, consistent with independent gating by four subunits assembled as a tetramer.
- Oswald, R. E. & Changeux, J.-P. Crosslinking of a-bungarotoxin to the acetylcholine receptor from *Torpedo* marmorata by ultraviolet light irradiation. FEBS Lett. 139, 225–229 (1982).
- Mao, B., Pear, M. R., McCammon, J. A. & Quiocho, F. A. Hinge-bending in L-arabinose-binding protein. The 'Venus's-flytrap' model. J. Biol. Chem. 257, 1131–1133 (1982).
- Sharff, A. J., Rodseth, L. E., Spurlino, J. C. & Quiocho, F. A. Crystallographic evidence of a large ligand-induced hingetwist motion between the two domains of the maltodextrin binding protein involved in active transport and chemotaxis. *Biochemistry* 31, 10657–10663 (1992).

- Oh, B.-H. et al. Three-dimensional structures of the periplasmic lysine/arginine/ornithine-binding protein with and without a ligand. J. Biol. Chem. 268, 11348–11355 (1993).
- Oh, B.-H., Ferro-Luzzi Ames, G. & Kim, S.-H. Structural basis for multiple ligand specificity of the periplasmic lysinearginine-, ornithine-binding protein. J. Biol. Chem. 269, 26232, 26230 (1004)
- 26323–26330 (1994).
 57. Schumacher, M. A., Choi, K. Y., Zalkin, H. & Brennan, R. G. Crystal structure of LacI member, PurR, bound to DNA: minor groove binding by α helices. *Science* 266, 763–770 (1004).
- Schumacher, M. A., Choi, K. Y., Lu, F., Zalkin, H. & Brennan, R. G. Mechanism of corepressor-mediated specific DNA binding by the purine repressor. Cell 83, 147–155 (1995).
- Lewis, M. et al. Crystal structure of the lactose operon repressor and its complexes with DNA and inducer. Science 271, 1247–1254 (1996).
 Mowbray, S. L. & Björkman, A. J. Conformational changes
- Mowbray, S. L. & Björkman, A. J. Conformational changes of ribose-binding protein and two related repressors are tailored to fit the functional need. J. Mol. Biol. 294, 487–499 (1999).
- Ábele, R., Svergun, D., Keinänen, K., Koch, M. H. J. & Madden, D. R. A molecular envelope of the ligand-binding domain of a glutamate receptor in the presence and absence of annist. *Biochemistry* 38, 10940–10957 (1999)
- absence of agonist. *Biochemistry* 38, 10949–10957 (1999).
 Patneau, D. K., Vyklicky, L. Jr. & Mayer, M. L. Hippocampal neurons exhibit cyclothiazide-sensitive rapidly desensitizing responses to kainate. *J. Neurosci.* 13, 3496–3509 (1993).
- Abele, R., Keinänen, K. & Madden, D. R. Agonist-induced isomerization in a glutamate receptor ligand-binding domain: a kinetic and mutagenetic analysis. J. Biol. Chem. 275, 21355–21363 (2000).
 The kinetics of iGluR agonist binding reveal an
 - The kinetics of iGluR agonist binding reveal an intermediate, fast docking step that precedes the slower cleft closure; key side chains are identified in both steps.
- Kunishima, N. et al. Structural basis of glutamate recognition by a dimeric metabotropic glutamate receptor. Nature 407, 971–977 (2000)
 - The first crystallographic analysis of the ligandbinding domain of a metabotropic glutamate receptor shows open and closed conformations for both the empty and agonist-bound forms.
- Wolf, A., Shaw, E. W., Nikaido, K. & Ferro-Luzzi Ames, G. The histidine-binding protein undergoes conformational changes in the absence of ligand as analyzed with conformation-specific monoclonal antibodies. J. Biol. Chem. 269, 23051–23058 (1994)
- Careaga, C. L., Sutherland, J., Sabeti, J. & Falke, J. J. Large amplitude twisting motions of an interdomain hinge – a disulfide trapping study of the galactose–glucose binding protein. *Biochemistry* 34, 3048–3055 (1995).
- Walmsley, A. R., Shaw, J. G. & Kelly, D. J. Perturbation of the equilibrium between open and closed conformations of the periplasmic C4-dicarboxylate binding protein from *Rhodobacter capsulatus. Biochemistry* 31, 11175–11181 (1992).
- Flocco, M. M. & Mowbray, S. L. The 1.9 Å X-ray structure of a closed unliganded form of the periplasmic glucose/galactose receptor from Salmonella typhimurium. J. Biol. Chem. 269, 8931–8936 (1994)
- Sack, J. S., Saper, M. A. & Quiocho, F. A. Periplasmic binding protein structure and function. Refined X-ray structures of the leucine/isoleucine/valine-binding protein and its complex with leucine. J. Mol. Biol. 206, 171–191 (1989).
- Duan, X. Q., Hall, J. A., Nikaido, H. & Quiocho, F. A. Crystal structures of the maltodextrin/maltose-binding protein complexed with reduced oligosaccharides: flexibility of tertiary structure and ligand binding. *J. Mol. Biol.* 306, 1115–1126 (2001).
- Vermersch, P. S., Lemon, D. D., Tesmer, J. J. G. & Quiocho, F. A. Sugar-binding and crystallographic studies of an arabinose-binding protein mutant (Met108Leu) that exhibits enhanced affinity and altered specificity. *Biochemistry* 30, 6861–6866 (1991).
 Ledvina, P. S., Tsai, A.-L., Wang, Z., Koehl, E. & Quiocho, F. A.
- Ledvina, P. S., Tsai, A.-L., Wang, Z., Koehl, E. & Quiocho, F. A. Dominant role of local dipolar interactions in phosphate binding to a receptor cleft with an electronegative charge surface: equilibrium, kinetic, and crystallographic studies. Protein Sci. 7, 2550–2559 (1998).
 Walmsley, A. R., Shaw, J. G. & Kelly, D. J. The mechanism of
- Walmsley, A. R., Shaw, J. G. & Kelly, D. J. The mechanism of ligand binding to the periplasmic C4-dicarboxylate binding protein (DctP) from Rhodobacter capsulatus. J. Biol. Chem. 267, 8064–8072 (1992).
- Turecek, R., Vlachova, V. & Vyklicky, L. Spontaneous openings of NMDA receptor channels in cultured rat hippocampal neurons. Eur. J. Neurosci. 9, 1999–2008 (1997).

- Zuo, J. et al. Neurodegeneration in Lurcher mice caused by mutation in δ2 glutamate receptor gene. Nature 388, 769–773 (1997).
- Zheng, Y., Brockie, P. J., Mellem, J. E., Madsen, D. M. & Maricq, A. V. Neuronal control of locomotion in C. elegans is modified by a dominant mutation in the GLR-1 ionotropic glutamate receptor. Neuron 24, 347–361 (1999).
- Kohda, K., Wang, Y. & Yuzaki, M. Mutation of a glutamate receptor motif reveals its role in gating and δ2 receptor channel properties. Nature Neurosci. 3, 315–322 (2000)
- channel properties. Nature Neurosci. 3, 315–322 (2000).
 Schwarz, M. K. et al. Dominance of the lurcher mutation in heteromeric kainate and AMPA receptor channels. Eur. J. Neurosci. 14, 861–868 (2001)
- Neurosci. 14, 861–868 (2001).
 Changeux, J.-P. & Edelstein, S. J. Allosteric mechanisms in normal and pathological nicotinic acetylcholine receptors. *Curr. Opin. Neurobiol.* 11, 369–377 (2001).
- Paas, Y. The macro- and microarchitectures of the ligandbinding domain of glutamate receptors. *Trends Neurosci.* 21, 117–125 (1998).
- Changeux, J.-P. & Édelstein, S. J. Allosteric receptors after
 Neuron 21, 950-980 (1998)
- years. Neuron 21, 959–980 (1998).
 Grosman, C., Zhou, M. & Auerbach, A. Mapping the conformational wave of acetylcholine receptor channel gating. Nature 403, 773–776 (2000).
- Monod, J., Wyman, J. & Changeux, J.-P. On the nature of allosteric transitions: a plausible model. *J. Mol. Biol.* 12, 88–118 (1965).
- Taverna, F. et al. The Lurcher mutation of an α-amino-3hydroxy-5-methyl-4-isoxazolepropionic acid receptor subunit enhances potency of glutamate and converts an antagonist to an agonist. J. Biol. Chem. 275, 8475–8479 (2000).
- Chapman, M. L., VanDongen, H. M. & VanDongen, A. M. Activation-dependent subconductance levels in the drk1 K channel suggest a subunit basis for ion permeation and gating. *Biophys. J.* 72, 708–719 (1997).
 Zheng, J. & Sigworth, F. J. Intermediate conductances
- Zheng, J. & Sigworth, F. J. Intermediate conductances during deactivation of heteromultimeric Shaker potassium channels. J. Gen. Physiol. 112, 457–474 (1998).
- channels. J. Gen. Physiol. 112, 457–474 (1998).
 Zheng, J. & Sigworth, F. J. Selectivity changes during activation of mutant Shaker potassium channels. J. Gen. Physiol. 110, 101–117 (1997).
- Luecke, H. & Quiocho, F. A. High specificity of a phosphate transport protein determined by hydrogen bonds. *Nature* 347, 402–406 (1990).
- Bräuner-Osborne, H., Egebjerg, J., Nielsen, E. O., Madsen, U. & Krogsgaard-Larsen, P. Ligands for glutamate receptors: design and therapeutic prospects. J. Med. Chem. 43, 2609–2645 (2000).
- Jayaraman, V., Keesey, R. & Madden, D. R. Ligand–protein interactions in the glutamate receptor. *Biochemistry* 39, 8693–8697 (2000).
- Madden, D. R., Thiran, S., Zimmermann, H., Romm, J. & Jayaraman, V. Stereochemistry of quinoxaline antagonist binding to a glutamate receptor investigated by Fourier transform infrared spectroscopy. J. Biol. Chem. 276, 37821–37826 (2001).
- Laube, B., Kuhse, J. & Betz, H. Evidence for a tetrameric structure of recombinant NMDA receptors. *J. Neurosci.* 18, 2954–2961 (1998).
- Mano, I. & Teichberg, V. I. A tetrameric subunit stoichiometry for a glutamate receptor-channel complex. Neuroreport 9, 327–331 (1998).
- 94. Safferling, M. et al. First images of a glutamate receptor ion channel: oligomeric state and molecular dimensions of GluRB homomers. Biochemistry 40, 13948–13953 (2001). This paper provides the first electron-microscopic views of an intact iGluR, showing it to be an elongated molecule of dimensions consistent with a dimer-ofdimers assembly.
- Partin, K. M. Domain interactions regulating AMPA receptor desensitization. J. Neurosci. 21, 1939–1948 (2001).
 Liu, D. T., Tibbs, G. R., Paoletti, P. & Siegelbaum, S. A.
- Liu, D. I., I Ibos, G. H., Paoletti, P. & Siegelbaum, S. A. Constraining ligand-binding site stoichiometry suggests that a cyclic nucleotide-gated channel is composed of two functional dimers. *Neuron* 21, 235–248 (1998).
- Perozo, E., Cortes, D. M. & Cuello, L. G. Structural rearrangements underlying K+-channel activation gating. Science 285, 73–78 (1999).
 - Electron paramagnetic resonance spectroscopic analysis allowed the identification of conformational
- changes associated with gating of a K* channel.
 Patneau, D. K. & Mayer, M. L. Kinetic analysis of interactions between kainate and AMPA: evidence for activation of a single receptor in mouse hippocampal neurons. *Neuron* 6, 785–798 (1991).
- Benveniste, M. & Mayer, M. L. Trapping of glutamate and glycine during open channel block of rat hippocampal neuron NMDA receptors by 9-aminoacridine. J. Physiol. (Lond.) 483, 367–384 (1995).

- 100. Lampinen, M., Pentikäinen, O., Johnson, M. S. & Keinänen, K. AMPA receptors and bacterial periplasmic amino acid-binding proteins share the ionic mechanism of ligand recognition. EMBO J. 17, 4704–4711 (1998).
- Morais-Cabral, J. H., Zhou, Y. & MacKinnon, R. Energetic optimization of ion conduction rate by the K* selectivity filter. Nature **414**, 37–42 (2001). 102. Lu, Z., Klem, A. M. & Ramu, Y. Ion conduction pore is
- conserved among potassium channels. Nature 413, 809-813 (2001).
- 103. Unwin, N. Acetylcholine receptor channel imaged in the open state. Nature 373, 37-43 (1995) Electron-microscopic analysis of different conformational states of the nAChR reveals a putative
- gating transition in the transmembrane domain M2. 104. Maeno, T., Edwards, C. & Anraku, M. Permeability of the endplate membrane activated by acetylcholine to some organic cations. *J. Neurobiol.* **8**, 173–184 (1977).
- 105. Dwyer, T. M., Adams, D. J. & Hille, B. The permeability of the endplate channel to organic cations in frog muscle. *J. Gen. Physiol.* **75**, 469–492 (1980).
- 106. Cohen, B. N., Labarca, C., Davidson, N. & Lester, H. A. Mutations in M2 alter the selectivity of the mouse nicotinic acetylcholine receptor for organic and alkali metal cations. J. Gen. Physiol. **100**, 373–400 (1992). 107. Burnashev, N., Villarroel, A. & Sakmann, B. Dimensions and
- ion selectivity of recombinant AMPA and kainate receptor channels and their dependence on Q/R site residues. J. Physiol. (Lond.) **496**, 165–173 (1996).
- Villaroel, A., Burnashev, N. & Sakmann, B. Dimensions of the narrow portion of a recombinant NMDA receptor
- channel. *Biophys. J.* **68**, 866–875 (1995). 109. Kuner, T., Beck, C., Sakmann, B. & Seeburg, P. H. Channellining residues of the AMPA receptor M2 segment: structural environment of the Q/R site and identification of the selectivity filter. *J. Neurosci.* **21**, 4162–4172 (2001).
- McCleskey, E. W. & Almers, W. The Ca channel in skeletal muscle is a large pore. Proc. Natl Acad. Sci. USA 82, 7149-7153 (1985).
- 111. Heginbotham, L., Abramson, T. & MacKinnon, R. A functional connection between the pores of distantly related ion channels as revealed by mutant K+ channels Science **258**, 1152–1155 (1992).
- 112. Heginbotham, L., Lu, Z., Abramson, T. & MacKinnon, R. Mutations in the K+ channel signature sequence. Biophys. J. **66**, 1061–1067 (1994).
- 113. Sommer, B. et al. Flip and flop: a cell-specific functional switch in glutamate-operated channels of the CNS. Science **249**, 1580–1585 (1990).
- 114. Hollmann, M., O'Shea-Greenfield, A., Rodgers, S. W. & Heinemann, S. Cloning by functional expression of a member of the glutamate receptor family. Nature 342, 643-648 (1989).
- Keinänen, K. et al. A family of AMPA-selective glutamate receptors. Science 249, 556–560 (1990).
- 116. Boulter, J. et al. Molecular cloning and functional expression of glutamate receptor subunit genes. Science **249**, 1033–1037 (1990).
- 117. Rose, C. R. & Konnerth, A. Self-regulating synapses. Nature 405, 413-415 (2000).
- 118. Bettler, B. et al. Cloning of a novel glutamate receptor subunit, GluR5: expression in the nervous system during development. *Neuron* **5**, 583–595 (1990).

- 119. Egebjerg, J., Bettler, B., Hermans-Borgmeyer, I. & Heinemann, S. Cloning of a cDNA for a glutamate receptor subunit activated by kainate but not AMPA. Nature 351,
- Werner, P., Voigt, M., Keinänen, K., Wisden, W. & Seeburg, P. H. Cloning of a putative high-affinity kainate receptor expressed predominantly in hippocampal CA3 cells. Nature **351**, 742–744 (1991).
- 121. Bettler, B. et al. Cloning of a putative glutamate receptor: a low affinity kainate-binding subunit. Neuron 8, 257-265 (1992).
- 122. Herb, A. et al. The KA-2 subunit of excitatory amino acid receptors shows widespread expression in brain and forms ion channels with distantly related subunits. *Neuron* **8**, 775–785 (1992).
- 123. Lerma, J., Paternain, A. V., Rodriguez-Moreno, A. & Lopez-Garcia, J. C. Molecular physiology of kainate receptors. *Physiol. Rev.* **81**, 971–998 (2001).
- 124. Moriyoshi, K. et al. Molecular cloning and characterization of the rat NMDA receptor. Nature **354**, 31–37 (1991). 125. Meguro, H. et al. Functional characterization of a
- heteromeric NMDA receptor channel expressed from cloned cDNAs. *Nature* **357**, 70–74 (1992).
- 126. Monyer, H. et al. Heteromeric NMDA receptors: molecular and functional distinction of subtypes. Science 256, 1217–1221 (1992).
- 127. Kutsuwada, T. et al. Molecular diversity of the NMDA receptor channel. Nature **358**, 36–41 (1992).
- 128. Ishii, T. et al. Molecular characterization of the family of the N-methyl-p-aspartate receptor subunits. J. Biol. Chem. 268 2836–2843 (1993).
- 129. Ciabarra, A. M. et al. Cloning and characterization of χ -1 a developmentally regulated member of a novel class of the ionotropic glutamate receptor family. J. Neurosci. 15, 6498-6508 (1995).
- 130. Sucher, N. J. et al. Developmental and regional expression pattern of a novel NMDA receptor-like subunit (NMDAR-L) in the rodent brain, J. Neurosci. 15, 6509-6520
- 131. Nishi, M., Hinds, H., Lu, H.-P., Kawata, M. & Hayashi, Y. Motoneuron-specific expression of NR3B, a novel NMDAtype glutamate receptor subunit that works in a dominantnegative manner. J. Neurosci. 21, RC185 (2001).
- 132. Johnson, J. W. & Ascher, P. Glycine potentiates the NMDA response in cultured mouse brain neurons. Nature 325, 529-531 (1987).
- 133. Mothet, J. P. et al. p-serine is an endogenous ligand for the glycine site of the N-methyl-p-aspartate receptor. Proc. Natl Acad. Sci. USA 97, 4926-4931 (2000).
- Nowak, L., Bregestovski, P., Ascher, P., Herbet, A. & Prochiantz, A. Magnesium gates glutamate-activated channels in mouse central neurones. Nature 307, 462-465 (1984)
- 135. Mayer, M. L., Westbrook, G. L. & Guthrie, P. B. Voltage-dependent block by Mg²⁺ of NMDA responses in spinal cord neurones. *Nature* 309, 261–263 (1984).
- 136. Wenthold, R. J., Yokotani, N., Doi, K. & Wada, K. Immunochemical characterization of the non-NMDA glutamate receptor using subunit-specific antibodies Evidence for a hetero-oligomeric structure in rat brain. J. Biol. Chem. **267**, 501–507 (1992).
- 137. Brose, N. et al. Differential assembly of coexpressed glutamate receptor subunits in neurons of rat

- cerebral cortex, J. Biol. Chem. 269, 16780-16784
- 138. Puchalski, R. B. et al. Selective RNA editing and subunit assembly of native glutamate receptors. Neuron 13,
- 131–147 (1994). 139. Seeburg, P. H. The TINS/TiPS lecture. The molecular biology of mammalian glutamate receptor channels. *Trends Neurosci.* **16**, 359–365 (1993).
- 140. Lomeli, H. et al. The rat δ -1 and δ -2 subunits extend the excitatory amino acid receptor family. FEBS Lett. **315**, 318–322 (1993).
- Kashiwabuchi, N. et al. Impairment of motor coordination, Purkinje cell synapse formation, and cerebellar long-term depression in GluR mutant mice. *Cell* **81**, 245–252 (1995).
- 142. Schuster, C. M. et al. Molecular cloning of an invertebrate glutamate receptor subunit expressed in Drosophila muscle. Science **254**, 112–114 (1991). 143. Maricq, A. V., Peckol, E., Driscoll, M. & Bargmann, C. I.
- Mechanosensory signalling in *C. elegans* mediated by the
- GLR-1 glutamate receptor. *Nature* **378**, 78–81 (1995). 144. Lam, H.-M. *et al.* Glutamate-receptor genes in plants.
- Nature **396**, 125–126 (1998). 145. Chiu, J., DeSalle, R., Lam, H. M., Meisel, L. & Coruzzi, G. Molecular evolution of glutamate receptors: a primitive signaling mechanism that existed before plants and animals diverged. *Mol. Biol. Evol.* **16**, 826–838 (1999).
- 146. Patneau, D. K. & Mayer, M. L. Structure-activity relationships for amino acid transmitter candidates acting at N-methyl -aspartate and quisqualate receptors. J. Neurosci. 10, 2385-2399 (1990).

Acknowledgement

This review is dedicated to the memory of Don C. Wiley, a great scientist, mentor and friend.

Online links

The following terms in this article are linked online to:

LocusLink: http://www.ncbi.nlm.nih.gov/LocusLink 81 | 82 | GluR1 | GluR2 | GluR3 | GluR4 | GluR5 | GluR6 | GluR7 | KA1 | KA2 | NR1 | NR2 | NR3

Protein Data Bank: http://www.rcsb.org/pdb/ 1BL8: potassium channel (KcsA) | 1EWK: mGluR subtype 1, complex with glutamate | 1FTJ: GluR2 S1S2J, complex with glutamate | 1FTK: GluR2 S1S2I, complex with kainate | 1FTL: GluR2 S1S2J, complex with DNQX | 1FTM: GluR2 S1S2J, complex with AMPA | 1FTO: GluR2 S1S2J, apo state | 1FW0: GluR2 S1S2J, complex with kainate | 1IIW: GluR0 ligand-binding core, apo state

FURTHER INFORMATION

Encyclopedia of Life Sciences: http://www.els.net/ AMPA receptors | ion channels | NMDA receptors Families of Transport Proteins:

http://www.biology.ucsd.edu/%7Emsaier/transport/toc.html Ion Channel Structure Research Group: http://www.mpimf-heidelberg.mpg.de/ion-channel/

Ligand-Gated Ion Channel Database

http://www.pasteur.fr/recherche/banques/LGIC/LGIC.html The Ion Channel Web Page:

http://phy025.lubb.ttuhsc.edu/Neely/ionchann.htm Access to this interactive links box is free online.