

Solving Linear Systems

August 2026

1 Introduction

Many problems in numerical analysis, engineering, and scientific computing can be expressed as a system of linear equations:

$$\begin{aligned}a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= b_1, \\a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= b_2, \\&\vdots \\a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n &= b_n.\end{aligned}$$

This system can be written compactly in matrix form as

$$A\mathbf{x} = \mathbf{b},$$

where

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}.$$

Here,

- A is the **coefficient matrix**,
- $\mathbf{x}$ is the **unknown vector**,
- $\mathbf{b}$ is the **right-hand-side vector**.

The objective is to determine $\mathbf{x}$ such that $A\mathbf{x} = \mathbf{b}$.

1.1 Direct and Iterative Methods

Direct methods obtain the solution through a finite sequence of arithmetic operations. In exact arithmetic, they produce the exact solution after a finite number of steps. Methods include Gaussian Elimination and LU Decomposition.

Iterative methods begin with an initial approximation $\mathbf{x}^{(0)}$ and repeatedly improve it

$$\mathbf{x}^{(0)} \rightarrow \mathbf{x}^{(1)} \rightarrow \mathbf{x}^{(2)} \rightarrow \cdots$$

until the approximation satisfies a chosen convergence criterion. Methods include Jacobi Method, Gauss-Seidel Method and Conjugate Gradient Method.

2 Gaussian Elimination

Gaussian Elimination is a **direct method** for solving a system of linear equations $A\mathbf{x} = \mathbf{b}$. The main idea is to transform the original system into an **upper triangular system**, which can then be solved using **back substitution**.

2.1 Forward Elimination

Consider the system

$$\begin{aligned}2x_1 + x_2 - x_3 &= 8, \\ -3x_1 - x_2 + 2x_3 &= -11, \\ -2x_1 + x_2 + 2x_3 &= -3.\end{aligned}$$

Its augmented matrix is

$$\left[\begin{array}{ccc|c} 2 & 1 & -1 & 8 \\ -3 & -1 & 2 & -11 \\ -2 & 1 & 2 & -3 \end{array} \right].$$

The goal of forward elimination is to eliminate the elements below the diagonal, transforming the matrix into the form

$$\left[\begin{array}{ccc|c} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \end{array} \right].$$

Pivot :

The **first pivot** is $a_{11} = 2$. To eliminate $a_{21} = -3$, calculate the multiplier

$$m_{21} = \frac{a_{21}}{a_{11}} = \frac{-3}{2}.$$

Then perform the row operation

$$R_2 - m_{21}R_1 \rightarrow R_2.$$

Similarly, to eliminate $a_{31} = -2$,

$$m_{31} = \frac{a_{31}}{a_{11}} = \frac{-2}{2} = -1,$$

and

$$R_3 - m_{31}R_1 \rightarrow R_3.$$

This produces

$$\left[\begin{array}{ccc|c} 2 & 1 & -1 & 8 \\ 0 & \frac{1}{2} & \frac{1}{2} & 1 \\ 0 & 2 & 1 & 5 \end{array} \right].$$

The **second pivot** is $a_{22} = \frac{1}{2}$. To eliminate $a_{32} = 2$, calculate

$$m_{32} = \frac{a_{32}}{a_{22}} = \frac{2}{1/2} = 4.$$

Then perform

$$R_3 - m_{32}R_2 \rightarrow R_3.$$

We obtain

$$\left[\begin{array}{ccc|c} 2 & 1 & -1 & 8 \\ 0 & \frac{1}{2} & \frac{1}{2} & 1 \\ 0 & 0 & -1 & 1 \end{array} \right].$$

The coefficient matrix is now in **upper triangular form**.

2.2 Back Substitution

The resulting system is

$$\begin{aligned}2x_1 + x_2 - x_3 &= 8, \\ \frac{1}{2}x_2 + \frac{1}{2}x_3 &= 1, \\ -x_3 &= 1.\end{aligned}$$

Starting from the last equation,

$$x_3 = -1.$$

Substitute $x_3 = -1$ into the second equation:

$$\frac{1}{2}x_2 + \frac{1}{2}(-1) = 1,$$

which gives

$$x_2 = 3.$$

Finally, substitute $x_2 = 3$ and $x_3 = -1$ into the first equation:

$$2x_1 + 3 - (-1) = 8,$$

so

$$x_1 = 2.$$

Therefore,

$$\mathbf{x} = \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix}.$$

2.3 Partial Pivoting

During forward elimination, the multiplier is calculated as

$$m_{ik} = \frac{a_{ik}}{a_{kk}},$$

where a_{kk} is the current **pivot**. If the pivot $a_{kk} = 0$, the multiplier cannot be calculated. Furthermore, if a_{kk} is very close to zero, division by a small number may amplify round-off errors and lead to numerical instability.

To avoid this problem, Gaussian Elimination is commonly implemented with **partial pivoting**. For each pivot column k , partial pivoting selects the element with the largest absolute value among the entries at or below the current pivot position:

$$|a_{pk}| = \max_{i=k,\dots,n} |a_{ik}|.$$

If $p \neq k$, rows k and p are swapped:

$$R_k \leftrightarrow R_p.$$

For example, consider

$$\left[\begin{array}{cc|c} 0.001 & 1 & 1 \\ 5 & 2 & 7 \\ -2 & 3 & 1 \end{array} \right].$$

In the first column,

$$|5| > |-2| > |0.001|,$$

so the second row is selected as the pivot row. After interchanging R_1 and R_2 ,

$$\left[\begin{array}{cc|c} 5 & 2 & 7 \\ 0.001 & 1 & 1 \\ -2 & 3 & 1 \end{array} \right].$$

The elimination can then proceed using 5 as the first pivot. Partial pivoting improves the numerical stability of Gaussian Elimination and avoids division by zero when a suitable nonzero pivot exists.

2.4 General Algorithm

For a general system

$$A\mathbf{x} = \mathbf{b},$$

Gaussian Elimination with partial pivoting consists of **forward elimination** followed by **back substitution**.

Forward Elimination

For each pivot column $k = 1, 2, \dots, n - 1$, perform the following steps.

1. Select the pivot row

Find the row p containing the largest absolute value in the current pivot column:

$$p = \operatorname{argmax}_{i=k, \dots, n} |a_{ik}|.$$

2. Check the pivot

If $|a_{pk}| < \varepsilon$, the matrix is singular or nearly singular, and the elimination cannot safely continue.

3. Swap rows

If $p \neq k$, interchange rows k and p : $R_k \leftrightarrow R_p$. The corresponding elements of the right-hand-side vector $\mathbf{b}$ must also be interchanged.

4. Eliminate the entries below the pivot

For $i = k + 1, \dots, n$, calculate the multiplier $m_{ik} = \frac{a_{ik}}{a_{kk}}$.

Then update the elements in row i :

$$a_{ij} - m_{ik}a_{kj} \rightarrow a_{ij}, \quad j = k, k + 1, \dots, n,$$

and update the corresponding element of the right-hand-side vector:

$$b_i - m_{ik}b_k \rightarrow b_i.$$

After forward elimination, the coefficient matrix is transformed into an upper triangular matrix.

*** Before performing back substitution, the final pivot must also be checked. If $|a_{nn}| < \varepsilon$, the matrix is singular or nearly singular.

Back Substitution

Starting from the last equation,

$$x_n = \frac{b_n}{a_{nn}}.$$

The remaining unknowns are calculated from bottom to top:

$$x_i = \frac{b_i - \sum_{j=i+1}^n a_{ij}x_j}{a_{ii}}, \quad i = n-1, n-2, \dots, 1.$$

3 LU Decomposition

Using Gaussian elimination, we eliminate the entries below the diagonal.

$$A = \begin{bmatrix} 3 & -1 & 2 \\ 6 & -1 & 5 \\ -9 & 7 & 3 \end{bmatrix}$$

Let $l_{21} = \frac{6}{3} = 2$, then we obtain

$$\begin{bmatrix} 3 & -1 & 2 \\ 0 & 1 & 1 \\ -9 & 7 & 3 \end{bmatrix}.$$

Next, let $l_{31} = \frac{-9}{3} = -3$, which gives

$$\begin{bmatrix} 3 & -1 & 2 \\ 0 & 1 & 1 \\ 0 & 4 & 9 \end{bmatrix}.$$

Finally, $l_{32} = \frac{4}{1} = 4$, so that

$$\begin{bmatrix} 3 & -1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 5 \end{bmatrix}.$$

The elimination multipliers are stored in the **lower triangular matrix**:

$$L = \begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -3 & 4 & 1 \end{bmatrix}.$$

Therefore,

$$\boxed{A = LU}$$

where

$$\boxed{\begin{bmatrix} 3 & -1 & 2 \\ 6 & -1 & 5 \\ -9 & 7 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -3 & 4 & 1 \end{bmatrix} \begin{bmatrix} 3 & -1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 5 \end{bmatrix}}$$

The entries of L are the elimination multipliers used during Gaussian elimination. The diagonal entries of L are all equal to 1 because each row retains its original leading coefficient during the elimination process.

The main advantage of LU decomposition is that it allows us to solve a linear system efficiently **when the same coefficient matrix is used with multiple right-hand sides**. Given $Ax = b$, and the decomposition $A = LU$, we obtain $LUx = b$ and $Ux = y$.

Then the system can be solved in two triangular systems: $Ly = b$, followed by $Ux = y$. The first system is solved by **forward substitution**, while the second is solved by **back substitution**.

3.1 LU Decomposition with Partial Pivoting

The previous example assumes that no row interchanges are required. However, in practical numerical computation, partial pivoting may be necessary for the same reasons discussed in Gaussian elimination. Consider

$$A = \begin{bmatrix} 0 & 2 & 1 \\ 1 & 1 & 2 \\ 2 & 1 & 1 \end{bmatrix}.$$

The first pivot is $a_{11} = 0$. Since the pivot is zero, we cannot perform the elimination directly. Using partial pivoting, we select the largest absolute value in the first column:

$$|2| > |1| > |0|.$$

Therefore, the third row is selected as the pivot row. We interchange R_1 and R_3 :

$$R_1 \leftrightarrow R_3,$$

giving

$$\begin{bmatrix} 2 & 1 & 1 \\ 1 & 1 & 2 \\ 0 & 2 & 1 \end{bmatrix}.$$

The first pivot is now $a_{11} = 2$. The elimination multipliers are

$$l_{21} = \frac{1}{2}, \quad l_{31} = \frac{0}{2} = 0.$$

Thus,

$$\begin{bmatrix} 2 & 1 & 1 \\ 0 & \frac{1}{2} & \frac{3}{2} \\ 0 & 2 & 1 \end{bmatrix}.$$

The second pivot is $a_{22} = \frac{1}{2}$. Therefore,

$$l_{32} = \frac{2}{1/2} = 4.$$

Performing the elimination, we obtain

$$U = \begin{bmatrix} 2 & 1 & 1 \\ 0 & \frac{1}{2} & \frac{3}{2} \\ 0 & 0 & -5 \end{bmatrix}.$$

The elimination multipliers are stored in

$$L = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & 4 & 1 \end{bmatrix}.$$

Because we interchanged the first and third rows, we also need a permutation matrix

$$P = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$$

The row interchange means that we decomposed PA , rather than A :

$$\boxed{PA = LU}$$

Indeed,

$$\boxed{\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 2 & 1 \\ 1 & 1 & 2 \\ 2 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & 4 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 0 & \frac{1}{2} & \frac{3}{2} \\ 0 & 0 & -5 \end{bmatrix}}$$

3.2 Solving Using LU Decomposition

Once the decomposition $PA = LU$ has been computed, solving $A\mathbf{x} = \mathbf{b}$ does not require performing Gaussian elimination again. Multiplying both sides by the permutation matrix gives $PA\mathbf{x} = P\mathbf{b}$. Since $PA = LU$, we obtain $LU\mathbf{x} = P\mathbf{b}$.

Rather than solving this system directly, we introduce an intermediate vector $\mathbf{y}$ and solve two triangular systems: $L\mathbf{y} = P\mathbf{b}$ followed by $U\mathbf{x} = \mathbf{y}$. The first system is solved by forward substitution, while the second is solved by back substitution.

Forward substitution :

Because L is lower triangular with ones on the diagonal, the unknowns are computed from top to bottom. For the system $L\mathbf{y} = P\mathbf{b}$, the equations are

$$\begin{aligned} y_1 &= (P\mathbf{b})_1, \\ y_2 &= (P\mathbf{b})_2 - l_{21}y_1, \\ y_3 &= (P\mathbf{b})_3 - l_{31}y_1 - l_{32}y_2, \\ &\vdots \\ y_n &= (P\mathbf{b})_n - \sum_{i=1}^{n-1} l_{ni}y_i. \end{aligned}$$

In general,

$$y_i = (P\mathbf{b})_i - \sum_{j=1}^{i-1} l_{ij}y_j, \quad i = 1, 2, \dots, n.$$

Since the diagonal entries of L are all equal to 1, no division is required.

Back substitution :

After obtaining y , we solve $U\mathbf{x} = \mathbf{y}$. Because U is upper triangular, the unknowns are computed from bottom to top. Starting from the last equation,

$$x_n = \frac{y_n}{u_{nn}}.$$

The remaining unknowns are calculated from bottom to top:

$$x_i = \frac{y_i - \sum_{j=i+1}^n u_{ij}x_j}{u_{ii}}, \quad i = n-1, n-2, \dots, 1.$$

Unlike forward substitution, division by the diagonal entries of U is necessary because they are generally not equal to one.

Regardless of coding, the lower-left entries store the multipliers of L , while the diagonal and upper-right entries store U . The diagonal of L is implicitly equal to 1. Thus, the matrix could be like

$$\begin{bmatrix} U & U & U \\ L & U & U \\ L & L & L \end{bmatrix}$$

So instead of allocating another $n \times n$ matrix, we overwrite those unused positions. This saves roughly half the memory while still allowing us to reconstruct both matrices whenever we need them.

4 Jacobi Method

The Jacobi Method, also known as the **Jacobi Iterative Method**, is a fundamental algorithm used to solve systems of linear equations $A\mathbf{x} = \mathbf{b}$.

Unlike Gaussian Elimination or LU Decomposition, which compute the solution through direct factorization, the Jacobi Method begins with an initial guess and repeatedly improves the approximation until it converges to the solution. Because each component of the new iterate is computed independently from the previous iterate, the method is easy to implement and naturally suitable for parallel computation.

4.1 Basic of Jacobi Method

For linear system $A\mathbf{x} = \mathbf{b}$, suppose the coefficient matrix is written as

$$A = D + L + U,$$

where D is the diagonal matrix containing the diagonal entries of A , L contains the entries below the diagonal, and U contains the entries above the diagonal.

$$D = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix}, \quad L = \begin{bmatrix} 0 & 0 & \dots & 0 \\ a_{21} & 0 & \dots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ a_{n1} & \dots & a_{n,n-1} & 0 \end{bmatrix}, \quad U = \begin{bmatrix} 0 & a_{12} & \dots & a_{1n} \\ 0 & 0 & \ddots & \vdots \\ \vdots & \ddots & \ddots & a_{n-1,n} \\ 0 & 0 & \dots & 0 \end{bmatrix}.$$

Substituting this decomposition into the original system gives

$$(D + L + U)\mathbf{x} = \mathbf{b}.$$

Rearranging,

$$D\mathbf{x} = \mathbf{b} - (L + U)\mathbf{x}.$$

Since D is diagonal, it is easy to invert, yielding the Jacobi iteration formula

$$\mathbf{x}^{(k+1)} = D^{-1}(\mathbf{b} - (L + U)\mathbf{x}^{(k)}).$$

4.2 Derive of the component-wise formula

To derive the component-wise iteration formula, consider the linear system

$$\begin{aligned} 10x_1 - x_2 + 2x_3 &= 6, \\ -2x_1 + 5x_2 + 2x_3 &= 6, \\ 2x_1 - x_2 + 10x_3 &= -10. \end{aligned}$$

The coefficient matrix is

$$A = \begin{bmatrix} 10 & -1 & 2 \\ -2 & 5 & 2 \\ 2 & -1 & 10 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 6 \\ 6 \\ -10 \end{bmatrix}.$$

Notice that the matrix is **strictly diagonally dominant**, so the Jacobi Method is expected to converge.

The coefficient matrix can be decomposed as $A = D + L + U$, where

$$D = \begin{bmatrix} 10 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 10 \end{bmatrix}, \quad L = \begin{bmatrix} 0 & 0 & 0 \\ -2 & 0 & 0 \\ 2 & -1 & 0 \end{bmatrix}, \quad U = \begin{bmatrix} 0 & -1 & 2 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{bmatrix}.$$

Based on the Jacobi iteration formula we can write

$$\begin{bmatrix} x_1^{(1)} \\ x_2^{(1)} \\ x_3^{(1)} \end{bmatrix} = \begin{bmatrix} 10 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 10 \end{bmatrix} \left(\begin{bmatrix} 6 \\ 6 \\ -10 \end{bmatrix} - \begin{bmatrix} 0 & -1 & 2 \\ -2 & 0 & 2 \\ 2 & -1 & 0 \end{bmatrix} \begin{bmatrix} x_1^{(0)} \\ x_2^{(0)} \\ x_3^{(0)} \end{bmatrix} \right).$$

Expanding the matrix equation gives

$$\begin{aligned} x_1^{(1)} &= \frac{1}{10}(6 - (-x_2^{(0)} + 2x_3^{(0)})), \\ x_2^{(1)} &= \frac{1}{5}(6 - (-2x_1^{(0)} + 2x_3^{(0)})), \\ x_3^{(1)} &= \frac{1}{10}(-10 - (2x_1^{(0)} - x_2^{(0)})). \end{aligned}$$

Thus, the Jacobi iteration is

$$\begin{aligned} x_1^{(k+1)} &= \frac{(6 + x_2^{(k)} - 2x_3^{(k)})}{10}, \\ x_2^{(k+1)} &= \frac{(6 + 2x_1^{(k)} - 2x_3^{(k)})}{5}, \\ x_3^{(k+1)} &= \frac{(-10 - 2x_1^{(k)} + x_2^{(k)})}{10}. \end{aligned}$$

Therefore, for a general $n \times n$ system, the Jacobi iteration is

$$\boxed{x_i^{(k+1)} = \frac{1}{a_{ii}}(b_i - \sum_{j \neq i} a_{ij}x_j^{(k)}).$$

4.3 Worked Example

From previous example we have

$$\begin{aligned} x_1^{(k+1)} &= \frac{(6 + x_2^{(k)} - 2x_3^{(k)})}{10}, \\ x_2^{(k+1)} &= \frac{(6 + 2x_1^{(k)} - 2x_3^{(k)})}{5}, \\ x_3^{(k+1)} &= \frac{(-10 - 2x_1^{(k)} + x_2^{(k)})}{10}. \end{aligned}$$

Choose the initial guess

$$\mathbf{x}^{(0)} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

Iteration 1 : Using $\mathbf{x}^{(0)}$

$$\begin{aligned} x_1^{(1)} &= \frac{(6 + 0 - 2(0))}{10} = 0.600 \\ x_2^{(1)} &= \frac{(6 + 0 - 0)}{5} = 1.200 \\ x_3^{(1)} &= \frac{(-10 - 2(0) + 0)}{10} = -1.000 \end{aligned} \quad \rightarrow \mathbf{x}^{(1)} = \begin{bmatrix} 0.6000 \\ 1.2000 \\ -1.0000 \end{bmatrix}$$

Iteration 2 : Using $\mathbf{x}^{(1)}$

$$\begin{aligned} x_1^{(2)} &= \frac{(6 + 1.2000 - 2(-1.0000))}{10} = 0.9200 \\ x_2^{(2)} &= \frac{(6 + 2(0.6000) - 2(-1.0000))}{5} = 1.8400 \\ x_3^{(2)} &= \frac{(-10 - 2(0.6000) + 1.2000)}{10} = -1.0000 \end{aligned} \quad \rightarrow \mathbf{x}^{(2)} = \begin{bmatrix} 0.9200 \\ 1.8400 \\ -1.0000 \end{bmatrix}$$

The approximations gradually approach the exact solution

$$\mathbf{x} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}.$$

4.4 Convergence of the Jacobi Method

Recall the matrix splitting $A = D + L + U$, the Jacobi iteration is

$$\mathbf{x}^{(k+1)} = D^{-1}\mathbf{b} - D^{-1}(L + U)\mathbf{x}^{(k)}.$$

Define the iteration matrix $T = -D^{-1}(L + U)$. Then the iteration becomes

$$\mathbf{x}^{k+1} = D^{-1}\mathbf{b} + T\mathbf{x}^{(k)}.$$

The behavior of the method is completely determined by the matrix T .

The Jacobi Method converges if the spectral radius of the iteration matrix satisfies $\rho(T) < 1$. Here, $\rho(T) = \max_i |\lambda_i|$ is the largest absolute value among all eigenvalues of T .

The Jacobi iteration can be written as $\mathbf{x}^{k+1} = \mathbf{c} + T\mathbf{x}^{(k)}$, where T is the iteration matrix, $\mathbf{c}$ is a constant vector, and $\mathbf{x}^{(k)}$ is the approximation after the k-th iteration.

Suppose the exact solution is $\mathbf{x}^*$. Since it satisfies the same iteration equation

$$\mathbf{x}^* = \mathbf{c} + T\mathbf{x}^*.$$

Define the error at the k-th iteration as

$$\mathbf{e}^{(k)} = \mathbf{x}^{(k)} - \mathbf{x}^*.$$

Subtracting the two equations eliminates the constant vector $\mathbf{c}$, giving

$$\begin{aligned} \mathbf{e}^{(k+1)} &= \mathbf{x}^{(k+1)} - \mathbf{x}^* \\ &= (\mathbf{c} + T\mathbf{x}^{(k)}) - (\mathbf{c} + T\mathbf{x}^*) \\ &= T(\mathbf{x}^{(k)} - \mathbf{x}^*) \\ &= T\mathbf{e}^{(k)} \end{aligned}$$

This recurrence shows that the error is transformed by the iteration matrix at each step. Applying the recurrence repeatedly gives

$$\begin{aligned}\mathbf{e}^{(1)} &= T\mathbf{e}^{(0)} \\ \mathbf{e}^{(2)} &= T\mathbf{e}^{(1)} = T^2\mathbf{e}^{(0)} \\ &\vdots \\ \mathbf{e}^{(k)} &= T^{(k-1)}\mathbf{e}^{(k-1)} = T^k\mathbf{e}^{(0)}\end{aligned}$$

Assume that the iteration matrix T has n linearly independent eigenvectors

$$\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n,$$

with corresponding eigenvalues

$$\lambda_1, \lambda_2, \dots, \lambda_n.$$

Since these eigenvectors form a basis of the vector space, any initial error can be written as a linear combination of them

$$\mathbf{e}^{(0)} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_n\mathbf{v}_n,$$

where $c_1, c_2, \dots, c_n$ are constants.

By the definition of an eigenvalue $T\mathbf{v}_i = \lambda\mathbf{v}_i$. Applying the matrix repeatedly gives

$$T^k\mathbf{v}_i = \lambda^k\mathbf{v}_i.$$

Substituting this into the error formula,

$$\begin{aligned}\mathbf{e}^{(k)} &= T^k\mathbf{e}^{(0)} \\ &= c_1(T^k\mathbf{v}_1) + c_2(T^k\mathbf{v}_2) + \dots + c_n(T^k\mathbf{v}_n) \\ &= c_1\lambda^k\mathbf{v}_1 + c_2\lambda^k\mathbf{v}_2 + \dots + c_n\lambda^k\mathbf{v}_n.\end{aligned}$$

Now consider the limit as $k \rightarrow \infty$. If every eigenvalue satisfies

$$|\lambda_i| < 1, \quad i = 1, 2, \dots, n.$$

then each term λ_i^k tends to zero $\lim_{k \rightarrow \infty} \lambda_i^k = 0$.

Consequently,

$$\lim_{k \rightarrow \infty} \mathbf{e}^{(k)} = \mathbf{0}.$$

The approximate solution converges to the exact solution $\lim_{k \rightarrow \infty} x^{(k)} = x^*$. Therefore, the Jacobi Method converges whenever all eigenvalues of the iteration matrix lie inside the unit circle, which is equivalent to the condition

$$\rho(T) = \max_i |\lambda_i| < 1.$$

4.5 Strictly diagonally dominant matrix

Let A be a strictly diagonally dominant matrix, meaning that

$$|a_{ii}| > \sum_{j \neq i} |a_{ij}|, \quad i = 1, 2, \dots, n.$$

Let λ be an arbitrary eigenvalue of T , with corresponding eigenvector $\mathbf{v} \neq 0$. By the definition of an eigenvalue, $T\mathbf{v} = \lambda\mathbf{v}$. Substituting $T = -D^{-1}(L + U)$ gives

$$-D^{-1}(L + U)\mathbf{v} = \lambda\mathbf{v}.$$

Multiplying both sides by D yields

$$-(L + U)\mathbf{v} = \lambda D\mathbf{v}.$$

For the i -th row, this becomes

$$-\sum_{j \neq i} a_{ij}v_j = \lambda a_{ii}v_i.$$

Now choose the component of v with the largest magnitude. Suppose it is the m -th component, so that

$$|v_m| = \max_{1 \leq j \leq n} |v_j| > 0.$$

Applying the previous equation to the m -th row,

$$\lambda a_{mm}v_m = -\sum_{j \neq m} a_{mj}v_j$$

Taking absolute values and applying the triangle inequality,

$$|\lambda||a_{mm}||v_m| = \left| \sum_{j \neq m} a_{mj}v_j \right| \leq \sum_{j \neq m} |a_{mj}||v_j|.$$

Since v_m is the largest component in magnitude, therefore

$$|\lambda||a_{mm}||v_m| \leq \sum_{j \neq m} |a_{mj}||v_m|.$$

Because $|v_m| > 0$, dividing both sides by $|v_m|$ gives

$$|\lambda||a_{mm}| \leq \sum_{j \neq m} |a_{mj}|.$$

or equivalently,

$$|\lambda| \leq \frac{\sum_{j \neq m} |a_{mj}|}{|a_{mm}|}.$$

Finally, since A is strictly diagonally dominant, $|a_{mm}| > \sum_{j \neq m} |a_{mj}|$, which implies

$$|\lambda| \leq \frac{\sum_{j \neq m} |a_{mj}|}{|a_{mm}|} < 1.$$

Since λ was chosen as an arbitrary eigenvalue of T , every eigenvalue satisfies $|\lambda| < 1$, which means $\rho(T) < 1$. Therefore, the Jacobi Method converges for any initial guess when A is strictly diagonally dominant.