

1
2
3
4
5
6
7

Native vs. Novel Threats

**Fear of the new: a predatory invader provokes stronger non-consumptive effects
on a keystone native prey than a native predator**

Olivier V. Raven Calum MacNeil Finnbar Lee
Robin Holmes Ian A. K. Kusabs Francis J. Burdon
Deniz Özkundakci

2026-07-22

Native vs. Novel Threats

Fear of the new: a predatory invader provokes stronger non-consumptive effects on a keystone native prey than a native predator



Olivier V. Raven, Calum MacNeil, Finnbar Lee, Robin Holmes, Ian A. K. Kusabs, Francis J. Burdon & Deniz Özkundakci

Abstract

The non-consumptive effects (NCEs) of predators on prey result in costly antipredator behaviours, including increased refuge use and reduced foraging. NCEs significantly influence prey population dynamics, community interactions, and affect ecosystem functioning, particularly when the prey are keystone species. In Aotearoa-New Zealand, the native freshwater crayfish kōura (*Paranephrops planifrons*) is a keystone species with an important functional role affecting sediment bioturbation, nutrient cycling, and benthic community structure. Kōura are preyed upon by multiple predators including native longfin eel (*Anguilla dieffenbachii*) and invasive brown bullhead catfish (*Ameiurus nebulosus*). Under laboratory conditions, we tested whether visual and chemical cues from these two predators induced different refuge-seeking and feeding behaviour responses by kōura. Predator presence produced in the dark a NCE on refuge use, with kōura exposed to catfish showing significantly greater refuge-seeking behaviour than those exposed to eels. This result is consistent with the generalisation theory triggered by unfamiliar predator cues rather than classic prey naivety. Our study indicates catfish incursions may provoke a greater NCE on kōura behaviours than resident eels. As catfish range expansion continues, these NCEs may represent an ecologically significant and underappreciated threat to kōura populations, compounding the direct predation pressure already documented for this invader.

Introduction

Biological invasions can have profound impacts on invaded ecosystems, particularly through the introduction of novel predators that prey upon naïve native species lacking coevolved defence mechanisms (Doherty et al. 2016). Native prey may be naïve not just to a specific invader, but to entire functional groups of predators, making them especially vulnerable to direct predation when an invasion introduces a predator type entirely absent from their evolutionary history (Sih et al. 2010; Carthey and Banks 2014). In addition to these consumptive effects, native prey may exhibit heightened fear responses to invasive predators, leading to the development of anti-predator behaviours such as increased refuge use and avoidance of open habitats (Kindinger and Albins 2017). These behavioural changes, termed non-consumptive effects (NCEs), can substantially influence population dynamics by reducing foraging activity, growth, and reproductive output (Preisser et al. 2005). In contrast, native predators generally elicit different responses because prey and predators have coevolved, resulting in behavioural and physiological adaptations that stabilise predator–prey interactions (Abrams 2000).

1 Chemical signals (kairomones) are a key component of NCEs (Kasumyan 2022). Freshwater
2 crayfish release and detect kairomones to help navigate aquatic environments and mediate
3 interactions with predatory fish (Blake and Hart 1993; Brönmark and Hansson 2012; Wood
4 and Moore 2020a). Kairomones carry signals via excreta, mucus, or skin, that crayfish use
5 to detect predators (Appelberg et al. 1993). Crayfish respond more strongly to signals from
6 predators that have consumed conspecifics and to alarm cues released when conspecifics are
7 injured or killed (Gherardi et al. 2011; Beattie and Moore 2018). The NCEs due to kairomone
8 detection include reduced growth, reduced activity levels (Adams 2007; Ramberg-Pihl and
9 Yurewicz 2020), and altered social dynamics like the use of refuge spaces (Driscoll et al. 2020).
10 Non-predatory fish can also trigger behaviour changes, likely through an increased awareness
11 of potential risk (Nyström 2005). However, the strength of these responses depends on prior
12 experience as wild-caught individual crayfish show stronger antipredator responses than farmed
13 conspecifics naïve to predators (Martin 2014), indicating that both learned behavior and
14 evolutionary history shape kairomone detection.

15 In Aotearoa-New Zealand (hereafter Aotearoa), the endemic northern freshwater crayfish
16 *Paranephrops planifrons* (White, 1842) is a keystone species impacted by non-native species.
17 This crayfish species is commonly known as kōura and is a taonga (culturally treasured) species
18 for Māori, the indigenous people of Aotearoa, with strong cultural values as a food resource
19 (Kusabs and Quinn 2009). Ecologically, kōura play a critical role as ecosystem engineers through
20 their effects on sediment dynamics, nutrient cycling, and benthic community structure (Collier
21 et al. 1997; Parkyn et al. 2001; Usio and Townsend 2004). Currently, kōura are threatened by
22 multiple stressors, including climate change, water-quality degradation and invasive species,
23 contributing to widespread population declines (Lee et al. 2025; Kusabs et al. 2026).

24 Kōura have a long coevolutionary history with the native longfin eel (*Anguilla dieffenbachii*)
25 (hereafter eel) and shortfin eel (*A. australis*) in Aotearoa (Brown 2009; MacNeil et al. 2025).
26 Reflecting this coevolutionary history, kōura can respond more strongly to eel kairomones than
27 to those of introduced non-native brown trout (*Salmo trutta*), even when sourced from streams
28 where both predators were present (Shave et al. 1994). Shave et al. (1994) suggest that kōura
29 predator recognition is a fixed behavioural trait shaped by coevolution. An invasive fish species
30 that threatens kōura is the brown bullhead catfish (*Ameiurus nebulosus*) (hereafter catfish).
31 Catfish were introduced to Aotearoa in 1877, and the invasive species has since spread to
32 multiple lakes and rivers in the North Island (McDowall 1990; Barnes and Hicks 2003; Dedual
33 2019). For example, in 2016/18, the presence of catfish was confirmed in two large volcanic
34 lakes, Lakes Rotoiti and Rotorua in the Bay of Plenty region, which have historically contained

1 large populations of kōura (Francis 2019; Lee et al. 2025). Catfish are highly tolerant of a wide
2 range of environmental conditions (Scott and Crossman 1973; Lee et al. 2025) and their broad
3 gape and opportunistic feeding behaviour enable the consumption of a wide variety of aquatic
4 prey (Scott and Crossman 1973; Dedual 2019). Given Aotearoa lacks native Siluriformes, the
5 recent and ongoing spread of catfish into North Island waterways represents a truly novel
6 predation threat to which kōura may lack adequate behavioural or sensory adaptations to
7 respond.

8 In this study, we used controlled laboratory experiments to test whether the presence of a novel
9 and native predator induced differences in kōura NCEs, specifically refuge-seeking and feeding
10 behaviours. Our experiments tested the following two hypotheses regarding kōura behavioural
11 responses to predator cues: 1) Kōura would show less shelter-seeking behaviour when presented
12 with catfish compared to eels, due to naivety towards a novel predator (Carthey and Banks
13 2014); or 2) kōura would respond more strongly to unfamiliar catfish cues than to the native
14 eel, consistent with a generalised fear response to a novel predator type (Sih et al. 2023).
15 Further, we expected that light regime would have an influence on shelter seeking behaviour,
16 as kōura, like many crayfish, are primarily nocturnal (Devcich 1979; Johnson et al. 2014). We
17 also predicted that kōura feeding rates would follow the patterns hypothesized above and either
18 be more in the presence of catfish than eels, reflecting stronger antipredator responses to the
19 native predator, or reduced in the presence of the unfamiliar invasive predator, reflecting a
20 generalised fear response to the novel predator. Overall, the refuge-use and feeding experiment
21 aimed to improve understanding of the behavioural mechanisms underlying kōura vulnerability
22 to invasive catfish, and how these responses may differ, and add to, those displayed in the
23 presence of native predators such as eels.

24 **Materials and Methods**

25 **Animal collection and preparation**

26 In mid-August 2025, 92 kōura were sourced from two catfish-free lakes, Ōkāreka and Tikitapu,
27 within the Rotorua Te Arawa Lakes area in the North Island of Aotearoa. Eels are rare or
28 absent in these lakes (Martin et al. 2007), potentially limiting the recent coevolutionary history
29 with local kōura populations. However, eels are a native predator that has a long record of
30 coexistence with kōura in Aotearoa (McDowall 1990; Feldmann and Pole 1994; Kaulfuss et
31 al. 2018). In this study our focus was on the macroevolutionary implications of the native

1 predator compared to a novel invader, regardless of the local coevolutionary history. In late
2 September 2025, catfish and eels were sourced from Lake Rotoroa in Hamilton in the Waikato
3 region of the North Island. Experiments were conducted at the Aquatic Research Centre
4 of the University of Waikato. All three species were held separately in holding tanks and
5 acclimated for a week in the laboratory conditions. These included a 10:14 light:dark cycle
6 reflecting the seasonal day-length and dechlorinated tap water with stable temperatures (15.6
7 °C), pH (8.4) and dissolved oxygen concentrations (10 mg l⁻¹). These parameters reflect typical
8 physiochemical conditions in the source lakes. Kōura and fish were fed every other day with
9 commercial carnivore pellets (Hikari Tropical, Kyorin Food Industries Ltd., Japan; animal
10 protein 47 %). Because kōura are a taonga (treasured) species, permission was obtained from Te
11 Arawa Lakes Trust and Te Komiti Whakahaere. All experimental procedures were conducted
12 under University of Waikato animal ethics committee approval (protocol number: 1232; see
13 Ethical Approval section).

14 **Refuge experiment**

15 Prior to the experiments all animals were individually weighed and measured. Only male kōura
16 were used as male crustaceans are usually the most bold and aggressive (Stein and Magnuson
17 1976; Figler et al. 2005) and restricting test animals to a single sex allowed a smaller overall
18 number of kōura to be used. Kōura had a mean ($\pm$ SD) orbital carapace length (OCL) of 26.0
19 mm ($\pm$ 3.08) (18.35-32.95 mm range) and a mean ($\pm$ SD) wet weight of 11.7 g ($\pm$ 4.25) (3–23
20 g range). Catfish ranged in length from 260-267 mm and 2.34–3.16 kg wet weight and eels
21 ranged in wet weight from 2.50-3.52 kg, with both these size ranges being able to consume
22 adult kōura (Hicks 1997; Barnes and Hicks 2003; Ryan 2009).

23 Experiments were conducted in 38 L glass tanks (600 $\times$ 200 $\times$ 220 mm) with each tank having
24 three sides (apart from front of tank) blacked out to block visual cues from neighbouring tanks.
25 Each tank was divided into two compartments by a 0.5 mm steel mesh divider, allowing visual
26 and chemical cues between compartments, while preventing any physical contact between fish
27 and kōura on either side of the barrier (Figure 1). All tanks were filled with ~26 L dechlorinated
28 tap water and had identical environmental regimes as previously detailed for acclimatisation
29 tanks, with water flow provided by an air stone. Each tank housed a single kōura on one side of
30 the barrier and either a single predator (catfish or eel) or no predator (control) on the other side.
31 A halved terracotta pot (90 mm diameter $\times$ 90 mm depth) was positioned on the kōura side of
32 the tank to serve as a potential refuge. To control for potential tank location or compartment

side effects, the position of predator treatments within the aquaria was systematically rotated between experimental rounds. The experiment was conducted across eight rounds (each lasting 5 hours), four rounds under light conditions (~400 lux, n = 32 kōura: 11 catfish, 11 eel, 10 control) and four separate rounds with different kōura under dark conditions (~70 lux, n = 32 kōura: 11 catfish, 11 eel, 10 control). Observations were made in the dark using red lighting positioned centrally above the tanks to minimize disturbance, as many aquatic species have limited sensitivity to red wavelengths and are unlikely to perceive this part of the spectrum. Following each individual experimental run, each tank was thoroughly cleaned and refilled with dechlorinated tap water before reuse. Each individual kōura was only exposed once to a fish predator and there were no mortalities of kōura or fish during experiments. All experiments were conducted within a three-week period during October 2025.

At the beginning of each experiment, an individual kōura was placed in the middle of the tank compartment containing the refuge. Kōura location was recorded at 21-time intervals over a 5-hour period, with observations spaced at progressively longer intervals to capture immediate behavioural responses at key timepoints (predator introduction at 60 min, removal at 240 min). The 0-minute recording was the individual's initial immediate location after its addition to the tank. Kōura location was recorded as either refuge (the animal was using the refuge, being inside it or immediately behind it and touching it) or non-refuge/ in the open (any other location in the tank where the animal was not in the refuge or touching the refuge). The control tanks had no fish added but at 60 minutes and again at 240 minutes, the water in the 'predator' compartment was mechanically disturbed by agitating the water with a small hand net for approximately 30 seconds. This was to simulate the disturbance time associated with adding and removing fish in the other treatments with the same size hand net. At the end of each experiment, the kōura was removed after the final 300-minute observation.

Feeding experiment

Experiments were conducted in glass tanks (320 × 170 × 170 mm), containing 5 L dechlorinated tap water and a halved terracotta pot (90 mm diameter × 90 mm depth). Catfish and eels had been held at the same density in separate identical holding tanks for a week prior to experiment. Fish exudates were collected by placing four fish (catfish or eel) in a 10 L bucket filled with water from the holding tanks for 15 minutes to concentrate the kairomones in the water. For eel and catfish treatments, 1 L from the buckets was added to each tank already holding 5 L of tap water, and for control an additional 1 L of dechlorinated tap water was added. Next, 10

1 Hikari Tropical carnivore pellets were added to each tank. Experimental trials showed pellets
2 remain intact in tap water for over 24 hours. In three rounds, a total of 46 male kōura were
3 measured and weighed. Their mean ($\pm$ SD) OCL was 27.93 mm ($\pm$ 4.80), with a range of 20.48
4 to 40.88 mm, and the mean ($\pm$ SD) wet weight of 15.83 g ($\pm$ 8.99), ranging from 6.00 to 42.00
5 g. Each kōura was then placed individually in a tank. To investigate how starvation period
6 influenced feeding motivation, each round kōura were starved for different periods (either 0, 3
7 or 4 days) prior to the feeding experiment. After 0 days starvation, 14 kōura were exposed to
8 eel water ($n = 4$), catfish water ($n = 6$) or control water ($n = 4$). After 3 days starvation, 16
9 kōura were exposed to eel water ($n = 6$), catfish water ($n = 5$) or control water ($n = 5$). After
10 4 days starvation, 16 kōura were exposed to eel water ($n = 5$), catfish water ($n = 5$) or control
11 water ($n = 6$). After 24 hours, when each experiment was terminated, the number of pellets
12 eaten (to the nearest quarter pellet) was counted and the feeding rate (pellets eaten per gram
13 of wet weight (WW) of kōura) calculated.

14 Data analysis

15 To analyse kōura refuge use, beta-binomial generalized linear mixed-effects models (GLMMs)
16 were fitted using the glmmTMB package in R ([Brooks et al. 2017](#); [R Core Team 2025](#)). To
17 address potential non-independence of repeated observations within tanks, raw observations
18 were first aggregated into three experimental periods (before, during, and after predator
19 introduction) per individual kōura, with the number of observations in refuge out of total
20 observations per period used as the response variable. Treatment (catfish, eel, control) and
21 period (before, during, after) were included as fixed effects, and tank and experimental round
22 as random intercepts. Because light condition and experimental round were fully confounded
23 by design (rounds 1–4 in light, rounds 5–8 in darkness), separate analyses were conducted
24 for each light condition to allow round as a random intercept and to test whether treatment
25 effects varied under each light condition. A beta-binomial distribution was used to account
26 for overdispersion in the binomial proportion data. Model fit was assessed using DHARMA
27 simulation-based residuals, including tests for overdispersion and zero-inflation ([Hartig et al.
28 2024](#)). Marginal means and pairwise contrasts between factor levels were obtained using the
29 emmeans package ([Lenth et al. 2026](#)), with p-values adjusted for multiple comparisons using
30 Sidak correction. Additionally, a combined GLMM (without round as random effect) was fitted
31 to test the overall effect of light condition across all treatments. Spearman rank correlations
32 with Bonferroni-adjusted p-values were used to investigate potential relationships between kōura
33 body size (OCL) and overall refuge use (all time intervals combined) in the three treatments.

1 Kōura feeding rate was analysed using a linear mixed model with pellets consumed per gram
2 wet weight as the response variable and starvation period (0, 3, 4) as a fixed effect, with
3 tank identity as a random intercept to account for potential tank-level clustering. Because
4 each starvation period was conducted in a separate experimental round, starvation period and
5 experimental round were fully confounded by design. Consequently, round was excluded as a
6 random effect. However, starvation period was included as a fixed effect to test for differences
7 in feeding across hunger states, acknowledging that observed effects may reflect round-specific
8 variation rather than hunger alone. Model selection using AICc indicated that starvation
9 period alone provided the best fit, with treatment not improving model fit (delta AICc =
10 14.38). Therefore, treatment effects were excluded from the main model and instead examined
11 separately within each starvation period using stratified linear models to avoid confounding
12 treatment comparisons with the starvation-round confounding. Model assumptions were verified
13 using residual diagnostics (Q-Q plots and residual plots) and testing normality of residuals
14 using the Shapiro-Wilk test. As residuals at day 0 starvation violated normality assumptions
15 ($p = 0.011$), a non-parametric Kruskal-Wallis test was executed. Pairwise comparisons for
16 starvation period were conducted using estimated marginal means with Sidak adjustment for
17 multiple comparisons.

18 Results

19 Refuge experiment

20 The proportion of time kōura spent in refuge varied across treatments, experimental periods,
21 and light conditions (Figure 2). Stratified analyses revealed that treatment effects on refuge use
22 differed between light and dark conditions. Under dark condition, kōura exposed to catfish used
23 refuge significantly more than those exposed to eels ($p = 0.004$; Table 1), however the control
24 treatment in the dark was not significant with either the catfish or the eel treatment ($p = 0.083$,
25 $p = 0.565$). In contrast, under light conditions, there were no significant differences in refuge
26 use between the three treatments, but refuge use differed significantly between experimental
27 periods, with lower refuge use before predator introduction compared to after predator removal
28 ($p = 0.021$). No significant period effects were observed in dark conditions. The combined
29 model showed that light regime had a significant effect on refuge use ($p = 0.005$), with kōura
30 using refuge more in light than in dark conditions across all treatments. Overall, under light
31 conditions, the pattern of refuge use in the control treatment was more similar to the catfish
32 treatment, whereas under dark conditions it more resembled the eel treatment (Fig. 3; Table

1 S1). There was also no relationship between kōura OCL and refuge use in any of the three
2 treatments ($r_s = -0.09$, $p = 0.693$, $r_s = 0.09$, $p = 0.680$ and $r_s = 0.16$, $p = 0.494$ for catfish, eel
3 and control treatments respectively, all NS).

4 For all treatments individual kōura were highly variable in their movement around tanks and
5 use of refuges, even in the time intervals comprising the first 60 minutes of the experiment,
6 before fish were introduced. The coefficient of variations (CVs) for kōura refuge use in the light
7 and the dark before fish introduction were similarly high, being 60.4 % and 58.2 % (both $n =$
8 32) respectively, confirming high baseline variation in kōura refuge use, regardless of treatment
9 type. Despite this inherent variation in individual kōura behaviour, clear patterns in kōura
10 refuge use were evident over the course of the experiment (Figure 3). Under light conditions,
11 the control showed the largest overall increase in refuge use (40% to 82%), driven primarily
12 by a sharp rise in the ‘after’ period, while the catfish treatment showed a more sustained
13 increase across all periods (52% to 73%) and the eel treatment the smallest increase (35%
14 to 53%). Under dark conditions, the pattern shifted, with the catfish treatment showing the
15 greatest increase (39% to 70%), while the eel treatment showed a modest and non-sustained
16 increase (27% to 41% during ‘exposure’, returning to 36% ‘after’), and the control remained
17 comparatively low throughout (37% to 47%).

18 Feeding experiment

19 The linear mixed model showed that starvation period had a significant effect on kōura feeding
20 rate (3 days: $p < 0.001$; 4 days: $p = 0.046$; Figure 4; **?@tbl-feeding-summary**). Treatment
21 was not included in the main model following model selection. Stratified analyses revealed no
22 significant treatment effects across starvation periods. Although residuals at day 0 violated
23 normality (Shapiro-Wilk $p = 0.0112$), the non-parametric Kruskal-Wallis test confirmed no
24 treatment effect ($p = 0.3165$). Starvation days 3 and 4 met normality assumptions and showed
25 no significant treatment effects ($p = 0.1495$ and 0.1573 respectively). After 4 days starvation,
26 the mean ($\pm$ SE) kōura feeding rate (pellets g WW^{-1}) for the control was over two and a half
27 times that of the catfish and eel treatments (0.148 ± 0.035 compared to 0.059 ± 0.039 and
28 0.052 ± 0.039 respectively).

Discussion

Prey modify their behaviour in response to predation risk, creating a ‘landscape of fear’ that affects foraging, movement, and habitat use independently of actual predation events (Laundré et al. 2001). Our study adds to a growing body of literature (i.e., (Sniegula et al. 2025)) showing invasive predators can generate stronger NCEs on native prey than native predators, with potentially far-reaching consequences on invaded ecosystems. The results in our experiments suggest the introduction of a novel predator may change the prey’s fear landscape in ways that a native predator does not. Kōura are a keystone species that act as ecosystem engineers in Aotearoa freshwaters, meaning that predator-induced changes in kōura behaviour may indirectly affect ecosystem-level properties (e.g., resources or other species) through non-consumptive pathways (Peacor and Werner 2001). Such indirect effects represent a form of behavioural trophic cascade (Schmitz et al. 1997), whereby predator presence alone drives ecosystem-level change. Our results also highlight that prey naivety does not necessarily manifest as a failure to respond to novel predators. Instead, naïve prey may exhibit an over-avoiding response consistent with generalisation theory (Sih et al. 2010; Ehlman et al. 2019), which potentially has important implications for predicting the ecological impacts of invasive species.

Refuge use

Under dark conditions, kōura demonstrated greater refuge-seeking behaviour when exposed to the novel catfish than to native eel. This result is notable because it contradicts the classic prey naivety hypothesis, which predicts that prey underestimate or fail to appropriately respond to novel predators due to a lack of shared evolutionary history (Carthey and Banks 2014). Instead, kōura seemingly exhibited an enhanced defensive response to the novel predator. Such over-responses to unfamiliar predators are less commonly reported but are consistent with fear generalisation theory, whereby experience with native predators primes a generalised heightened vigilance response to any unfamiliar threat (Ehlman et al. 2019; Sih et al. 2023). For naïve prey, unfamiliar cues could trigger such a response because the consequences of ignoring a potential predation threat could be fatal, even when, in reality, the actual predation risk is absent (Dawkins and Krebs 1979; Lima and Dill 1990). Such generalised responses are often overcautious and energetically costly because prey may initially treat uncertain threats as highly dangerous until further experience is acquired (Sih et al. 2010). Unlike classic prey naivety scenarios where the novel predator poses no actual threat, catfish genuinely predate kōura, meaning the heightened refuge response carries a real energetic cost but is not disproportionate

1 to the actual risk of predation. Elevated refuge use is itself a meaningful NCE, reducing
2 foraging time with potential consequences for growth, reproduction, and population persistence
3 ([Preisser et al. 2005](#); [Sheriff et al. 2020](#)). However, this behaviour may simultaneously confer
4 protection against direct predation as experimental evidence shows that shelter significantly
5 reduces catfish predation on crayfish ([Adams 2007](#); [Zhao et al. 2024](#)). This result suggests that
6 enhancing or restoring habitat complexity in catfish-invaded lakes could mitigate both direct
7 predation and fear-driven reductions in foraging, though direct field testing is needed.

8 Before fish were added to the tanks, refuge use by kōura was similar across treatments but
9 highly variable at the individual level, reflecting substantial behavioural variation. The control
10 treatment showed increasing refuge use over time, particularly under light conditions, which
11 is unsurprising given that kōura are nocturnally adapted and need to remain hidden during
12 daylight to avoid visual predators such as cormorants ([Devcich 1979](#)). Importantly, treatment
13 effects on refuge use depended on light condition. Under dark conditions, when kōura are
14 naturally more active, catfish exposure resulted in significantly greater refuge use than eel
15 exposure. Control refuge use was intermediate between the two predator treatments, though
16 the reasons for this pattern are unclear from the current data. The contrasting responses of
17 kōura to catfish and eels indicates biologically meaningful differences in NCE between exposure
18 to a novel and native predator.

19 Unlike the response when exposed to catfish, kōura reduced their refuge use in the presence
20 of eels. This contrasting response likely reflects the long evolutionary relationship between
21 kōura and eels, though the strength of coevolutionary adaptation in these specific lakes remains
22 uncertain. The lakes naturally have restricted access to the coast limiting the number of
23 eels present and historical vulcanism has affected these dynamics with a potential historical
24 separation approximately 700 years ago following the Kaharoa eruption (~AD 1314; ([Hogg
25 et al. 2003](#))). Nevertheless, kōura probably were able to recognise chemical and visual cues
26 from the eels and respond in a proportionate rather than precautionary way ([Shave et al.
27 1994](#)). Instead of hiding, kōura may rely on contextual information, such as predator size,
28 movement patterns, or chemical concentrations, to modulate their response ([Wood and Moore
29 2020b](#)). This calibrated response makes ecological sense as longfin eels are the most widely
30 distributed fish in the freshwaters of Aotearoa ([McDowall 1990](#)), and a persistent high-intensity
31 refuge response to eels would likely impose substantial foraging costs. Whether shaped by deep
32 coevolutionary history or by recognition of eels as an established native predator, kōura show a
33 nuanced, context-dependent response to eels rather than an over-generalised avoidance strategy.
34 This pattern was most pronounced in the dark conditions, where kōura refuge use in all three

1 treatments began similarly but changed progressively over the course of the experiment. Kōura
2 in the catfish treatment steadily increased their refuge use, while kōura in the eel treatment
3 showed similar behaviour to those in the control treatment, with decreases to baseline levels by
4 the end of the observation period. This result suggests that under conditions most relevant
5 to natural activity by kōura, chemical recognition of the familiar eel predator is sufficient to
6 enable a calibrated, low-cost response, while the unfamiliar catfish continues to elicit sustained
7 precautionary refuge use (Shave et al. 1994; Wood and Moore 2020a). Beyond predator
8 recognition, the reduced refuge use observed in eel-exposed kōura may not reflect a lack of
9 response but could instead represent active escape-searching behaviour (i.e. a “flight” response).
10 If this was the behavioural motivation, kōura would be attempting to move away from an
11 area where they perceived themselves as vulnerable in the confined tank space, rather than
12 retreating to a refuge where they might be prone to a known predator.

13 **Feeding behaviour**

14 NCEs are not a single process but emerge from the interaction between sensory ecology and
15 environmental context. The absence of a kairomone effect on feeding in this study likely reflects
16 a combination of methodological constraints rather than an absence of NCEs on foraging
17 behaviour. The feeding experiment, in which predators were presented solely through chemical
18 cues without visual information, did not demonstrate a clear fish predator NCE on kōura. This
19 may have been due to the concentration of predator kairomones not being high enough to exceed
20 detection thresholds (Brown et al. 2004), the starvation conditions during the experiment, the
21 restricted time frame of the experiment, or a combination of these factors. Kōura can readily
22 detect necrotic fish tissue and readily leave refuges to scavenge on catfish, eel, and marine fish
23 carcasses with little discrimination between fish species, indicating that chemosensory responses
24 in kōura may depend on cue type, context, and concentration rather than predator identity
25 alone (MacNeil et al. 2025). The concentration of live fish exudates in our experiment relative
26 to natural conditions is difficult to determine, while kairomones were concentrated by confining
27 fish in a bucket, in our experimental tanks they were subsequently diluted in 5 L of water,
28 not replenished, and potentially degraded over the 24-hour period. Whether the resulting
29 concentration reflects, exceeds or falls below ecologically relevant levels in lakes where kōura
30 and predators coexist remains unclear, and future studies should attempt to quantify natural
31 kairomone concentrations to better calibrate experimental designs (Achtymichuk et al. 2025).
32 In addition, many crayfishes show a stronger response to alarm cues of conspecifics (Stebbing
33 et al. 2010; Gherardi et al. 2011; Ramberg-Pihl and Yurewicz 2020), and to kairomones from

1 predators on a conspecifics diet (Beattie and Moore 2018). Fish predators in our experiments
2 were maintained on the same diet as kōura for an extended acclimation period prior to exposure
3 (see Methods), which may have further dampened or negated the threat signals received from
4 kairomones (Smee and Weissburg 2006). Furthermore, the pre-experiment starvation period
5 for kōura may have been insufficient to elicit a very strong feeding response, as crayfish such
6 as kōura are opportunistic foragers which are necessarily adapted to going for prolonged
7 periods without feeding (Hazlett 2003). The length of time since feeding can influence foraging
8 behaviour, with previous studies indicating that crayfish will be more active foragers after
9 being starved for at least three days (Hazlett 2003). In the current study, the longest starvation
10 period was four days and at this point there was a clear discrepancy between the control feeding
11 rate and the much lower feeding rates in the two fish treatments. Allowing for ethics and
12 welfare concerns, if we had extended the starvation period, this may have elicited a stronger
13 and prolonged response. Additionally, the stress of confinement in an artificial setting may itself
14 have altered normal kōura feeding responses, as elevated baseline stress hormones are known
15 to modify foraging behaviour independently of predator cue exposure (Clinchy et al. 2012).
16 Finally, feeding was measured as a single endpoint after 24 hours, providing no information on
17 whether kōura initially suppressed feeding after kairomone exposure before eventually resuming
18 normal feeding activity (Wright et al. 2010). Further experiments using higher kairomone
19 concentrations, longer starvation periods, and continuous feeding monitoring would help clarify
20 whether non-visual predator cues generate NCEs on kōura foraging behaviour.

21 Conclusions

22 As ecosystems become increasingly novel under global change, the reliability of historical
23 predator–prey cues may decline, amplifying behavioural mismatches (Sih et al. 2011). The
24 present study provides experimental evidence that invasive brown bullhead catfish have the
25 potential to exert stronger NCEs on kōura, an endemic freshwater crayfish when compared
26 to their native predator, the longfin eel. The heightened refuge-seeking response to catfish
27 likely reflects a generalised antipredator response triggered by unfamiliar chemical and visual
28 cues, consistent with a fear generalisation framework. In contrast, the reduced response to eels
29 suggests that familiarity with native predators enables kōura to make more proportionate risk
30 assessments, although their behaviour may have also been an antipredator strategy. Future
31 studies should particularly include kōura from eel-inhabited waterways, which would determine
32 whether populations with a recent coevolutionary history with native eels display similar
33 proportionate responses, thereby confirming whether the patterns observed here reflect broader

1 adaptive differences across kōura populations with distinct predatory histories. Additionally,
2 comparative studies of kōura sourced from catfish-invaded versus catfish-free catchments would
3 be worthwhile to assess whether repeated exposure over multiple generations has resulted
4 in rapid evolution or behavioural plasticity in antipredator responses, and whether catfish-
5 experienced kōura show more calibrated responses to catfish presence than naïve kōura. Such
6 evolutionary dynamics could have important implications for predicting kōura persistence in
7 catfish-invaded systems (Sih et al. 2010). From a management perspective, translocation of
8 kōura from catfish-experienced source populations into catchments facing imminent catfish
9 invasion may represent a more feasible strategy for seeding populations with individuals
10 capable of more adaptive responses. Although we found no significant NCE of fish exudates
11 on feeding behaviour, our study suggests further investigations of non-visual predator cues on
12 kōura foraging behaviour is warranted. Because kōura are a keystone species that ecosystem
13 engineer, even a small increase in refuge use driven by catfish presence could have cascading
14 effects on sediment dynamics, decomposition, and nutrient cycling in invaded lakes. As brown
15 bullhead catfish continue to spread across Aotearoa’s rivers and lakes, management strategies
16 aimed at protecting kōura should account for the behavioural cost of catfish NCEs alongside
17 direct predation pressure as an underappreciated and ecologically significant dimension to this
18 invasion.

19 **Acknowledgements**

20 We thank Te Arawa Lakes Trust and Te Komiti Whakahaere for the opportunity to collect
21 and study the treasured kōura. We are grateful to Soweeta Fort-D’ath and William Anaru
22 (Te Arawa Lakes Trust), and Tihini Grant (Ngāti Pikiao). We also thank Nick Ling and Ben
23 Roche (University of Waikato) for assisting with catfish and eel collection.

24 **Funding statement**

25 This research was supported by the Fish Futures programme funded through a Ministry of
26 Business, Innovation and Employment grant (CAWX2101) with additional funding provided by
27 the Bay of Plenty Regional Council under the Toi huarewa Waimāori - Bay of Plenty Regional
28 Council Chair in Lake and Freshwater Science programme.

1 **Authors' contributions**

2 Study conception and design were undertaken by O.V.R., C.M., and R.H. Experiments were
3 executed by O.V.R. and C.M. Data analysis was performed by O.V.R. and F.L. I.A.K.K.
4 initiated the concept of this study for inclusion in the FishFutures programme, obtained iwi
5 consent, and provided local ecological knowledge and kōura specimens. Manuscript drafting
6 was led by O.V.R. All authors contributed to manuscript revision, approved the results, and
7 agreed on the final manuscript.

8 **Ethical approval**

9 This study received ethical approval by University of Waikato animal ethics committee, protocol
10 number: 1232. Additional permission to conduct experiments with kōura was granted by Te
11 Arawa Lakes Trust and Te Komiti Whakahaere.

12 **Declarations**

13 The corresponding author has declared that none of the authors has any competing interests.

14 **Data availability statement**

15 The code and derived data supporting the findings of this study are openly available on GitHub
16 (https://github.com/OlivierRaven/Koura_Fear) and a rendered version of the analysis is hosted
17 at https://olivierraven.github.io/Koura_Fear/. The primary raw dataset is archived on Zenodo
18 (<https://doi.org/10.5281/zenodo.19716761>). Peer review documents, including the cover letter
19 and reviewer responses, are available in the manuscript repository in the interest of open and
20 transparent science.

21 **Tables**

Table 1: Pairwise contrasts for kōura refuge use in light and dark conditions. Results are shown separately for each light condition from beta-binomial GLMMs. Odds ratios and p-values are shown for treatment and period comparisons. P-values are Sidak-adjusted for multiple comparisons within each factor.

Condition	Contrast	Odds ratio	SE	df	Null	z value	p value
Light	control / catfish	0.863	0.396	Inf	1	-0.322	0.944
Light	control / eel	1.641	0.734	Inf	1	1.108	0.509
Light	catfish / eel	1.903	0.866	Inf	1	1.414	0.333
Light	before / during	0.463	0.194	Inf	1	-1.836	0.158
Light	before / after	0.297	0.135	Inf	1	-2.672	0.021
Light	during / after	0.641	0.294	Inf	1	-0.969	0.597
Dark	control / catfish	0.393	0.172	Inf	1	-2.133	0.083
Dark	control / eel	1.542	0.655	Inf	1	1.019	0.565
Dark	catfish / eel	3.924	1.693	Inf	1	3.170	0.004
Dark	before / during	0.728	0.286	Inf	1	-0.807	0.698
Dark	before / after	0.527	0.215	Inf	1	-1.569	0.259
Dark	during / after	0.723	0.290	Inf	1	-0.809	0.698

¹ Source: [Article Notebook](#)

Figures

Supplementary information

Supplementary tables are provided in the thesis appendices. For standalone access, render supplementary.qmd in this chapter's folder.

Abrams PA (2000) The evolution of predator-prey interactions: Theory and evidence. Annual Review of Ecology and Systematics 31:79–105. <https://doi.org/10.1146/annurev.ecolsys.31.1.79>

Achtymichuk GH, Fish KM, Ferrari MCO (2025) The power of concentration: Antipredator responses to diluted frozen crayfish alarm cues provide insights on ecologically relevant




Figure 1: (a) Aquarium layout for kōura refuge seeking behaviour experiment. Each experimental round consisted of eight aquaria run simultaneously. Each tank contained a single kōura (K) and refuge (half terracotta plant pot) on one side of a mesh divider. On the other side was either a single catfish (C), eel (E) or a control with no fish. The mesh divider allowed visual and chemical cues while preventing direct physical contact between tank occupants. (b) Experiments were repeated in light and dark (red light), with photographs showing a catfish in the light regime (left) and an eel in the dark regime (red light) (right) separated from kōura by mesh dividers.




Figure 2: Predicted marginal probabilities of refuge use by treatment and experimental period in light and dark conditions. Results are shown separately for light (top) and dark (bottom) conditions, each fitted with a beta-binomial GLMM. Points represent model-estimated marginal means with 95% confidence intervals shown as error bars. Different letters (a, b) indicate statistically significant differences based on Tukey-adjusted pairwise comparisons within each factor. The dashed line at 0.5 marks the threshold where kōura spend half their time in refuge.




Figure 3: Kōura refuge use over the course of the experiment, showing the proportion of time spent in refuge across three periods (before, during, and after predator exposure) for each treatment (catfish: dark grey solid line; eel: medium grey dashed line; control: light grey dot-dash line) under light (top) and dark (bottom) conditions. Black dashed vertical lines indicate the start and end of predator exposure and grey vertical lines show individual observation times.




Figure 4: Mean ($\pm$ SE) kōura feeding rate (pellets consumed per gram wet weight kōura) across predator treatments (catfish, eel, control) and starvation periods (0, 3, and 4 days).

- 1 concentrations and updates to methodology. PLOS One 20:e0340001. <https://doi.org/10.1371/journal.pone.0340001>
- 2
- 3 Adams SB (2007) Direct and indirect effects of channel catfish (*Ictalurus punctatus*) on native
4 crayfishes (Cambaridae) in experimental tanks. The American Midland Naturalist 158:85–96.
5 [https://doi.org/10.1674/0003-0031\(2007\)158%5B85:daieoc%5D2.0.co;2](https://doi.org/10.1674/0003-0031(2007)158%5B85:daieoc%5D2.0.co;2)
- 6 Appelberg M, Soederbaeck B, Odelstroem T (1993) Predator detection and perception of
7 predation risk in the crayfish *Astacus astacus* L. Nordic journal of freshwater research
8 68:55–62
- 9 Barnes GE, Hicks BJ (2003) Brown bullhead catfish (*Ameiurus nebulosus*) in Lake Taupo. In:
10 Managing invasive freshwater fish in New Zealand: Proceedings of a workshop hosted by
11 Department of Conservation, 10–12 May 2001, Hamilton. Department of Conservation
- 12 Beattie MC, Moore PA (2018) Predator recognition of chemical cues in crayfish: Diet and
13 experience influence the ability to detect predation threats. Behaviour 155:505–530. <https://doi.org/10.1163/1568539x-00003501>
- 14
- 15 Blake MA, Hart PJB (1993) The behavioural responses of juvenile signal crayfish *Pacifastacus*
16 *leniusculus* to stimuli from perch and eels. Freshwater Biology 29:89–97. <https://doi.org/10.1111/j.1365-2427.1993.tb00747.x>
- 17
- 18 Brönmark C, Hansson L-A (2012) Chemical ecology in aquatic systems. Oxford University
19 Press
- 20 Brooks M, Bolker B, Kristensen K, et al (2017) [glmmTMB: Generalized linear mixed models](#)
21 [using template model builder](#). CRAN: Contributed Packages
- 22 Brown GE, Poirier J-F, Adrian JC (2004) Assessment of local predation risk: The role
23 of subthreshold concentrations of chemical alarm cues. Behavioral Ecology 15:810–815.
24 <https://doi.org/10.1093/beheco/arh084>
- 25 Brown LA (2009) [Habitat determinants and predatory interactions of the endemic freshwater](#)
26 [crayfish \(koura, *Paranephrops planifrons*\) in the lower North Island, New Zealand](#). Master's

1 thesis, Massey University

2 Carthey AJR, Banks PB (2014) Naïveté in novel ecological interactions: Lessons from theory and
 3 experimental evidence. *Biological Reviews* 89:932–949. <https://doi.org/10.1111/brv.12087>

4 Clinchy M, Sheriff MJ, Zanette LY (2012) Predator-induced stress and the ecology of fear.
 5 *Functional Ecology* 27:56–65. <https://doi.org/10.1111/1365-2435.12007>

6 Collier KJ, Parkyn SM, Rabeni CF (1997) Koura: A keystone species. *Water and Atmosphere*
 7 5:18–20

8 Dawkins R, Krebs JR (1979) Arms races between and within species. *Proceedings of the Royal*
 9 *Society of London B Biological Sciences* 205:489–511. [https://doi.org/10.1098/rspb.1979.](https://doi.org/10.1098/rspb.1979.0081)
 10 0081

11 Dedual M (2019) [Summary of the impacts and control methods of Brown Bullhead catfish](#)
 12 [\(*Ameiurus nebulosus* Lesueur 1815\) in New Zealand and overseas](#). Bay of Plenty Regional
 13 Council

14 Devcich AA (1979) An ecological study of *Paranephrops planifrons* White (Decapoda: Parastaci-
 15 dae) in Lake Rotoiti, North Island. PhD thesis, University of Waikato

16 Doherty TS, Glen AS, Nimmo DG, et al (2016) Invasive predators and global biodiversity loss.
 17 *Proceedings of the National Academy of Sciences* 113:11261–11265. [https://doi.org/10.](https://doi.org/10.1073/pnas.1602480113)
 18 1073/pnas.1602480113

19 Driscoll J, Kola M, Mathews L (2020) Perception of alarm cues influences the outcome of
 20 shelter competition in crayfish. *Ethology* 126:584–592. <https://doi.org/10.1111/eth.13011>

21 Ehlman SM, Trimmer PC, Sih A (2019) Prey responses to exotic predators: Effects of old risks
 22 and new cues. *The American Naturalist* 193:575–587. <https://doi.org/10.1086/702252>

23 Feldmann RM, Pole M (1994) A new species of *Paranephrops* White, 1842: A fossil freshwater
 24 crayfish (Decapoda: Parastacidae) from the Manuhirikia Group (Miocene), Central Otago,
 25 New Zealand. *New Zealand Journal of Geology and Geophysics* 37:163–167. <https://doi.org/10.1080/00288409408839511>

- 1 [org/10.1080/00288306.1994.9514611](https://doi.org/10.1080/00288306.1994.9514611)
- 2 Figler MH, Blank GS, Peeke HVS (2005) Shelter competition between resident male red
3 swamp crayfish *Procambarus clarkii* (Girard) and conspecific intruders varying by sex
4 and reproductive status. *Marine and Freshwater Behaviour and Physiology* 38:237–248.
5 <https://doi.org/10.1080/10236240500376477>
- 6 Francis LB (2019) [Evaluating the effects of invasive brown bullhead catfish \(*Ameiurus nebulosus*\)](#)
7 [on kōura \(freshwater crayfish, *Paranephrops planifrons*\) in Lake Rotoiti](#). Master’s thesis,
8 University of Waikato
- 9 Gherardi F, Mavuti KM, Pacini N, et al (2011) The smell of danger: Chemical recognition of fish
10 predators by the invasive crayfish *Procambarus clarkii*. *Freshwater Biology* 56:1567–1578.
11 <https://doi.org/10.1111/j.1365-2427.2011.02595.x>
- 12 Hartig F, Lohse L, leite M de S (2024) [DHARMa: Residual diagnostics for hierarchical](#)
13 [\(multi-level / mixed\) regression models](#)
- 14 Hazlett BA (2003) The effects of starvation on crayfish responses to alarm odor. *Ethology*
15 109:587–592. <https://doi.org/10.1046/j.1439-0310.2003.00902.x>
- 16 Hicks BJ (1997) Food webs in forest and pasture streams in the Waikato region, New Zealand:
17 A study based on analyses of stable isotopes of carbon and nitrogen, and fish gut contents.
18 *New Zealand Journal of Marine and Freshwater Research* 31:651–664. [https://doi.org/10.](https://doi.org/10.1080/00288330.1997.9516796)
19 [1080/00288330.1997.9516796](https://doi.org/10.1080/00288330.1997.9516796)
- 20 Hogg AG, Higham TFG, Lowe DJ, et al (2003) A wiggle-match date for Polynesian settlement
21 of New Zealand. *Antiquity* 77:116–125. <https://doi.org/10.1017/s0003598x00061408>
- 22 Johnson MF, Rice SP, Reid I (2014) The activity of signal crayfish (*Pacifastacus leniusculus*)
23 in relation to thermal and hydraulic dynamics of an alluvial stream, UK. *Hydrobiologia*
24 724:41–54. <https://doi.org/10.1007/s10750-013-1708-1>
- 25 Kasumyan AO (2022) Fish as sources of kairomones-chemical signals for aquatic animals.
26 *Journal of Ichthyology* 62:289–315. <https://doi.org/10.1134/s0032945222020084>

- 1 Kaulfuss U, Lee DE, Wartho J-A, et al (2018) Geology and palaeontology of the Hindon Maar
2 Complex: A Miocene terrestrial fossil *Lagerstätte* in southern New Zealand. Palaeogeography,
3 Palaeoclimatology, Palaeoecology 500:52–68. <https://doi.org/10.1016/j.palaeo.2018.03.022>
- 4 Kindinger TL, Albins MA (2017) Consumptive and non-consumptive effects of an invasive
5 marine predator on native coral-reef herbivores. Biological Invasions 19:131–146. <https://doi.org/10.1007/s10530-016-1268-1>
- 6
- 7 Kusabs IAK, Duggan IC, Bruere AC, et al (2026) Invasive catfish linked to the decline of
8 freshwater crayfish in two Aotearoa New Zealand lakes. Biological Invasions 28: <https://doi.org/10.1007/s10530-026-03852-0>
- 9
- 10 Kusabs IA, Quinn JM (2009) Use of a traditional Māori harvesting method, the tau kōura,
11 for monitoring kōura (freshwater crayfish, *Paranephrops planifrons*) in Lake Rotoiti, North
12 Island, New Zealand. New Zealand Journal of Marine and Freshwater Research 43:713–722.
13 <https://doi.org/10.1080/00288330909510036>
- 14 Laundré JW, Hernández L, Altendorf KB (2001) Wolves, elk, and bison: Reestablishing the
15 “landscape of fear” in Yellowstone National Park, U.S.A. Canadian Journal of Zoology
16 79:1401–1409. <https://doi.org/10.1139/z01-094>
- 17 Lee F, Kusabs IAK, Perry GLW, MacNeil C (2025) Identifying refugia from the synergistic
18 threats of climate change and invasive species. Web Ecology 25:221–239. <https://doi.org/10.5194/we-25-221-2025>
- 19
- 20 Lenth RV, Piaskowski J, Banfai B, et al (2026) `emmeans`: Estimated marginal means, aka
21 least-squares means
- 22 Lima SL, Dill LM (1990) Behavioral decisions made under the risk of predation: A review and
23 prospectus. Canadian Journal of Zoology 68:619–640. <https://doi.org/10.1139/z90-092>
- 24 MacNeil C, Lee F, Kusabs I, Holmes R (2025) The biter bit; scavenging behaviour of native
25 freshwater crayfish on the carrion of native and introduced fish predators in Aotearoa-New
26 Zealand. Knowledge & Management of Aquatic Ecosystems 29. <https://doi.org/10.1051/kmae/2025022>
- 27

- 1 Martin CW (2014) Naïve prey exhibit reduced antipredator behavior and survivorship. PeerJ
2 2:e665. <https://doi.org/10.7717/peerj.665>
- 3 Martin M, Boubée J, Kusabs I (2007) [Taonga and mahinga kai of the Te Arawa lakes: A review](#)
4 [of current knowledge - tuna](#). National Institute of Water & Atmospheric Research Ltd,
5 Hamilton, New Zealand
- 6 McDowall RM (1990) [New Zealand freshwater fishes: A natural history and guide](#)
- 7 Nyström P (2005) Non-lethal predator effects on the performance of a native and an exotic
8 crayfish species. Freshwater Biology 50:1938–1949. [https://doi.org/10.1111/j.1365-2427.](https://doi.org/10.1111/j.1365-2427.2005.01438.x)
9 [2005.01438.x](#)
- 10 Parkyn SM, Collier KJ, Hicks BJ (2001) New Zealand stream crayfish: Functional omnivores
11 but trophic predators? Freshwater Biology 46:641–652. [https://doi.org/10.1046/j.1365-](https://doi.org/10.1046/j.1365-2427.2001.00702.x)
12 [2427.2001.00702.x](#)
- 13 Peacor SD, Werner EE (2001) The contribution of trait-mediated indirect effects to the net
14 effects of a predator. Proceedings of the National Academy of Sciences 98:3904–3908.
15 <https://doi.org/10.1073/pnas.071061998>
- 16 Preisser EL, Bolnick DI, Benard MF (2005) Scared to death? The effects of intimidation and
17 consumption in predator-prey interactions. Ecology 86:501–509. [https://doi.org/10.1890/04-](https://doi.org/10.1890/04-0719)
18 [0719](#)
- 19 R Core Team (2025) [R: A language and environment for statistical computing](#). R Foundation
20 for Statistical Computing, Vienna, Austria
- 21 Ramberg-Pihl NC, Yurewicz KL (2020) Behavioral responses of northern crayfish (*Faxonius*
22 *virilis*) to conspecific alarm cues and predator cues from smallmouth bass (*Micropterus*
23 *dolomieu*). Marine and Freshwater Behaviour and Physiology 53:1–16. [https://doi.org/10.](https://doi.org/10.1080/10236244.2020.1717338)
24 [1080/10236244.2020.1717338](#)
- 25 Ryan P (2009) [Eels | Te Ara Encyclopedia of New Zealand](#)

- 1 Schmitz OJ, Beckerman AP, O'Brien KM (1997) Behaviorally mediated trophic cascades:
2 Effects of predation risk on food web interactions. *Ecology* 78:1388–1399. [https://doi.org/](https://doi.org/10.1890/0012-9658(1997)078%5B1388:bmtceo%5D2.0.co;2)
3 [10.1890/0012-9658\(1997\)078%5B1388:bmtceo%5D2.0.co;2](https://doi.org/10.1890/0012-9658(1997)078%5B1388:bmtceo%5D2.0.co;2)
- 4 Scott WB, Crossman EJ (1973) *Freshwater fishes of Canada*. Fisheries Research Board of
5 Canada
- 6 Shave CathyR, Townsend CR, Crowl TA (1994) Anti-predator behaviours of a freshwater
7 crayfish (*Paranephrops zealandicus*) to a native and an introduced predator. *New Zealand*
8 *Journal of Ecology* 18:1–10
- 9 Sheriff MJ, Peacor SD, Hawlena D, Thaker M (2020) Non-consumptive predator effects on
10 prey population size: A dearth of evidence. *Journal of Animal Ecology* 89:1302–1316.
11 <https://doi.org/10.1111/1365-2656.13213>
- 12 Sih A, Bolnick DI, Luttbeg B, et al (2010) Predator-prey naïveté, antipredator behavior, and
13 the ecology of predator invasions. *Oikos* 119:610–621. [https://doi.org/10.1111/j.1600-](https://doi.org/10.1111/j.1600-0706.2009.18039.x)
14 [0706.2009.18039.x](https://doi.org/10.1111/j.1600-0706.2009.18039.x)
- 15 Sih A, Chung HJ, Neylan I, et al (2023) Fear generalization and behavioral responses to multiple
16 dangers. *Trends in Ecology & Evolution* 38:369–380. [https://doi.org/10.1016/j.tree.2022.](https://doi.org/10.1016/j.tree.2022.11.001)
17 [11.001](https://doi.org/10.1016/j.tree.2022.11.001)
- 18 Sih A, Ferrari MCO, Harris DJ (2011) Evolution and behavioural responses to human-induced
19 rapid environmental change. *Evolutionary Applications* 4:367–387. [https://doi.org/10.](https://doi.org/10.1111/j.1752-4571.2010.00166.x)
20 [1111/j.1752-4571.2010.00166.x](https://doi.org/10.1111/j.1752-4571.2010.00166.x)
- 21 Smee DL, Weissburg MJ (2006) Hard clams (*Mercenaria mercenaria*) evaluate predation risk
22 using chemical signals from predators and injured conspecifics. *Journal of Chemical Ecology*
23 32:605–619. <https://doi.org/10.1007/s10886-005-9021-8>
- 24 Sniegula S, Konczarek D, Bonk M, et al (2025) Non-consumptive effects of native, alien and
25 invasive alien crayfish on damselfly egg life history and carry-over effects on larval physiology.
26 *NeoBiota* 97:215–235. <https://doi.org/10.3897/neobiota.97.139760>

- 1 Stebbing PD, Watson GJ, Bentley MG (2010) The response to disturbance chemicals and
2 predator odours of juvenile and adult signal crayfish *Pacifastacus leniusculus* (Dana).
3 Marine and Freshwater Behaviour and Physiology 43:183–195. [https://doi.org/10.1080/](https://doi.org/10.1080/10236244.2010.487301)
4 [10236244.2010.487301](https://doi.org/10.1080/10236244.2010.487301)
- 5 Stein RA, Magnuson JJ (1976) Behavioral response of crayfish to a fish predator. Ecology
6 57:751–761. <https://doi.org/10.2307/1936188>
- 7 Usio N, Townsend CR (2004) Roles of crayfish: Consequences of predation and bioturbation
8 for stream invertebrates. Ecology 85:807–822. <https://doi.org/10.1890/02-0618>
- 9 Wood TC, Moore PA (2020b) Big and bad: How relative predator size and dietary information
10 influence rusty crayfish (*Faxonius rusticus*) behavior and resource-use decisions. Canadian
11 Journal of Zoology 98:62–72. <https://doi.org/10.1139/cjz-2019-0089>
- 12 Wood TC, Moore PA (2020a) Fine-tuned responses to chemical landscapes: Crayfish use
13 predator odors to assess threats based on relative size ratios. Ecosphere 11:e03188. <https://doi.org/10.1002/ecs2.3188>
14 <https://doi.org/10.1002/ecs2.3188>
- 15 Wright TF, Eberhard JR, Hobson EA, et al (2010) Behavioral flexibility and species invasions:
16 The adaptive flexibility hypothesis. Ethology Ecology & Evolution 22:393–404. <https://doi.org/10.1080/03949370.2010.505580>
17 <https://doi.org/10.1080/03949370.2010.505580>
- 18 Zhao M, Feng G, Wang H, et al (2024) The influence of shelter type and coverage on crayfish
19 (*Procambarus clarkii*) predation by catfish (*Silurus asotus*): A controlled environment study.
20 Animals 14:1147. <https://doi.org/10.3390/ani14081147>