

The importance of structure: mesocosm evidence for shelter design in freshwater crayfish restoration

Olivier V. Raven Robin Holmes Ian A. K. Kusabs
Francis J. Burdon Deniz Özkundakci

2026-07-08

Freshwater crayfish are keystone species that rank among the most threatened invertebrate groups globally. While habitat loss and predation by non-native predators feature prominently as drivers of crayfish population declines, the optimal design of habitat enhancement structures for their restoration remains largely unknown. To better understand freshwater crayfish habitat preferences, our study focused on the endemic kōura (*Paranephrops planifrons*) found in Aotearoa New Zealand. We used four mesocosm experiments to examine how shelter structure, material composition, individual body size, and group dynamics influence refuge selection in kōura. Individual kōura strongly preferred a wooden log with enclosed drilled holes, but when housed in mixed-size groups, their preference shifted to flat and piled stone configurations that provided more complex refugia for individuals of different sizes. Flat stone structures were preferred over split wood pieces of similar dimensions, and this was strongest for large individuals. When structural dimensions were held constant, no significant preference was detected between concrete and wooden structures, indicating that material composition alone does not drive refuge selection. The findings of these mesocosm experiments suggest that stone pile configurations warrant field evaluations as habitat enhancement structures for kōura restoration. The stone piles create structural complexity that can provide habitat for different size classes and thus may add critical refuge from non-native predators for kōura populations.

1 **Table of contents**

2	Introduction	5
3	Materials and Methods	7
4	Animal collection and tank setup	7
5	Experiment 1: Individual preference	8
6	Experiment 2: Group dynamics	9
7	Experiment 3: Wood vs Stone	9
8	Experiment 4: Material comparison	10
9	Data analysis	10
10	Results	12
11	Experiment 1: Individual preference	12
12	Experiment 2: Group dynamics	12
13	Experiment 3: Wood vs Stone	13
14	Experiment 4: Material comparison	13
15	Discussion	14
16	Structure matters more than material	14
17	Body size and competitive dynamics	15
18	Restoration implications	16
19	Conclusion	17
20	Acknowledgements	17
21	Funding statement	17
22	Authors' contributions	17
23	Ethical approval	18
24	Declarations	18
25	Data availability statement	18
26	Figures	18

1 Introduction

2 Freshwater biodiversity is declining faster than in any other ecosystem globally, driven by
3 habitat loss, pollution, and invasive species (Dudgeon *et al.*, 2006). Crayfish are among the
4 most threatened freshwater invertebrates, with 32 % of species assessed as threatened with
5 extinction due to these same pressures (Richman *et al.*, 2015). As keystone species and
6 ecosystem engineers, crayfish play crucial roles in freshwater food webs through shredding of
7 leaf litter, bioturbation of sediments, and nutrient cycling, and their loss can have profound
8 cascading effects on freshwater ecosystems (Momot, 1995; Collier *et al.*, 1997; Parkyn *et al.*,
9 2001). The increasing degradation of freshwater habitats therefore not only threatens crayfish
10 populations but also risks disrupting the ecosystem processes crayfish sustain.

11 Habitat complexity is a key driver of crayfish population persistence, as structurally diverse
12 habitats provide physical refugia from predators, foraging opportunities, and shelter during
13 moulting when individuals are particularly vulnerable (Hammond *et al.*, 2006). Shelter
14 availability directly increases crayfish survival and growth under predation pressure (Adams,
15 2007), making refuge quality a critical consideration for both natural habitat persistence and
16 restoration design. Crayfish use a wide variety of natural and artificial structures as shelter,
17 including rocks, woody debris, and even anthropogenic objects such as bottles and discarded
18 appliances (Parkyn *et al.*, 2009; Adams, 2014), demonstrating considerable flexibility in habitat
19 use. While this diversity of potential refugia is well documented, preferences among different
20 structure types remain poorly understood, even though knowledge of habitat preferences is
21 fundamental to designing effective enhancement structures for restoration. In the context of
22 freshwater restoration, increasing structural complexity through providing shelters has improved
23 crayfish abundance and restocking success in degraded streams and lakes (Huolila *et al.*, 1997;
24 Johnsen and Taugbøl, 2008), and the availability of coarser substrate has also been shown to
25 improve juvenile survival rates (Mazlum *et al.*, 2017). These findings indicate that enhancing
26 habitat complexity can be a useful management tool for threatened crayfish populations, yet
27 systematic comparisons of shelter configurations for freshwater crayfish restoration remain
28 largely absent from the literature.

29 Shelter selection in crayfish is strongly influenced by body size, as individuals actively select
30 refugia proportional to their dimensions. Crayfish individuals tend to select stones more than
31 3 times longer and 1.25 times wider than their carapace length, with larger males consistently
32 selecting larger stones (Streissl and Hödl, 2002). Social dynamics further complicate shelter use
33 when refuge availability is limited, as competitive interactions among conspecifics determine

1 which individuals gain access to preferred shelters. Dominance hierarchies based on body size
 2 and reproductive status show that males and maternal females often are able to take over shelters
 3 from non-maternal females, while adults consistently outcompete juveniles for shelter access
 4 (Figler *et al.*, 2005). Body size and shelter dimensions have direct competitive consequences as
 5 crayfish in undersized shelters initiated and won more fights than those in appropriately sized
 6 shelters, suggesting that shelter fit influences an individual's assessment of its own fighting
 7 ability (Percival and Moore, 2010). Body size ratios between contestants are a strong predictor
 8 of shelter takeover success, although prior ownership can offset size disadvantages (Ranta and
 9 Lindström, 1993), particularly when both individuals hold high-quality shelters (Takahashi *et al.*,
 10 2019). At the population level, habitat complexity buffers these competitive effects, with
 11 juveniles in structurally complex habitats showing higher survival, faster growth, and greater
 12 moulting rates as more shelter availability reduces agonistic interactions (Olsson and Nyström,
 13 2009). Together, shelter use in crayfish reflects an interaction between individual body size and
 14 social competitive dynamics, suggesting that effective habitat enhancement structures must
 15 provide a range of shelter sizes to support different life stages and size classes simultaneously.

16 The material and structural properties of shelters can influence their value as refugia. In a
 17 study comparing shelter configurations on reef fish assemblages, the availability and size of
 18 enclosed spaces mattered more than total structure volume (Hylkema *et al.*, 2020). Additionally,
 19 enclosed structures such as PVC pipes provide greater protection against predatory fish than
 20 stone or vegetation structures, likely due to their consistent hole dimensions (Zhao *et al.*, 2024).
 21 However, the use of synthetic materials like PVC pipes in restoration ecology brings ecological
 22 concerns, particularly in contexts where introducing non-native materials may conflict with
 23 conservation values or indigenous cultural frameworks (Wehi and Lord, 2017; Cooke *et al.*,
 24 2023). Natural materials such as stones and wood offer ecologically compatible alternatives,
 25 yet the importance of structural configuration versus material composition for crayfish shelter
 26 preference is poorly understood.

27 In Aotearoa-New Zealand (hereafter Aotearoa), the endemic freshwater crayfish kōura
 28 (*Paranephrops planifrons*) faces conservation pressure from habitat loss, water quality
 29 degradation, and predation by invasive species (Lee *et al.*, 2025; Kusabs *et al.*, 2026). Kōura
 30 are a taonga (culturally treasured) species for Māori, the indigenous people of Aotearoa, and
 31 hold significant ecological and cultural value in freshwater ecosystems (Kusabs *et al.*, 2015a;
 32 Kusabs and Quinn, 2009). In their natural habitat, kōura are positively associated with
 33 overhanging banks, coarse substrates, and woody debris (Usio and Townsend, 2000; Jowett
 34 *et al.*, 2008; Parkyn *et al.*, 2009; Raven *et al.*, (a) in review). Kōura have been shown to

1 significantly increase refuge use in the presence of predators (Shave *et al.*, 1994; Raven *et al.*,
2 (b) in review). Habitat restoration by increasing shelter availability represents a promising
3 management tool for kōura recovery, and the use of natural materials aligns with ecological
4 and cultural values.

5 In this study, four mesocosm experiments were conducted to identify kōura habitat preferences.
6 We assessed how different structures varying in material and configuration influence refuge
7 selection, accounting for individual behaviour, body size, and group interactions. We predicted
8 that 1) Structurally complex structures providing enclosed crevices would be preferred over
9 isolated or open structures, as enclosed spaces reduce predation risk; 2) structure preference
10 would be size-dependent, with larger kōura selecting structures with larger refuge spaces (Streissl
11 and Hödl, 2002; Percival and Moore, 2010); 3) stone-based structures would be preferred over
12 wood due to greater structural stability, and 4) material type alone would not influence refuge
13 preference between concrete and wooden bricks. The results provide practical guidance for
14 designing cost-effective, ecologically appropriate habitat enhancement structures to enhance
15 kōura abundance and support population recovery.

16 **Materials and Methods**

17 **Animal collection and tank setup**

18 In mid-August 2025, 72 kōura were sourced from Lakes Ōkāreka and Tikitapu, located within
19 the Rotorua Te Arawa Lakes area in the North Island of Aotearoa. Collected animals were
20 transported to the Aquatic Research Centre at the University of Waikato in Hamilton. Individ-
21 uals were assigned to one of three size classes based on orbital carapace length (OCL): small (<
22 22 mm), medium (22–30 mm), and large (> 30 mm). Each animal was marked with a unique
23 identification number written on the carapace using an orange oil-based paint marker (Sharpie,
24 medium point) (Ramalho *et al.*, 2010).

25 Kōura were acclimatised for 12 days in outdoor concrete holding tanks (Ø 1.4 m), each filled to
26 0.4 m with dechlorinated tap water conditioned for three months, paving sand as substrate,
27 shelters, and were covered with a single layer of shade cloth. Each holding tank housed
28 10–12 individuals. Kōura were fed every second day with commercial carnivore pellets (Hikari
29 Tropical, Kyorin Food Industries Ltd., Japan; animal protein 47 %). Water temperature and
30 dissolved oxygen remained stable throughout the experimental period (temperature: 12 ± 1.2
31 °C; dissolved oxygen: 99.9 ± 1.7 %).

1 Experiments were conducted in six larger outdoor concrete tanks (1425 L; Ø 1.85 m) with the
2 same substrate and water conditions, covered with a double layer of shade cloth to prevent
3 escape and minimise solar heating. Fresh dechlorinated tap water and an air stone were
4 added between rounds. To limit thermal stress, experiments were conducted during the
5 Australasian spring (September to October). Because kōura are a taonga (treasured) species,
6 permission to conduct the experiments was obtained from Te Arawa Lakes Trust and Te Komiti
7 Whakahaere. All experimental procedures were conducted under University of Waikato animal
8 ethics committee approval (protocol number: 1207; see Ethical Approval section).

9 **Experiment 1: Individual preference**

10 This experiment assessed individual kōura structure preference by offering single animals a
11 choice among four structures (Figure 1). The structure-types were: SingleStone (ten quarried
12 stones (150–250 mm) placed individually across the quadrant with sufficient spacing between
13 stones that no crevices formed between them), FlatStone (ten quarried stones arranged tightly
14 without stacking), StonePile (ten quarried stones stacked to form a pile), and WoodLog (a
15 tānekaha (*Phyllocladus trichomanoides*) log (500 mm long, 150 mm diameter) containing six
16 drilled holes (70 mm deep, 32 mm diameter)). If a kōura was not using a structure, TankWall
17 was noted as its location which represents the wall of the tank or when the kōura was in the
18 open. TankWall functioned as an indicator of structure rejection rather than a preference.
19 Six experimental tanks were used, each housing a single medium-sized kōura. Across five
20 experimental replicates, a total of 30 kōura were tested, consisting of 15 females with mean
21 OCL and weight $\pm$ SD: 26.9 ± 2 mm, range: 23.2–29.8 mm, 12.5 ± 3.3 g and 15 males: $26.1 \pm$
22 2.2 mm, range: 22.6–30 mm, 11.9 ± 3.1 g. Each replicate lasted 22 hours, beginning at noon
23 (12:00) and ending at 10:00 the following morning.

24 Each experimental tank was divided into four quadrants, with one structure type placed at
25 random (randomizer.org) in each quadrant. For each trial, a single kōura was placed inside
26 a vertically oriented plastic tube (50 mm long, open at one end and sealed at the other)
27 positioned at the centre of the tank, with the sealed end resting on the substrate. This ensured
28 a standardised starting position from which the kōura could exit in any direction (MacNeil *et*
29 *al.*, 2025). The time taken for the kōura to locate its initial refuge location was recorded. On
30 the following day, the final location of the kōura was noted after which it was removed from
31 the tank. Tanks were reset and structure placements re-randomised prior to the next trial.
32 Each individual kōura was used once in a single trial and not reused across temporal replicate

1 blocks. To confirm nocturnal activity, three digital cameras (GoPro Hero 3) were placed in one
2 tank and recorded for five hours at the end of each of the five days.

3 **Experiment 2: Group dynamics**

4 Building on Experiment 1, this experiment examined how group dynamics and body size
5 influenced structure preference when kōura were housed together. The same six experimental
6 tanks were used and configured with the identical structure types and randomisation procedure
7 as in Experiment 1 (Figure 1). In contrast to the single-animal trials, each tank housed a
8 group of six male kōura, as male-only groups simplified competitive dynamics and size class
9 assignment given limited female availability. Groups were comprised of two small, two medium,
10 and two large individuals. Size classes were defined by OCL: small (mean OCL and weight
11 $\pm$ SD: 19.2 ± 2 mm, range: 15.8–21.6 mm, 3.9 ± 2.1 g), medium (26.4 ± 1.6 mm, 23.5–29.5
12 mm, 11.6 ± 2.4 g), and large (36.3 ± 4.2 mm, 30.7–43.3 mm, 31.8 ± 11.2 g). Each group was
13 introduced to the centre of the tank within an open-top plastic container ($150 \times 150 \times 85$
14 mm). The container was removed, and the initial location of each individual was recorded. On
15 the following day, the final location of each kōura was documented, before all individuals were
16 removed from the tank. This procedure was repeated across five experimental replicate blocks,
17 resulting in 30 group-trials in total (six groups $\times$ five blocks). Structure placements and tank
18 assignments were re-randomised between blocks, such that each group was housed in a different
19 tank in each successive round. The same six groups were reused across all five blocks, with
20 individuals staying with their assigned group between replicates. No mortality occurred during
21 the experiment.

22 **Experiment 3: Wood vs Stone**

23 Using the same male kōura groups from Experiment 2, this experiment narrowed the choice to
24 two structure types, stone and wood, to directly compare preference between these materials.
25 The structure types were: (1) a flat stone structure (FlatStone) constructed from twelve stones
26 placed tightly together, and (2) a wood structure consisting of twelve wood splits (WoodSplit)
27 made from the tānekaha logs (Figure 1). Each group was released at the centre of the tank,
28 and the initial structure selected by each kōura was recorded. The following day, the final
29 location of each kōura was documented. This procedure was repeated across three experimental
30 replicate blocks, resulting in 18 group-trials in total (six groups $\times$ three blocks). Structure

1 placements and tank assignments were re-randomised between blocks, such that each group
2 was housed in a different tank in each successive round. One small individual died in the first
3 block and was replaced with a size-matched individual to maintain group composition. One
4 individual could not be located at the end of the final block and was assumed to have escaped
5 the tank.

6 **Experiment 4: Material comparison**

7 This experiment tested whether kōura showed a preference between perforated concrete bricks
8 and perforated wooden blocks to further identify the effect of material on structure preference.
9 Two perforated building bricks (ConcreteBrick) of $80 \times 80 \times 225$ mm with five holes ($\varnothing$ 30 mm)
10 were tested against two wooden blocks (WoodBlock) with drilled holes made from Western red
11 cedar (*Thuja plicata*), constructed to match the building bricks in external dimensions and
12 hole size. To prevent floating, a solid clay brick of identical dimensions was placed on top of
13 each WoodBlock. ConcreteBrick did not require additional weight (Figure 1). New groups of
14 ten kōura and four experiment tanks were used for this experiment, comprising a total of 40
15 individuals consisting of 18 females with mean OCL and weight $\pm$ SD: 26.2 ± 3.1 mm, range:
16 $20.2\text{--}30.8$ mm, 12.2 ± 4.2 g and 22 males: 26 ± 2.1 mm, range: $20.9\text{--}29.5$ mm, 11.6 ± 3.2 g.
17 Due to reduced animal availability, individuals were drawn from a slightly broader size range
18 than medium-size individuals, resulting in unbalanced size class representation (small: $n = 4$,
19 medium: $n = 35$, large: $n = 1$). In each of the four experimental tank quadrants one structure
20 was placed, with similar structure types positioned opposite to one another. Each group of
21 kōura was released at the centre of the tank and the following day, the final location of each
22 kōura was documented. This experiment was repeated across four temporal blocks using the
23 same four groups of ten kōura, resulting in 16 group-trials in total (four groups $\times$ four blocks).
24 Structure placements and tank assignments were re-randomised between blocks, such that each
25 group was housed in a different tank in each successive round.

26 **Data analysis**

27 Experiment 1: To test the difference between the four structure types selected by the single
28 kōura, a multinomial goodness-of-fit test was performed using the XNomial package (Engels,
29 2014). TankWall was excluded from preference tests as it represents the absence of structure
30 choice rather than a structure type preference. Additionally, pairwise likelihood-ratio G-tests

1 with Bonferroni correction were used to identify which structure types differed significantly
2 from each other. Bayesian categorical mixed-effects models were used to test whether body
3 size or sex affected structure preference across all experiments, fitted using the brms package
4 (Bürkner *et al.*, 2025). For Experiments 1 and 2, where kōura chose among four structure
5 types, a categorical family was used. For Experiments 3 and 4, where only two structure types
6 were directly compared, a Bernoulli family was used. Model convergence was assessed using
7 the Gelman-Rubin diagnostic (all $\hat{R} = 1.00$) and divergent transition counts (all experiments:
8 0 divergences), confirming reliable posterior sampling (Gelman and Rubin, 1992). To test
9 whether structure type use differed between sexes, the Bayesian categorical mixed-effects model
10 was fitted with structure type as the outcome, sex as a fixed effect and tank and experimental
11 round as random intercepts. A permutation-based symmetry test was applied to assess whether
12 structure type transitions between initial location and next-day location were directional rather
13 than random.

14 Experiment 2: To test whether overall structure preference existed at the group level, one-
15 sample Wilcoxon signed-rank tests were applied to group-round proportions for each structure
16 type, testing against a null expectation of 0.25 (equal distribution among four structure
17 types). P-values were Bonferroni adjusted for multiple comparisons. Normal approximation
18 was used due to ties arising from discrete group counts. To test whether size classes affected
19 structure preference while accounting for repeated measurement of individuals and groups, the
20 Bayesian categorical mixed-effects model was fitted with structure type as the outcome, size
21 class as a fixed effect, and group identity, individual identity nested within group, and tank
22 and experimental round as random intercepts. The Stuart-Maxwell test was used to test for
23 statistical differences between initial and next-day location.

24 Experiment 3: To test the difference between FlatStone and WoodSplit the Wilcoxon signed-
25 rank test was applied to group-round proportions for each structure type, testing against a null
26 expectation of 0.5. P-values were Bonferroni adjusted. To test whether size classes affected
27 structure preference while accounting for repeated measurement of individuals and groups, the
28 Bayesian Bernoulli mixed-effects model was fitted with size class as a fixed effect, following the
29 same random effects structure as described for Experiment 2. The Stuart-Maxwell test was
30 used to test for statistical differences between initial and next-day location.

31 Experiment 4: To test whether kōura showed a preference between ConcreteBrick and Wood-
32 Block, the Wilcoxon signed-rank test was applied to group-round proportions for each structure
33 type, testing against a null expectation of 0.5 and p-values were Bonferroni adjusted. A paired
34 Wilcoxon signed-rank test was used to directly compare group-round proportions between

ConcreteBrick and WoodBlock. As size class was unbalanced, the effect of sex on structure preference was tested using the Bayesian Bernoulli mixed-effects model with sex as a fixed effect, following the same random effects structure as described for Experiment 2. Finally, as only next-day locations were recorded in this experiment, no directional shift analysis was performed.

Results

Experiment 1: Individual preference

Individual kōura showed a strong preference for WoodLog, selecting it significantly more often than SingleStone, FlatStone, and StonePile (pairwise G-test, Bonferroni adjusted: $p < 0.001$, 0.011, 0.004, respectively; overall multinomial test: $p < 0.001$; Figure 2). No significant differences were found among the stone structure types. Overall, WoodLog was selected by 67 % of kōura at refuge (males: 80 %; females: 53 %), and no kōura were found at TankWall. Sex had no credible effect on structure preference ($\beta = 3.32$, 95% CI: -1.09 to 9.3; posterior probability = 0.92) with credible intervals spanning zero for all structure types. The symmetry test indicated that structure type use shifted between initial and next-day location ($p < 0.001$), suggesting kōura actively relocated to preferred structure types overnight rather than remaining at their initial location. The video footage showed kōura emerging, walking and climbing on the structures after sunset.

Experiment 2: Group dynamics

When housed in mixed-size groups, kōura showed a significant overall preference for stone-based structures (Figure 3). FlatStone was used significantly more ($p = 0.004$), while SingleStone ($p < 0.001$) and WoodLog ($p = 0.039$) were significantly avoided. StonePile did not differ from random expectation ($p = 1.000$) and did not differ significantly from FlatStone ($p = 0.117$). No kōura were found at TankWall, indicating active structure selection across the group. Size classes had limited but credible effect on structure preference. Medium-sized kōura showed a weaker preference for FlatStone over SingleStone compared to small kōura ($\beta = -1.77$, 95% CI: -3.7 to -0.14; posterior probability = 0.99), while no other size class effect was credible. Overall, FlatStone remained the most selected structure type across all size classes, but its strength of preference declined from small (posterior probability of top choice = 82 %) to medium (53

1 %), and with large kōura showing similar preference between FlatStone and StonePile (42
2 % vs 38 %). Structure type use shifted directionally between initial and next-day location
3 (Stuart-Maxwell test: $p = 0.001$), indicating that kōura actively relocated to preferred structure
4 types overnight.

5 **Experiment 3: Wood vs Stone**

6 Kōura consistently selected FlatStone over WoodSplit, with the preference growing with body
7 size (Figure 4). At the group level, FlatStone was used significantly more than expected ($p =$
8 0.001), while WoodSplit was significantly avoided ($p = 0.004$). Only one kōura was found at
9 TankWall. The Bayesian model indicated that preference for FlatStone strengthened with body
10 size, with the posterior probability of FlatStone being the top-ranked structure increasing from
11 small (66 %) to medium (79 %), to large kōura (99 %). Large kōura showed a particularly
12 strong and credible preference for FlatStone over WoodSplit ($\beta = -2.55$, 95% CI: -4.7 to -0.9;
13 posterior probability = > 0.99), while medium kōura did not differ from small kōura. Structure
14 type use shifted directionally between initial and next-day location (Stuart-Maxwell test: $p =$
15 0.003), with kōura relocating toward FlatStone overnight.

16 **Experiment 4: Material comparison**

17 Kōura used both structure types, showing no significant preference between ConcreteBrick and
18 WoodBlock ($p = 0.137$; Figure 5), with mean group proportions of 47 % and 53 % respectively.
19 Neither structure type differed significantly from equal distribution ($p = 0.273$). 9 % of kōura
20 were found at TankWall, which is higher than in any previous experiment, likely reflecting
21 the smaller total structure size in this experiment rather than active structure avoidance. Sex
22 had no credible effect on preference between structure types ($\beta = -0.18$, 95% CI: -1.24 to 0.84;
23 posterior probability = 0.65), indicating that both structures were equally suitable refuge for
24 male and female kōura.

1 Discussion

2 Structure matters more than material

3 The consistent finding across all experiments that structural stability and cavities complexity de-
4 termined preference, regardless of whether the comparison involved material type, configuration
5 or group size, suggests that kōura, like many refuge-dependent crustaceans (Zimmer-Faust and
6 Spanier, 1987; Beck, 1995; Streissl and Hödl, 2002; Pirtle *et al.*, 2012), are primarily selecting
7 for physical shelter quality rather than material familiarity. When hole dimensions were held
8 constant between concrete and wooden perforated structures, no preference for either material
9 was detected, demonstrating that material composition alone may not drive refuge selection
10 when structure is equivalent. Neither structure type was conditioned prior to experiments,
11 meaning biofilm colonisation that occurs in natural lake environments was absent. As stone and
12 wood develop different biofilm communities (Tank and Winterbourn, 1996), material preference
13 in field settings could change. The wooden structures required a clay brick placed on top to
14 prevent floating, creating a difference with the concrete structures, though the absence of any
15 preference suggests this did not meaningfully influence the result. This material-independence
16 broke down when material affected structural properties: when flat-laid stones were compared
17 against similarly sized split wood pieces, kōura showed a clear preference for stone. The wood
18 splits, being less dense than stone, sank but created less structurally stable refuge places in the
19 sandy substrate than the stones, as larger kōura were able to move them [O.V. Raven, pers.
20 obs.].

21 In group settings, structures providing the most crevice complexity were strongly preferred,
22 with flat-laid stones accommodating the most kōura, followed by stone piles. Individual kōura
23 showed a different preference, consistently choosing the perforated wooden log over all stone
24 options, which is consistent with our first prediction that structurally complex and enclosed
25 refugia would be preferred as enclosed spaces reduce predation risk (Jurcak *et al.*, 2016). The
26 individual drilled holes in the wooden log offered protection similar to what has been found for
27 perforated brick and PVC tube shelters in response to the threat of catfish predation (Adams,
28 2007; Zhao *et al.*, 2024). The directional shift in structure type use, between initial locations
29 shortly after release and their location following a night within the tank, confirmed across
30 all experiments that kōura actively relocated overnight, supporting the interpretation that
31 next-day locations reflect genuine refuge preference rather than chance displacement.

32 The shift from the perforated wooden log in individual trials to stone structures in group trials

1 was unexpected. The wooden log contained six individual drilled holes, so capacity was not
2 limiting for the group of six kōura. The preference shift may partly reflect size mismatch
3 between fixed hole dimensions, and the range of body sizes present in each group, with only
4 medium-sized individual finding the holes well-suited, while small and large individuals were
5 better served by the range of crevices created by the stone structures (Streissl and Hödl, 2002).
6 However, a more fundamental explanation may relate to social behaviour. The wooden log
7 was the most selected structure in round 1 of Experiment 2 (17 out of 36 times), similar to
8 Experiment 1, but this preference did not persist in subsequent rounds. This shift suggests that
9 once kōura had experienced sharing the tank, they adjusted their shelter use toward structures
10 allowing closer conspecific proximity. Gregarious shelter behaviour has been documented in
11 juvenile burrowing crayfish (Hazlett *et al.*, 2007; Reynolds *et al.*, 2013; Clay *et al.*, 2017) and is
12 well established in lobsters including the New Zealand rock lobster (*Jasus edwardsii* Hutton,
13 1875) (Butler *et al.*, 1999), where conspecific chemical cues attract individuals to share refugia
14 and find refugia significantly faster, reducing predation exposure (Zimmer-Faust and Spanier,
15 1987; Childress and Herrnkind, 2001). Whether kōura exhibit similar gregarious tendencies
16 remains understudied, but the preference shift observed in this study, might suggest that social
17 context alters shelter preferences.

18 **Body size and competitive dynamics**

19 Size class consistently influenced refuge preference across experiments, but the pattern differed by
20 context. In Experiment 3, larger kōura showed a stronger preference for flat stones over wooden
21 splits than smaller individuals, consistent with size dependent shelter selection documented in
22 crayfish (Streissl and Hödl, 2002; Percival and Moore, 2010). This may reflect larger kōura
23 being more capable of asserting dominance over preferred stable refugia through competitive
24 displacement, which is consistent with the size-based dominance hierarchy documented in kōura
25 populations (Ranta and Lindström, 1993; Figler *et al.*, 2005; Stewart and Tabak, 2011). In
26 Experiment 2, where four structure types were available, small kōura showed the strongest
27 preference for flat stones while large kōura distributed more evenly between flat stones and stone
28 piles, suggesting that the large continuous surface area of the flat stones provides refuge spaces
29 proportional to smaller body dimensions (Streissl and Hödl, 2002), while larger individuals find
30 adequate crevice space in either stone configuration. The absence of sex effects in Experiments 1
31 and 4 suggests that refuge preference in kōura is driven primarily by body size and competitive
32 dominance rather than sex (Figler *et al.*, 1999). When refuge availability is limited, competitive
33 hierarchies displace smaller and recently moulted individuals from preferred refugia, leaving

1 them exposed (Ranta and Lindström, 1993; Figler *et al.*, 2005). Providing sufficient refuge
2 spaces with a range of crevice size can reduce agonistic interactions and benefit the whole
3 population (Olsson and Nyström, 2009), supporting complex structure configurations over
4 structures with fixed hole dimensions.

5 **Restoration implications**

6 With the threat from invasive predatory fish, such as catfish preying on kōura (Kusabs *et*
7 *al.*, 2026), enhancing habitat complexity may reduce top-down control on kōura populations.
8 Additional habitat as used here can provide shelter spaces and offers a promising management
9 tool, as refuge availability directly influences predation survival (Adams, 2007; Zhao *et al.*,
10 2024). The present study demonstrates that structural configuration and stability drive
11 refuge preference more strongly than material composition, with stone-based configurations
12 preferred by groups and no preference detected between materials when dimensions were equal,
13 suggesting that stone-based habitat enhancement structures warrant field evaluation in lake
14 environments.

15 The broader restoration literature supports both the structural principle and the choice of
16 natural materials. Structurally stable crevices are important for freshwater species like crayfish
17 (Huolila *et al.*, 1997; Streissl and Hödl, 2002), but also for marine environments, as reef fish,
18 marine shrimp, and lobsters have all been shown to benefit from habitat enhancement structures
19 such as artificial reefs (Spanier, 1994; Magnelia *et al.*, 2008; Hylkema *et al.*, 2020; Dickson *et*
20 *al.*, 2023). Structures made from stone and wood have been shown to outperform synthetic
21 alternatives and bare substrates (Huolila *et al.*, 1997; Magnelia *et al.*, 2008; Cooke *et al.*, 2023;
22 Dickson *et al.*, 2023), and do not contribute to plastic pollution in the environment (Cooke
23 *et al.*, 2023). Locally sourced natural materials also align with *kaitiakitanga*, which is the
24 Māori principle of environmental guardianship and preserves the natural character of culturally
25 significant lake environments (Clapcott *et al.*, 2018). Furthermore, existing habitat surveys
26 suggest that kōura are associated with coarser substrates (Kusabs *et al.*, 2015b; Olsson *et al.*,
27 2006; Jowett *et al.*, 2008), and cobbles have previously been used in stream and lake restoration
28 (Huolila *et al.*, 1997; Johnsen and Taugbøl, 2008). For in-lake deployment stone pile structures
29 are likely to be the most practical configuration, due to their elevated structure, stone piles are
30 less likely to be buried into soft substrates than flat individual rocks (Garron *et al.*, 2024). Field
31 validation in lake environments remains an essential next step, as in-lake conditions including

1 water movement, natural substrate, and predator pressure may modify kōura preferences in
2 ways not captured under controlled conditions.

3 **Conclusion**

4 Across four mesocosm experiments, structural configuration and crevice complexity consistently
5 predicted kōura shelter preference more strongly than material composition. The shift from
6 individual preference for enclosed wooden log shelters to group preference for stone configurations
7 shows that social context plays a key role in determining shelter value for kōura. Size-dependent
8 preferences further suggest that structures accommodating multiple body sizes simultaneously
9 benefit whole populations rather than only dominant individuals. Stone-based structures using
10 locally sourced materials are promising candidates for field evaluation, aligning with ecological
11 and cultural values associated with restoring this taonga species. Field validation remains
12 the essential next step for translating these findings into effective kōura habitat restoration
13 practice.

14 **Acknowledgements**

15 We thank Te Arawa Lakes Trust and Te Komiti Whakahaere for the opportunity to collect
16 and study the treasured kōura. We are grateful to Soweeta Fort-D’ath and William Anaru (Te
17 Arawa Lakes Trust), and Tihini Grant (Ngāti Pikiao).

18 **Funding statement**

19 This research was supported by the Fish Futures programme funded through a Ministry of
20 Business, Innovation and Employment grant (CAWX2101) with additional funding provided by
21 the Bay of Plenty Regional Council under the Toi huarewa Waimāori - Bay of Plenty Regional
22 Council Chair in Lake and Freshwater Science programme.

23 **Authors’ contributions**

24 Study conception and design were undertaken by O.V.R., R.H. and F.J.B. Experiments were
25 conducted by O.V.R. Data analysis was performed by O.V.R. I.A.K.K. obtained iwi consent,
26 provided local ecological knowledge and kōura specimens. Manuscript drafting was led by

1 O.V.R. All authors contributed to manuscript revision, approved the results, and the final
2 manuscript.

3 **Ethical approval**

4 This study received ethical approval by University of Waikato animal ethics committee, protocol
5 number: 1207. Additional permission to conduct experiments with kōura was granted by Te
6 Arawa Lakes Trust and Te Komiti Whakahaere.

7 **Declarations**

8 The corresponding author has declared that none of the authors has any competing interests.

9 **Data availability statement**

10 The code and derived data supporting the findings of this study are openly available on GitHub
11 (https://github.com/OlivierRaven/Koura_Mesocosm) and a rendered version of the analysis
12 is hosted at https://oliverraven.github.io/Koura_Mesocosm/. The primary raw dataset
13 is archived on Zenodo (<https://doi.org/10.5281/zenodo.20754344>). Peer review documents,
14 including the cover letter and reviewer responses, are available in the manuscript repository in
15 the interest of open and transparent science.

16 **Figures**

17 Adams SB. 2007. Direct and indirect effects of channel catfish (*Ictalurus punctatus*) on native
18 crayfishes (Cambaridae) in experimental tanks. *The American Midland Naturalist* **158**:
19 85–96. [https://doi.org/10.1674/0003-0031\(2007\)158\[85:daieoc\]2.0.co;2](https://doi.org/10.1674/0003-0031(2007)158[85:daieoc]2.0.co;2)
20 Adams SB. 2014. Crayfish use of trash versus natural cover in incised, sand-bed streams.
21 *Environmental Management* **53**: 382–392. <https://doi.org/10.1007/s00267-013-0197-3>
22 Beck MW. 1995. Size-specific shelter limitation in stone crabs: A test of the demographic
23 bottleneck hypothesis. *Ecology* **76**: 968–980. <https://doi.org/10.2307/1939360>
24 Bürkner P-C, Gabry J, Weber S, Johnson A. 2025. [brms: Bayesian regression models using](#)
25 [Stan](#).



Figure 1: Overhead photographs of experimental tank configurations for each of the four mesocosm experiments. Experiment 1 (Individual preference) and Experiment 2 (Group dynamics): four shelter types, WoodLog (tānekaha log with drilled holes), FlatStone (stones arranged flat), StonePile (stones stacked), and SingleStone (individual stones placed separately). Experiment 3 (Wood vs Stone): WoodSplit shelter (tānekaha wood splits) and FlatStone shelter placed in opposing halves of the tank. Experiment 4 (Material comparison): ConcreteBrick (perforated concrete bricks) and WoodBlock (perforated wooden blocks of identical dimensions) placed in opposing quadrants of the tank.




Figure 2: Proportion of individual kōura (Experiment 1) found at each structure type the next day, shown separately for female (white bars) and male (black bars) kōura. Bars show observed proportions with sample sizes (n) shown on bars. Letters above bars indicate results of pairwise G-tests with Bonferroni correction within each sex group, structure types sharing the same letter within a panel do not differ significantly from each other, while structure types with different letters differ significantly in use. No significant difference in overall structure preference was detected between sexes (see results).




Figure 3: Structure type preference of group kōura (Experiment 2). (A) Proportion of kōura found at each structure type at L1 across all groups and rounds. The dashed line indicates the null expectation of equal distribution (0.25). Significance symbols indicate deviation from null expectation (Wilcoxon signed-rank test, Bonferroni adjusted): ↑ used more than expected, ↓ used less than expected. Sample sizes shown in white. (B) Posterior predicted probability of structure type use by size class from Bayesian categorical mixed model, with 95 % credible intervals.




Figure 4: Structure type preference in stone versus wood comparison (Experiment 3). (A) Proportion of kōura found at each structure type at next-day location across all groups and rounds. The dashed line indicates the null expectation of equal distribution (0.5). Significance symbols as in Figure 3. (B) Posterior predicted probability of structure type use by size class from Bayesian categorical mixed model, with 95 % credible intervals. ↑ used more than expected, ↓ used less than expected. TankWall excluded from model predictions.




Figure 5: Material preference (Experiment 4). (A) Proportion of kōura found at each structure type at next-day location across all groups and rounds. The dashed line indicates the null expectation of equal distribution (0.5). (B) Posterior predicted probability of structure type use by sex from Bayesian categorical mixed model, with 95 % credible intervals. TankWall excluded from model predictions.

- 1 Butler M, MacDiarmid A, Booth J. 1999. The cause and consequence of ontogenetic changes
2 in social aggregation in New Zealand spiny lobsters. *Marine Ecology Progress Series* **188**:
3 179–191. <https://doi.org/10.3354/meps188179>
- 4 Childress MJ, Herrnkind WF. 2001. The guide effect influence on the gregar-
5 iousness of juvenile caribbean spiny lobsters. *Animal Behaviour* **62**: 465–472.
6 <https://doi.org/10.1006/anbe.2001.1760>
- 7 Clapcott J, Ataria J, Hepburn C, Hikuroa D, Jackson A, Kirikiri R, Williams E. 2018. Mātau-
8 ranga Māori: Shaping marine and freshwater futures. *New Zealand Journal of Marine and*
9 *Freshwater Research* **52**: 457–466. <https://doi.org/10.1080/00288330.2018.1539404>
- 10 Clay M, Stoeckel J, Helms B. 2017. The role of abiotic and biotic cues in burrow habitat selec-
11 tion by juvenile crayfish. *Behaviour* **154**: 1177–1196. [https://doi.org/10.1163/1568539x-](https://doi.org/10.1163/1568539x-00003463)
12 [00003463](https://doi.org/10.1163/1568539x-00003463)
- 13 Collier KJ, Parkyn SM, Rabeni CF. 1997. Koura: A keystone species. *Water and Atmosphere*
14 **5**: 18–20.
- 15 Cooke SJ, Piczak ML, Vermaire JC, Kirkwood AE. 2023. On the troubling use of plastic
16 ‘habitat’ structures for fish in freshwater ecosystems – or – when restoration is just littering
17 (D Lesbarrères, Ed). *FACETS* **8**: 1–19. <https://doi.org/10.1139/facets-2022-0210>
- 18 Dickson J, Franken O, Watson MS, Monnich B, Holthuijsen S, Eriksson BK, Govers LL, Heide
19 T van der, Bouma TJ. 2023. Who lives in a pear tree under the sea? A first look at tree reefs
20 as a complex natural biodegradable structure to enhance biodiversity in marine systems.
21 *Frontiers in Marine Science* **10**: 1213790. <https://doi.org/10.3389/fmars.2023.1213790>
- 22 Dudgeon D, Arthington AH, Gessner MO, Kawabata Z, Knowler DJ, Lévêque C, Naiman
23 RJ, Prieur-Richard A, Soto D, Stiassny MLJ, et al. 2006. Freshwater biodiversity: Im-
24 portance, threats, status and conservation challenges. *Biological Reviews* **81**: 163–182.
25 <https://doi.org/10.1017/s1464793105006950>
- 26 Engels B. 2014. XNomial: Exact goodness-of-fit test for multinomial data with fixed probabili-
27 ties. *CRAN: Contributed Packages*. <https://doi.org/10.32614/cran.package.xnomial>
- 28 Figler MH, Blank GS, Peeke HVS. 2005. Shelter competition between resident male red
29 swamp crayfish *Procambarus clarkii* (girard) and conspecific intruders varying by sex
30 and reproductive status. *Marine and Freshwater Behaviour and Physiology* **38**: 237–248.
31 <https://doi.org/10.1080/10236240500376477>
- 32 Figler MH, Cheverton HM, Blank GS. 1999. Shelter competition in juvenile red swamp crayfish
33 (*Procambarus clarkii*): The influences of sex differences, relative size, and prior residence.
34 *Aquaculture* **178**: 63–75. [https://doi.org/10.1016/s0044-8486\(99\)00114-3](https://doi.org/10.1016/s0044-8486(99)00114-3)
- 35 Garron C, Pangaud A, Taubaty M, Rolan-Meynard M, Argillier C, Raymond J, Dublon J,

- 1 Westrelin S. 2024. Towards the development of a standardized method to monitor lakeshore
2 hydromorphological restoration measures. *Aquatic Conservation: Marine and Freshwater*
3 *Ecosystems* **34**. <https://doi.org/10.1002/aqc.4152>
- 4 Gelman A, Rubin DB. 1992. Inference from iterative simulation using multiple sequences.
5 *Statistical Science* **7**: 457–472. <https://doi.org/10.1214/ss/1177011136>
- 6 Hammond KS, Hollows JW, Townsend CR, Lokman PM. 2006. Effects of temperature and water
7 calcium concentration on growth, survival and moulting of freshwater crayfish, *Paranephrops*
8 *zealandicus*. *Aquaculture* **251**: 271–279. <https://doi.org/10.1016/j.aquaculture.2005.05.032>
- 9 Hazlett BA, Lawler S, Edney G. 2007. Agonistic behavior of the crayfish *Euastacus armatus*
10 and *Cherax destructor*. *Marine and Freshwater Behaviour and Physiology* **40**: 257–266.
11 <https://doi.org/10.1080/10236240701562412>
- 12 Huolila M, Marjomäki TJ, Laukkanen E. 1997. The success of crayfish stocking in a
13 dredged river with and without artificial shelter increase. *Fisheries Research* **32**: 185–189.
14 [https://doi.org/10.1016/s0165-7836\(97\)00046-5](https://doi.org/10.1016/s0165-7836(97)00046-5)
- 15 Hylkema A, Debrot AO, Osinga R, Bron PS, Heesink DB, Izioka AK, Reid CB, Rip-
16 pen JC, Treibitz T, Yuval M, et al. 2020. Fish assemblages of three common
17 artificial reef designs during early colonization. *Ecological Engineering* **157**: 105994.
18 <https://doi.org/10.1016/j.ecoleng.2020.105994>
- 19 Johnsen SI, Taugbøl T. 2008. Add stones, get crayfish – is it that simple? *Freshwater Crayfish*
20 **16**: 47–50.
- 21 Jowett IG, Parkyn SM, Richardson J. 2008. Habitat characteristics of crayfish (*Paranephrops*
22 *planifrons*) in New Zealand streams using generalised additive models (GAMs). *Hydrobiolo-*
23 *gia* **596**: 353–365. <https://doi.org/10.1007/s10750-007-9108-z>
- 24 Jurcak AM, Lahman SE, Wofford SJ, Moore PA. 2016. Behavior of crayfish. In *Biology and*
25 *ecology of crayfish* CRC Press;
- 26 Kusabs IA, Hicks BJ, Quinn JM, Hamilton DP. 2015a. Sustainable management of freshwater
27 crayfish (kōura, *Paranephrops planifrons*) in Te Arawa (Rotorua) lakes, North Island, New
28 Zealand. *Fisheries Research* **168**: 35–46. <https://doi.org/10.1016/j.fishres.2015.03.015>
- 29 Kusabs IAK, Duggan IC, Bruere AC, Scholes P, Te Kurapa TW, Grant TM, Burdon FJ. 2026.
30 Invasive catfish linked to the decline of freshwater crayfish in two Aotearoa New Zealand
31 lakes. *Biological Invasions* **28**. <https://doi.org/10.1007/s10530-026-03852-0>
- 32 Kusabs IA, Quinn JM. 2009. Use of a traditional Maori harvesting method, the tau kōura,
33 for monitoring kōura (freshwater crayfish, *Paranephrops planifrons*) in Lake Rotoiti, North
34 Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* **43**: 713–722.
35 <https://doi.org/10.1080/00288330909510036>

- 1 Kusabs IA, Quinn JM, Hamilton DP. 2015b. Effects of benthic substrate, nutrient enrichment
2 and predatory fish on freshwater crayfish (kōura, *Paranephrops planifrons*) population
3 characteristics in seven Te Arawa (Rotorua) lakes, North Island, New Zealand. *Marine and*
4 *Freshwater Research* **66**: 631–643. <https://doi.org/10.1071/MF14148>
- 5 Lee F, Kusabs IAK, Perry GLW, MacNeil C. 2025. Identifying refugia from the syn-
6 ergistic threats of climate change and invasive species. *Web Ecology* **25**: 221–239.
7 <https://doi.org/10.5194/we-25-221-2025>
- 8 MacNeil C, Lee F, Kusabs I, Holmes R. 2025. The biter bit; scavenging behaviour
9 of native freshwater crayfish on the carrion of native and introduced fish predators
10 in Aotearoa-New Zealand. *Knowledge & Management of Aquatic Ecosystems*: 29.
11 <https://doi.org/10.1051/kmae/2025022>
- 12 Magnelia SJ, Jesus MJD, Schlechte JW, Cummings GC, Duty JL. 2008. Comparison of plastic
13 pipe and juniper tree fish attractors in a central texas reservoir.
- 14 Mazlum Y, Guner Gurlek O, Sirin S. 2017. Effect of different substrates on survival and growth
15 of juveniles of the freshwater narrow-clawed crayfish *Astacus leptodactylus* eschscholtz, 1823
16 (Decapoda, Astacidae). *Crustaceana* **90**: 1289–1302. [https://doi.org/10.1163/15685403-](https://doi.org/10.1163/15685403-00003724)
17 [00003724](https://doi.org/10.1163/15685403-00003724)
- 18 Momot WT. 1995. Redefining the role of crayfish in aquatic ecosystems. *Reviews in Fisheries*
19 *Science* **3**: 33–63. <https://doi.org/10.1080/10641269509388566>
- 20 Olsson K, Nyström P. 2009. Non-interactive effects of habitat complexity and adult crayfish on
21 survival and growth of juvenile crayfish (*Pacifastacus leniusculus*). *Freshwater Biology* **54**:
22 35–46. <https://doi.org/10.1111/j.1365-2427.2008.02089.x>
- 23 Olsson K, Stenroth P, Nyström P, Holmqvist N, McIntosh AR, Winterbourn MJ. 2006. Does
24 natural acidity mediate interactions between introduced brown trout, native fish, crayfish
25 and other invertebrates in West Coast New Zealand streams? *Biological Conservation* **130**:
26 255–267. <https://doi.org/10.1016/j.biocon.2005.12.019>
- 27 Parkyn SM, Collier KJ, Hicks BJ. 2001. New Zealand stream crayfish: Functional omnivores
28 but trophic predators? *Freshwater Biology* **46**: 641–652. [https://doi.org/10.1046/j.1365-](https://doi.org/10.1046/j.1365-2427.2001.00702.x)
29 [2427.2001.00702.x](https://doi.org/10.1046/j.1365-2427.2001.00702.x)
- 30 Parkyn SM, Meleason MA, Davies-Colley RJ. 2009. Wood enhances crayfish (*Paranephrops*
31 *planifrons*) habitat in a forested stream. *New Zealand Journal of Marine and Freshwater*
32 *Research* **43**: 689–700. <https://doi.org/10.1080/00288330909510034>
- 33 Percival D, Moore P. 2010. Shelter size influences self-assessment of size in crayfish,
34 *Orconectes rusticus*: Consequences for agonistic fights. *Behaviour* **147**: 103–119.
35 <https://doi.org/10.1163/000579509x12512685881053>

- 1 Pirtle JL, Eckert GL, Stoner AW. 2012. Habitat structure influences the survival and predator-
2 prey interactions of early juvenile red king crab *Paralithodes camtschaticus*. *Marine Ecology*
3 *Progress Series* **465**: 169–184. <https://doi.org/10.3354/meps09883>
- 4 Ramalho RO, McClain WR, Anastacio PM. 2010. An effective and simple method of temporarily
5 marking crayfish. *Freshwater Crayfish* **17**: 57–60.
- 6 Ranta E, Lindström K. 1993. Body size and shelter possession in mature signal crayfish,
7 *Pacifastacus leniusculus*. *Annales Zoologici Fennici* **30**: 125–132.
- 8 Raven OV, Burdon FJ, Kusabs IAK, Holmes R, Özkundakci D. (b) in review. [Littoral habitat](#)
9 [structure influences freshwater crayfish populations in five volcanic lakes of Aotearoa New](#)
10 [Zealand](#).
- 11 Raven OV, MacNeil C, Lee F, Holmes R, Kusabs IAK, Burdon FJ, Özkundakci D. (a) in review.
12 [Fear of the new: A predatory invader may provoke stronger non-consumptive effects on a](#)
13 [keystone native prey than a native predator](#).
- 14 Reynolds J, Souty-Grosset C, Alastair A. 2013. Ecological roles of crayfish in freshwater and ter-
15 restrial habitats. *Freshwater Crayfish* **19**: 197–218. [https://doi.org/2010.5869/fc.2013.v19-](https://doi.org/2010.5869/fc.2013.v19-2.197)
16 [2.197](#)
- 17 Richman NI, Böhm M, Adams SB, Alvarez F, Bergey EA, Bunn JJS, Burnham Q, Cordeiro J,
18 Coughran J, Crandall KA, et al. 2015. Multiple drivers of decline in the global status of
19 freshwater crayfish (decapoda: astacidea). *Philosophical Transactions of the Royal Society*
20 *B: Biological Sciences* **370**: 20140060. <https://doi.org/10.1098/rstb.2014.0060>
- 21 Shave CR, Townsend CR, Crowl TA. 1994. Anti-predator behaviours of a freshwater crayfish
22 (*Paranephrops zealandicus*) to a native and an introduced predator. *New Zealand Journal*
23 *of Ecology* **18**: 1–10.
- 24 Spanier E. 1994. What are the characteristics of a good artificial reef for lobsters? *Crustaceana*
25 **67**: 173–186.
- 26 Stewart C, Tabak J. 2011. Size-based dominance hierarchies in the New Zealand freshwater
27 crayfish (koura) *Paranephrops zealandicus*. *New Zealand Journal of Marine and Freshwater*
28 *Research* **45**: 281–285. <https://doi.org/10.1080/00288330.2011.559661>
- 29 Streissl F, Hödl W. 2002. Habitat and shelter requirements of the stone crayfish, *Austropotamo-*
30 *bius torrentium* schrank. *Hydrobiologia* **477**: 195–199. <https://doi.org/10.1023/A:1021094309738>
- 31 Takahashi K, Yamaguchi E, Fujiyama N, Nagayama T. 2019. The effects of quality of shelters
32 and prior residence on marmorkrebs (marbled crayfish). *Journal of Experimental Biology*.
33 <https://doi.org/10.1242/jeb.197301>
- 34 Tank JL, Winterbourn MJ. 1996. Microbial activity and invertebrate colonisation of wood in a
35 New Zealand forest stream. *New Zealand Journal of Marine and Freshwater Research* **30**:

1 271–280. <https://doi.org/10.1080/00288330.1996.9516714>

2 Usio N, Townsend CR. 2000. Distribution of the New Zealand crayfish *Paranephrops*
3 *zealandicus* in relation to stream physico-chemistry, predatory fish, and inverte-
4 brate prey. *New Zealand Journal of Marine and Freshwater Research* **34**: 557–567.
5 <https://doi.org/10.1080/00288330.2000.9516957>

6 Wehi PM, Lord JM. 2017. Importance of including cultural practices in ecological restoration.
7 *Conservation Biology* **31**: 1109–1118. <https://doi.org/10.1111/cobi.12915>

8 Zhao M, Feng G, Wang H, Shen C, Fu Y, Zhang Y, Zhang H, Yao Y, Chen J, Xu W.
9 2024. The influence of shelter type and coverage on crayfish (*Procambarus clarkii*) pre-
10 dation by catfish (*Silurus asotus*): A controlled environment study. *Animals* **14**: 1147.
11 <https://doi.org/10.3390/ani14081147>

12 Zimmer-Faust RK, Spanier E. 1987. Gregariousness and sociality in spiny lobsters: Implications
13 for den habitation. *Journal of Experimental Marine Biology and Ecology* **105**: 57–71.
14 [https://doi.org/10.1016/s0022-0981\(87\)80029-1](https://doi.org/10.1016/s0022-0981(87)80029-1)