

Littoral habitat structure influences freshwater crayfish populations in five volcanic lakes of Aotearoa New Zealand

Olivier V. Raven Francis J. Burdon Ian A. K. Kusabs
Robin Holmes Deniz Özkundakci

2026-06-29

Freshwater crayfish are increasingly threatened by habitat degradation, declining water quality, and invasive species. In Aotearoa New Zealand, lake populations of the endemic northern freshwater crayfish *Paranephrops planifrons* (kōura) have declined by up to 91 %. Identifying key habitat requirements is essential to improve the conservation of crayfish facing multiple threats. To assess littoral habitat associations, we surveyed shoreline habitats in five Aotearoa lakes and modelled kōura occupancy, abundance, and biomass using generalised additive mixed models. Kōura occupancy increased with shoreline habitat complexity, particularly coarser substrate types and greater riparian vegetation cover. Elevated summer surface water temperatures were negatively associated with kōura occupancy and abundance, indicating reduced habitat suitability during warm periods. pH was positively correlated with kōura abundance and biomass. Our results suggest that the composition and physical structure of littoral habitats are important drivers of kōura distribution and abundance. As climatic drivers resulting in warmer water temperatures and reduced dissolved oxygen are difficult to manage, targeted protection and restoration of coarse substrate shoreline habitats may provide a practical conservation response.

Table of contents

1	1 Introduction	3
2	2 Materials and Methods	5
3	2.1 Study location	5
4	2.2 Data collection	6
5	2.3 Data analyses	7
6	3 Results	9
7	3.1 Kōura occupancy	10
8	3.2 Kōura CPUE	11
9	3.3 Kōura BPUE	11
10	4 Discussion	12
11	4.1 Habitat associations with kōura	12
12	4.2 Fish associations as indicator of habitat quality	14
13	4.3 Associations with physico-chemical drivers	15
14	4.4 Conclusions	16
15	5 Acknowledgements	17
16	5.1 Funding statement	17
17	5.2 Authors' contributions	17
18	5.3 Ethical approval	18
19	5.4 Declarations	18
20	5.5 Data availability statement	18
21	6 Tables	18
22	7 Figures	22
23	8 Supplementary information	22

1 Introduction

Freshwater crayfish (Astacidea) play a vital role in freshwater ecosystems (Momot, 1995). They are ecosystem engineers, exerting substantial influence over key ecological processes, including nutrient cycling, organic matter decomposition and bioturbation (Momot, 1995; Collier et al., 1997; Parkyn et al., 2001). They are keystone species due to their broad trophic niche as predators and detritivores, and their ability to modify habitats through sediment excavation for burrow construction, removal of macrophytes, and alteration of macroinvertebrate communities (Statzner et al., 2003; Creed & Reed, 2004; Usio & Townsend, 2004; Albertson & Daniels, 2018). Nearly one-third of the world’s crayfish species are currently threatened with extinction (Richman et al., 2015). Despite their ecological importance and imperilled status, crayfish are frequently overlooked in restoration and conservation initiatives (Taylor et al., 2019; Danilović et al., 2022).

In Aotearoa New Zealand (hereafter Aotearoa), the endemic northern freshwater crayfish *Paranephrops planifrons* White, 1842, also known by its Māori name kōura, inhabits a wide range of freshwater environments, including rivers, streams and lakes across both main islands (Hopkins, 1970). Beyond its ecological role, kōura hold deep cultural significance, having historically been an important food source and trade commodity for iwi (Māori communities; Hiroa, 1921; Kusabs & Quinn, 2009)). However, kōura populations are increasingly threatened by climate change, invasive species and declining water quality (Kusabs et al., 2015a; Lee et al., 2025). These declines are likely the result of multiple interacting environmental stressors, including declining water quality, deoxygenation and the introduction and consequent impacts of non-native species (Kusabs & Quinn, 2009). For example, in Lakes Rotorua and Rotoiti, located in the Bay of Plenty region of Aotearoa’s North Island, kōura populations have declined by 76 % and 91 %, respectively, over the past two decades (Kusabs et al., 2026).

In 2016, the invasive North American brown bullhead catfish (*Ameiurus nebulosus* Lesueur, 1819) was confirmed to be present in Lakes Rotorua and Rotoiti (Grayling, 2020), where it now occurs in large numbers and preys heavily on kōura (Francis, 2019; Kusabs et al., 2026). To reduce predation risk, kōura typically seek refuge in deeper lake habitats during the day and migrate to the more productive shallow areas at night to forage (Devcich, 1979). However, summer lake stratification has led to oxygen depletion in deeper waters, forcing kōura to spend

24 more time in shallow habitats where predation risk is greater (Dedual, 2002). Climate change
25 is expected to intensify these effects by strengthening thermal stratification and increasing the
26 extent and duration of anoxic conditions in bottom waters as water temperatures rise (Woolway
27 et al., 2021; Zhang et al., 2024). In addition, dense beds of invasive macrophytes may obstruct
28 kōura migration pathways between deep and shallow habitats (Hessen et al., 2004; Kusabs &
29 Quinn, 2009) and further reduce oxygen availability during seasonal macrophyte die-off events
1 (Vincent et al., 1984; Hessen et al., 2004; Hamilton et al., 2005).

2 Declines in kōura populations have cascading ecological impacts and implications for Māori
3 cultural practices, including reduced opportunities for customary harvest and disruption of
4 traditional activities (Kusabs et al., 2015a; Kusabs & Quinn, 2009; Kusabs et al., 2026). To
5 effectively conserve and restore populations requires an improved understanding of where kōura
6 occur within lakes and what habitats are most important for sustaining healthy populations.
7 Previous research in stream environments has shown that kōura preferentially occupy struc-
8 turally complex habitats characterised by bank undercuts, woody debris, leaf litter, boulders
9 and low flow velocities (Usio & Townsend, 2000; Parkyn & Collier, 2004; Jowett et al., 2008;
10 Parkyn et al., 2009). In contrast, habitats used by kōura in lakes remains poorly understood,
11 with previous studies largely focused on deeper offshore habitats (Kusabs et al., 2015b; Devcich,
12 1979; Kusabs & Quinn, 2009). These studies suggest that benthic substrate characteristics may
13 be more important than nutrient enrichment or predator abundance — specifically non-native
14 rainbow trout (*Oncorhynchus mykiss* Walbaum, 1792) — in determining kōura distribution
15 (Kusabs et al., 2015b). The littoral zone of lakes has received little attention, despite it being
16 the most productive zone and supporting key life history processes of kōura (Strayer & Findlay,
17 2010), including nocturnal foraging and use by ovigerous females prior to juvenile release
18 (Devcich, 1979). Consequently, the littoral zone is likely to represent an important habitat
19 within lakes for kōura.

20 In this study, we examine which littoral habitat characteristics are associated with kōura
21 occurrence, abundance and biomass across five Rotorua Te Arawa Lakes. We hypothesised
22 that structurally diverse littoral habitats, including coarse substrates and native macrophytes,
23 would be positively associated with kōura occurrence and abundance, as they provide physical
24 refuge and foraging opportunities. Conversely, invasive submerged macrophytes were expected
25 to have negative associations with kōura as these dense macrophytes hinder movement and
26 reduce native biodiversity by displacing native macrophytes, altering habitat structure, and
27 modifying ecological processes. We further expected that kōura abundance would be lower in
28 lakes where catfish are present, as catfish are a known predator on benthic invertebrates and

native fauna. By quantifying these habitat associations within the littoral zone, this study aims to inform targeted habitat protection and restoration strategies for the conservation of this culturally and ecologically important species.

2 Materials and Methods

2.1 Study location

Study sites were selected in five lakes located in the Rotorua Te Arawa Lakes area in the North Island of Aotearoa. The Lakes Rotorua, Rotoiti, Rotoehu, Rotomā, and Ōkāreka are located on a warm- temperate volcanic plateau (280-355 m) where winters are mild and year-round temperatures typically varies from 4°C to 24°C. The lakes span a wide range of morphometric and trophic conditions from shallow polymictic eutrophic systems (Rotoehu, maximum depth 14 m) to deep stratified oligotrophic lakes (Rotomā, maximum depth 83 m). Shorelines are characterised by a mix of native riparian vegetation, exotic pasture, and low-density residential development, with several lakes also supporting stands of emergent macrophytes. Detailed lake characteristics and shoreline composition are summarised in Table 1. All five lakes have populations of kōura, and brown bullhead catfish was present in Lakes Rotorua and Rotoiti at time of sampling (Kusabs et al., 2026).

To survey these lakes, a stratified-random survey design was developed. The entire accessible shoreline of each lake was first mapped using aerial photos and then ground truthed and classified into dominant habitat types. Classification used the following criteria: a habitat was considered dominant “rocky” if the combined percentage of bedrock, boulders, and cobble exceeded 25 % by surface area; “sandy” if sand was the dominant sediment (50 %) and rocky conditions were not met; “muddy” if soft mud or organic matter was dominant and rocky/sandy conditions were not met; and “emergent macrophyte” if emergent native macrophytes exceeded 25 % of the shoreline cover by surface area. Shoreline sites exceeding two meters in depth or classified as geothermal were excluded, as these conditions were either unsuitable for kōura or inaccessible for field sampling.

For each lake, the total shoreline length of each habitat type within this sample frame was calculated in a projected coordinate system (New Zealand Transverse Mercator 2000; EPSG:2193). To ensure balanced representation of littoral habitat types across systems and to avoid confounding lake identity with sampling effort, an equal number of sites ($n = 12$) was surveyed in

each lake, with sites allocated proportionally among the dominant shoreline habitat types. All selected sites were fully accessible for fieldwork, and therefore an oversample was not required. Within each habitat type, site locations were generated using the Generalized Random Tessellation Stratified (GRTS) design (Stevens & Olsen, 2004) executed using the spsurvey package in R (Dumelle et al., 2023; R Core Team, 2025), producing unbiased random coordinates without enforcing fixed spacing between sites. The final coordinates of the 12 random survey sites per lake were extracted directly from the GRTS output (Figure 1).

2.2 Data collection

In total, 60 sampling sites were sampled once in the Austral spring/summer from October 2024 to February 2025. Each sampling site measured 15×10 m and extended from the shoreline into the lake. At locations with unstable lake margins or depths exceeding 1 m, observations and measurements were conducted from a boat. At each site, physical, chemical and biological parameters were recorded (Table 2).

Physical characteristics were assessed using both visual estimates and field measurements. Substrate composition was visually estimated with an underwater viewing chamber, as water clarity was sufficiently clear at all sites, using a standardised classification scheme, categorising bedrock, boulders (>256 mm), cobble (64–256 mm), gravel (2–64 mm), sand (0.063–2 mm), mud (<0.063 mm), and organic matter. A substrate index was then calculated from these estimates using a formula adapted to include fine organic matter and mud (Jowett & Richardson, 1990; Jowett et al., 2008; Harding et al., 2009):

$$\text{Substrate index} = 0.08 \times (\text{bedrock}) + 0.07 \times (\text{boulders}) + 0.06 \times (\text{cobble}) + 0.04 \times (\text{gravel}) + 0.03 \times (\text{sand}) + 0.02 \times (\text{organic matter})$$

Larger substrate classes were assigned higher weights because coarse substrates such as cobble and boulders create greater interstitial pore spaces and stable structures that kōura use as refuge from predators. Coarse substrate availability has been shown to mediate crayfish vulnerability to predation and directly influence crayfish abundance in lakes (Usio & Townsend, 2000; Nyström et al., 2006), whereas fine substrates such as silt and mud alone offer limited physical refugia. Fine substrates were assigned near-zero coefficients (0.01–0.03), making their contribution to the index minimal relative to coarse substrates (0.06–0.08), consistent with the positive-weighting structure of the original index (Jowett & Richardson, 1990). Riparian

vegetation, overhanging tree cover, and wood debris was visually estimated as percentage cover within each site, along with emergent, submerged, and turf vegetation cover. To assess proximity to deeper waters, slope was calculated as horizontal distance from the shoreline to the 5 m depth contour derived from bathymetric maps. Chemical parameters included water temperature ($^{\circ}\text{C}$), dissolved oxygen (mg L^{-1}), and conductivity ($\mu\text{S cm}^{-1}$), measured with a YSI ProSolo meter (Pro2030 Dissolved Oxygen and Conductivity Meter, YSI Inc., Yellow Springs, OH, USA) and pH was determined using a handheld pH meter (EC-PCTestr35, Eutech Instruments Pte Ltd, Paisley, UK).

Catch data for kōura and fish were collected using two single-winged, double-throated fyke nets (600 mm high D-shaped entrance hoop) and two collapsible bait traps baited with trout flesh (480×250 mm; two 50 mm entrance openings). Fyke nets were placed perpendicular to the shoreline at 10 m intervals, with bait traps positioned between them. All equipment was deployed simultaneously at each site for a 24-h period. The use of two fyke nets and two baited bait traps per site was intended to maximise detection probability across habitat types. Captured kōura were measured for orbital carapace length (OCL) using a digital calliper (0.01 mm precision; 150 mm) and for wet mass using a digital scale (± 1 g; 5 kg capacity, Anko Brand, Kmart Group, Mulgrave, VIC, Australia). Catfish and other fish species, including eels (*Anguilla* spp.), goldfish (*Carassius auratus* (Linnaeus, 1758)), common smelt (*Retropinna retropinna* (Richardson, 1848)), kōaro (*Galaxias brevipinnis* Günther, 1866), rainbow trout, common bully (*Gobiomorphus cotidianus* McDowall & Fulton, 1978), and mosquitofish (*Gambusia affinis* (S. F. Baird & Girard, 1853)), were recorded for abundance and measured for total length, and weighed for wet mass. Catch per unit effort (CPUE) and biomass per unit effort (BPUE) were calculated for all kōura and fish based on the number of nets and traps used at each site (Kusabs et al., 2015b). Missing kōura weights due to failure of scale ($n = 81$) were estimated using a $\log -\log$ length-weight regression fitted to individuals with both measurements ($n = 240$), following standard approaches for weight length relationships in aquatic organisms (Figure 6) (Froese, 2006). Predicted values were back-transformed with a lognormal bias correction factor ($10^{\wedge\{0.5 \wedge 2\}}$) (Quinn & Keough, 2002).

$$\log_{10}(\text{Weight [g]}) = -2.938 + 2.864 \times \log_{10}(\text{OCL length [mm]})$$

2.3 Data analyses

Because each site was sampled only once, imperfect detection could not be estimated separately from occupancy. Presence–absence data therefore reflect observed detections at the time of sampling, and sites where kōura were not captured were treated as absences in the models. As a result, inference focuses on relative associations between kōura occurrence and environmental or biotic predictors, rather than on estimating true occupancy probabilities.

Kōura presence or absence, CPUE, and BPUE were analysed using mixed-effects models to account for non-independence among sampling sites (Bolker et al., 2009). To test for differences among lakes, kōura presence was modelled using a binomial generalised linear mixed model (GLMM; logit link). CPUE and BPUE were modelled using Tweedie GLMM (log link), which is appropriate for continuous non-negative data with high frequency of zeros, as common in ecological count and biomass data (Shono, 2008). These models were fitted using the glmmTMB package (Brooks et al., 2017), with habitat type included as a random intercept. To identify environmental and biotic predictors associated with kōura distribution and abundance, GAMMs were implemented with the mgcv package (Wood, 2004, 2017). To reduce multicollinearity, predictors were screened using Variance Inflation Factors (VIFs) calculated from linear models fitted to the fixed-effect design matrix. A conservative threshold of $VIF > 5$ was applied (Quinn & Keough, 2002; Kutner et al., 2005). Across all model sets, dissolved oxygen expressed as percentage saturation (DO %) exceeded this threshold and was excluded from further analyses, while dissolved oxygen concentration (DO mg L⁻¹) was retained.

For occupancy analyses, kōura presence–absence was modelled using a binomial GAMM with a logit link function (Zuur et al., 2009; Wood, 2017). For CPUE and BPUE, a hurdle modelling approach was adopted due to the high frequency of zero observations in the data. In this approach, occurrence (zero vs non-zero values) was modelled using a binomial GAMM with a logit link, followed by modelling of positive CPUE or BPUE values using Gamma GAMMs with a log link. This framework allows occurrence and positive abundance to be modelled separately and is appropriate for datasets characterised by many zero observations. Hurdle models are commonly used with freshwater crayfish catch data, which often possess these features (Rosewarne et al., 2013). Environmental and biotic predictors considered included substrate index, slope to 5 m depth, riparian vegetation, overhanging trees, wood cover, aquatic vegetation classes (emergent native, submerged native, submerged non-native, and emergent non-native vegetation), water temperature (°C), dissolved oxygen (mg L⁻¹), pH, specific conductivity (µS cm⁻¹), and the presence of catfish, eels, goldfish, and common smelt.

Continuous predictors were centred and standardised prior to modelling to improve convergence and facilitate comparison of effect sizes and were modelled as smooth terms using thin plate regression splines ($bs = 'ts'$), which apply additional shrinkage to penalise overly complex relationships and can shrink smooth terms fully to zero during selection (Wood, 2017). Basis dimensions (k) were set based on the number of unique values per predictor and adjusted where necessary to balance flexibility and parsimony. Binary predictors (fish presence variables) were included as parametric terms.

Model selection was performed using maximum likelihood estimation with stepwise backward elimination, sequentially removed and the model refitted until all remaining terms met the retention threshold ($p = 0.05$). Reduced models were compared with their respective full models using likelihood ratio tests and Akaike Information Criterion (AIC) scores. Final selected models were refitted using restricted maximum likelihood (REML), which accounts for the degrees of freedom associated with fixed effects and provides less biased estimates of variance components, resulting in more stable smoothing parameter estimates in models with random effects. Lake identity was included as a random effect in all GAMMs to account for spatial clustering and non-independence among observations and was not subject to the variable selection procedure (Zuur et al., 2009).

Model diagnostics were performed to evaluate model fit and assumptions. Residuals were inspected to assess homogeneity of variance and normality (Zuur et al., 2009). Concurvity, the degree to which smooth terms can be approximated by other smooths in the model, was assessed to identify potential instability in smooth term estimates (Wood, 2017). Basis dimension checks using effective degrees of freedom (edf) were performed to ensure smoothing terms had sufficient flexibility to capture underlying patterns without overfitting (Zuur et al., 2009). For binomial components, model discrimination was evaluated using Receiver Operating Characteristic and Area Under the Curve (ROC/AUC), where values approaching 1 indicate perfect discrimination between presences and absences (Fielding & Bell, 1997). For positive-only abundance models, predicted versus observed values were compared to assess how well the model reproduces observed CPUE and BPUE values across the range of the data.

To provide context for the association between catfish presence and kōura metrics detected in the GAMM, a descriptive comparison of kōura presence rate, CPUE, and BPUE was conducted across catfish-present and catfish-absent sites, and across lakes with and without catfish.

3 Results

A total of 321 kōura were captured across 33 of the 60 surveyed sites within the five Rotorua Te Arawa lakes. Kōura OCL ranged from 7–53 mm and individual body mass from 0.2–87 g. Kōura were detected in all five lakes; however, abundance varied among lakes, with total counts ranging from 19 to 112 individuals. Mixed-effects models indicated a significant effect of lake on kōura occupancy (binomial GLMM: likelihood ratio test, $\chi^2 = 13.76$, $p = 0.008$), CPUE (Tweedie GLMM: $\chi^2 = 12.60$, $p = 0.013$), and BPUE (Tweedie GLMM: $\chi^2 = 9.66$, $p = 0.047$; Figure 2).

The five lakes differed substantially in environmental and biotic characteristics (Table 3). Physical habitat structure varied among lakes, with differences in substrate index, nearshore slope, and littoral vegetation cover. Emergent macrophytes were absent from Lake Rotorua but present in the other lakes, particularly Ōkāreka, while submerged macrophyte cover at the shore was rare in Rotorua and Rotoiti but more extensive in Rotoehu and Rotomā. Water chemistry also differed markedly among lakes. Surface water temperatures were highest in Rotorua and lowest in Rotomā and Ōkāreka, while pH ranged from circum-neutral conditions in Rotorua to more alkaline conditions in Rotoehu and Rotoiti. Specific conductivity showed strong contrasts, being highest in Rotoehu and lowest in Ōkāreka. Dissolved oxygen concentrations in the littoral zone varied less among lakes but were generally lowest in Rotorua. Fish populations differed between lakes, reflecting invasion history. Catfish were detected only in Lakes Rotorua and Rotoiti, whereas goldfish occurred in all lakes but were most abundant in Rotoehu. Common bullies were found in every survey location. Trout were found sporadically in Rotorua, Rotoiti and Ōkāreka and kōaro were only caught from Ōkāreka.

3.1 Kōura occupancy

The full binomial GAMM for kōura occupancy explained 65 % of the deviance ($AIC = 51$; adj. $R^2 = 0.647$). Stepwise ML reduction ($\alpha = 0.05$) retained riparian vegetation, substrate index, temperature, and specific conductivity as smoother terms, lake identity as a random effect, and the presence of goldfish (negative) and catfish (positive) as parametric terms. The final model refitted using REML explained 57.9 % of the deviance (adj. $R^2 = 0.592$) and retained positive influences of riparian vegetation ($p = 0.026$), substrate index ($p = 0.001$), specific conductivity ($p = 0.002$) and a negative influence of temperature ($p < 0.001$), as significant smoother terms (Figure 3 a). The random effect of lake identity was not statistically significant ($p = 0.622$),

indicating limited residual variation between lakes after accounting for environmental and biotic predictors. Among the parametric covariates, goldfish had a significant negative relationship ($p = 0.017$), and catfish presence showed a significant positive relationship ($p = 0.014$). Model diagnostics showed no evidence of overfitting ($\text{edf} < k$) and good discrimination ($\text{AUC} = 0.946$; Figure 3 b). Concurvity diagnostics revealed high collinearity between specific conductivity and lake identity (worst-case = 1.00, observed = 0.94), and between temperature and lake identity (worst-case = 0.95, observed = 0.88), suggesting both predictors partly reflect between-lake differences. Residual diagnostics showed no systematic patterns in variance across fitted values, consistent with acceptable model fit for a binomial GAMM. The calibration plot indicated slight under-prediction at low probabilities and slight over-prediction at mid-range probabilities (Figure 3 c). The binomial components of the CPUE and BPUE hurdle models produced the same structure and effect directions as the occupancy model, so only the positive Gamma components are reported here.

3.2 Kōura CPUE

The full Gamma GAMM for positive CPUE explained 48.4 % of the deviance ($\text{AIC} = 122$; adj. $R^2 = 0.19$; $n = 33$). Stepwise selection ($\alpha = 0.05$) reduced the model to temperature, pH, and emergent native macrophytes as smoother terms, lake identity as a random effect, and the presence of common smelt (negative) as a parametric effect. This reduced model achieved a lower AIC (115) and explained 47.3 % of the deviance (adj. $R^2 = 0.194$). Likelihood ratio tests indicated that the reduced model was statistically equivalent to the full model ($p = 0.950$), and it was therefore selected as the final CPUE model.

The final REML-fitted model explained 47.3 % of the deviance (adj. $R^2 = 0.195$) and retained positive influence of pH ($p < 0.001$) and negative influence of temperature ($p = 0.019$) and emergent native macrophytes ($p < 0.001$) as significant smooth predictors (Figure 4 a). The lake identity random effect was not significant ($p = 0.482$), indicating minimal additional between lake variation once environmental covariates were included. Presence of common smelt had a significant negative parametric relationship ($p = 0.041$). Concurvity diagnostics indicated strong correlation between temperature and pH, but both variables were retained because they were ecologically interpretable and contributed significantly to model fit. Model diagnostics showed no evidence of overfitting ($\text{edf} < k$), and predictive accuracy was moderate (Figure 4 b). Residual diagnostics showed no systematic patterns in residual variance across fitted values.

27 The model tended to slightly over-predict very low CPUE values and under predict the highest
28 observations.

29 3.3 Kōura BPUE

30 The full Gamma GAMM for positive BPUE explained 97.4 % of the deviance ($AIC = 266$; adj.
1 $R^2 = 0.76$; $n = 33$). Stepwise ML selection ($\alpha = 0.05$) retained pH, emergent native macrophytes,
2 and the lake identity random effect. The resulting reduced model had a higher AIC (318) and
3 explained 26 % of the deviance (adj. $R^2 = 0.141$). As the full model was overfitting the data
4 the reduced model was chosen for inference. The final REML-fitted BPUE model explained
5 25.9 % of the deviance (adj. $R^2 = 0.14$) and retained a positive influence of pH ($p = 0.002$)
6 and a negative influence of emergent native macrophytes ($p = 0.041$) as significant, near-linear
7 smoothing (Figure 5 a). The emergent macrophyte effect, while statistically significant, was
8 close to the threshold and should be interpreted with caution given the distributional limitations
9 of the model. The lake identity random effect was not significant ($p = 0.607$), indicating minimal
10 additional between lake variation after accounting for environmental predictors. Diagnostic
11 checks showed appropriate basis dimensions ($edf < k$), stable convergence, and acceptable levels
12 of observed concurvity, though minor departure from distributional assumptions was noted
13 at the lower tail, likely reflecting a small number of high-biomass sites. Predictive accuracy
14 was moderate (Figure 5 b), with a tendency for the model to under predict the highest BPUE
15 values.

16 4 Discussion

17 We found that local and site-scale habitat characteristics were consistently associated with
18 kōura occurrence and abundance across the five surveyed lakes. Although initial mixed-effects
19 models identified significant differences in kōura occupancy, Catch Per Unit Effort (CPUE),
20 and Biomass Per Unit Effort (BPUE) among lakes, these differences were no longer evident
21 once local environmental and biotic predictors were incorporated into the Generalised Additive
22 Mixed-Models (GAMMs). However, interpreting the non-significance of lake identity in the final
23 models requires caution. In the occupancy model, both temperature and specific conductivity
24 showed near-perfect collinearity with lake identity, suggesting these predictors likely absorbed
25 between-lake differences rather than capturing purely within-lake habitat associations. Given
26 that predictors were often correlated at the site scale, the GAMMs are best interpreted as

identifying habitat associations rather than mechanistic drivers of kōura distribution. The predictors substrate index and riparian vegetation cover, which varied considerably within lakes, likely reflect genuine within-lake habitat associations. Consequently, targeted management actions that improve local habitat structure and address key biotic pressures at finer spatial scales have the potential to deliver local gains, particularly where whole-lake interventions are not feasible or where local habitat degradation is the dominant stressor.

4.1 Habitat associations with kōura

Kōura occurrence was consistently associated with coarser substrate types and greater riparian vegetation cover. Substrate heterogeneity was moderately negatively correlated with the substrate index ($r = -0.48$), indicating that the index reflects dominance of coarse substrates rather than proportional mixing of particle sizes. Higher substrate index values therefore reflect the presence of large particles like bedrock, boulders and cobbles, which create large interstitial pore spaces that function as physical refugia, providing shelter from predators (Kusabs et al., 2015b; White & Irvine, 2003; Haschenburger & Roest, 2009). The relationship between substrate size and benthic invertebrates is not linear (Quinn & Hickey, 1990), where larger pore spaces between coarse particles may favour adult kōura, smaller interstitial spaces within mixed substrates may provide refugia for juveniles, suggesting that heterogeneous substrates support kōura across life stages. However, mixed substrates containing both coarse and fine particles can reduce pore space availability as smaller particles infill void spaces between larger ones (Haschenburger & Roest, 2009), meaning that dominance of coarse substrate rather than substrate heterogeneity is the more relevant driver of refugia availability for adults. Additionally coarse substrates may facilitate burrowing by kōura at the margins of rocky areas where softer sediments are accessible, providing an additional form of refuge (Usio & Townsend, 2000). Similar preferences for coarse substrates have been reported in stream environments (Olsson et al., 2006; Jowett et al., 2008), although the optimal substrate size may differ between habitat types. In streams, kōura show a non-linear response to substrate size with an optimum around 130 mm, with probability of occurrence declining at boulder substrates larger than 256 mm, likely because larger substrates are associated with higher flow velocities that kōura actively avoid (Jowett et al., 2008). Coarse particles in streams are also relatively stable under typical flow conditions, as their size reduces the likelihood of transport during high flows. In lakes, where flow is minimal, kōura may preferentially use larger substrates such as boulders, as this constraint is removed and larger particles create more accessible interstitial refugia. However, substrate stability in these volcanic lakes remains relevant in wave-exposed littoral

25 zones, as pumice substrates, being lighter than other lithologies of similar size, are more likely
26 to redistribution during wave action, potentially reducing the stability of interstitial refugia.
27 Although fyke net deployment is more challenging in highly structured rocky habitats and may
28 lead to conservative abundance estimates, the consistent positive association with the substrate
29 index across all five lakes highlights the importance of coarse littoral substrates for kōura.

30 This refugia mechanism also helps explain why emergent macrophyte habitats were negatively
31 associated with kōura abundance and biomass despite providing vertical structure. Emergent
32 native macrophytes, including raupō (*Typha orientalis* C.Presl) and giant spike rush (*Eleocharis*
1 *sphacelata* R.Br.), often form in relatively homogeneous habitats, characterised by fine sediments
2 and high organic matter accumulation in sheltered littoral areas (Chambers, 1987; Lishawa et al.,
3 2023). Such conditions may limit the availability of effective refuges, reduce foraging efficiency,
4 and increase oxygen demand associated with organic decomposition, with dissolved oxygen also
5 likely to exhibit greater unmeasured daily variability within macrophyte stands (Bunch et al.,
6 2010; Schrank & Lishawa, 2019). Therefore, rocky shorelines maintain accessible interstitial
7 pore spaces that function as refugia, whereas macrophyte-dominated habitats accumulate fine
8 sediments that infill potential void spaces and reduce refugia availability. However, the negative
9 association between emergent native macrophytes and kōura abundance and biomass should be
10 interpreted with caution. Only 15 of 60 sites had emergent macrophyte cover and in the CPUE
11 and BPUE models this was 7 of 33 positive sites, meaning model estimates for this predictor
12 carry greater uncertainty than the smooth term confidence intervals suggest. That a consistent
13 negative effect was nonetheless detected across both CPUE and BPUE models suggests the
14 relationship is ecologically meaningful despite the limited representation of this habitat type in
15 the dataset.

16 4.2 Fish associations as indicator of habitat quality

17 Associations between kōura abundance and the presence of certain fish species further support
18 the importance of habitat-mediated processes rather than direct biotic interactions. Gold-
19 fish presence was negatively associated with kōura occupancy, and common smelt presence
20 was negatively associated with kōura CPUE. Goldfish are typically associated with warmer,
21 vegetated littoral habitats with finer substrates (Fry & Hart, 1948; Ford & Beitingger, 2005),
22 conditions that were also unfavourable for kōura in this study. Goldfish presence is therefore
23 best interpreted as an indicator of habitat conditions that are unsuitable for kōura rather
24 than as a direct driver of kōura distribution. Similarly, common smelt are primarily pelagic

25 (Rowe, 1993), and their association with reduced kōura abundance is unlikely to reflect direct
26 interactions. Instead, this pattern may reflect broader lake-scale trophic or environmental
27 gradients, such as increased predation pressure mediated through shared predators (Schoen et
28 al., 2015).

29 A weak positive association between catfish presence and kōura occupancy was detected in
30 the GAMM, although this pattern should be interpreted cautiously. Catfish were restricted to
31 lakes Rotorua and Rotoiti, occurring at only 6 of 24 sites within these two lakes. Examining
32 this association across multiple spatial scales reveals a scale-dependent pattern consistent with
1 (Levin, 1992), who demonstrated that ecological relationships could shift direction depending on
2 the scale of analysis. At the lake scale, kōura presence, CPUE, and BPUE were all substantially
3 lower in catfish-invaded lakes than in catfish-free lakes, suggesting a negative association. At
4 the site scale within invaded lakes, catfish-present sites also had lower kōura presence rates and
5 abundance than catfish-absent sites. The positive parametric term in the GAMM therefore likely
6 resulted from habitat covariate adjustment, where catfish and kōura co-occur in coarse substrate
7 littoral habitats that support higher benthic invertebrate biomass and historically higher kōura
8 densities (Barnes, 1996; Barnes & Hicks, 2003). Importantly, the positive GAMM term does
9 not preclude strong negative impacts of catfish during earlier stages of invasion; instead, current
10 distributions may represent a residual distribution following substantial historical declines, with
11 kōura persisting at reduced densities in habitats where stable coexistence is possible (Kusabs et
12 al., 2026). Overall, fish associations observed in this study appear to reflect shared responses to
13 habitat structure and productivity rather than direct biotic regulation of kōura populations.

14 4.3 Associations with physico-chemical drivers

15 Elevated summer water temperatures were negatively associated with kōura abundance in
16 both occupancy and CPUE models, indicating that thermal conditions impose an important
17 constraint on littoral habitat suitability. Lakes were surveyed at different times across the
18 spring-summer period, so between-lake temperature differences partly reflect when each lake
19 was sampled, not only differences in thermal regime. Kōura are known to be temperature-
20 sensitive, with activity increasing above approximately 10 °C and temperatures exceeding ~21
21 °C considered thermally stressful, and probably leading to aestivation (Devcich, 1979; Angilletta
22 et al., 2004; Hammond et al., 2006; Hesni et al., 2009). During the study period, surface water
23 temperatures in several lakes exceeded this threshold, including a mean summer temperature

24 of 23.15 °C in Lake Rotorua, conditions likely to reduce suitable habitat availability during
25 warm periods.

26 In stratified lakes, elevated surface temperatures may further constrain kōura habitat by
27 intensifying thermal stratification and limiting oxygen availability in deeper waters ([Wetzel,](#)
28 [2001](#)). Lake Rotoiti and Ōkāreka, for example, experience extended stratification and reduced
29 bottom-water oxygen concentrations during the summer periods ([Westernhagen et al., 2010](#)),
30 conditions that may restrict kōura to warmer nearshore or surface habitats where thermal stress
31 is also elevated. Although vertical movements and depth-specific habitat use were not assessed
32 directly, the negative association between temperature and kōura abundance observed here is
1 consistent with broader constraints on habitat suitability arising from the combined effects of
2 thermal stress and reduced oxygen availability during prolonged stratification, rather than from
3 surface temperature alone. Importantly, this temperature effect was detected independently of
4 littoral habitat structure, suggesting that elevated summer temperatures may compress kōura
5 into a narrower subset of thermally and structurally suitable microhabitats, where local habitat
6 characteristics may provide limited buffering but also increase vulnerability to predation and
7 other stressors.

8 Because sampling was restricted to the spring/summer period, seasonal shifts in temperature
9 tolerance and habitat use could not be evaluated. Nevertheless, projected climate warming is
10 expected to increase surface water temperatures and extend the duration of thermal stratification
11 in lakes, thereby intensifying summer thermal stress in shallow littoral zones and further reducing
12 suitable kōura habitat during summer and autumn ([Jane et al., 2021](#)).

13 In contrast to temperature effects on occurrence and abundance, water chemistry variables
14 were more strongly associated with kōura abundance and biomass. They increased with
15 higher pH and specific conductivity, with pH showing positive associations in both CPUE and
16 BPUE models and conductivity positively associated with occupancy. Elevated pH values
17 indicate carbonate-rich conditions that may enhance calcium availability, which is essential
18 for exoskeleton formation, moulting, and growth in kōura ([Hammond et al., 2006](#)). Further,
19 specific conductivity reflects the overall ionic composition of the water, which in volcanic lake
20 systems may indicate differences in catchment geology and geothermal inputs, though given its
21 near-perfect collinearity with lake identity the mechanistic interpretation of this association
22 remains uncertain. Favourable chemical conditions are more likely to influence the amount
23 of biomass that littoral habitats can support, rather than acting as strict determinants of
24 kōura presence. This interpretation is consistent with physiological constraints associated

25 with calcification and moulting, processes that become increasingly demanding as individuals
26 increase in size ([Greenaway, 1985](#)).

27 4.4 Conclusions

28 Our study found that local littoral habitat complexity, particularly coarse substrate availability
1 and riparian vegetation cover, were consistently associated with kōura occurrence and abundance
2 in nearshore habitats. Elevated summer water temperature was negatively associated with
3 kōura occurrence and abundance, consistent with thermal sensitivity of kōura, though this
4 association partly reflects between-lakes differences and should be interpreted cautiously.
5 Together, these findings suggest that both local habitat complexity and thermal conditions
6 shape kōura distribution within the littoral zone. This study contributes to understanding of
7 how habitat complexity structures influence freshwater crayfish populations and demonstrates
8 that not all forms of structural complexity equally support biodiversity. Different forms
9 of structural complexity likely support different species and ecological functions, meaning
10 that maintaining diversity of habitat types within the littoral zone is important for broader
11 biodiversity conservation ([Meerhoff & González-Sagrario, 2022](#)). As shoreline modification,
12 sedimentation, and invasive macrophyte expansion continue to reduce variation and structural
13 diversity in littoral habitats, understanding which components of habitat complexity are
14 functionally important for target species becomes critical for effective restoration planning.

15 Climate warming is expected to impose increasing constraints on freshwater crayfish populations
16 through rising summer water temperatures, prolonged periods of thermal stratification, and
17 intensified interactions with non-native species that are better adapted to warmer conditions
18 ([Capinha et al., 2013](#); [Lee et al., 2025](#)). While such broad-scale drivers are difficult to
1 manage directly, local habitat conditions represent a tractable management lever. The strong
1 association between kōura occurrence and coarse substrate availability suggests that an absence
2 of interstitial refuge availability may act as a limiting factor freshwater crayfish populations in
3 these systems. Accordingly, management actions that prioritise the protection and enhancement
1 of coarse littoral substrates and riparian vegetation, including the targeted placement of rocks
2 in areas already used by freshwater crayfish ([Johnsen & Taugbøl, 2008](#)), may help support the
3 persistence and local recovery of populations in lakes and represent a practical approach to
4 conserving functionally important habitat complexity in littoral ecosystems.

5 Acknowledgements

We thank Te Arawa Lakes Trust and Te Komiti Whakahaere for the opportunity to study kōura in the Rotorua Te Arawa lakes. We are grateful to Soweeta Fort-D’ath and William Anaru (Te Arawa Lakes Trust), and Tihini Grant (Ngāti Pikiao), for facilitating access to the lakes and supporting the field programme. We appreciate Joe Butterworth’s assistance with fieldwork, and the support of Andy Bruere and the Bay of Plenty Regional Council in helping to fund the fieldwork. We thank Calum MacNeil and three anonymous reviewers for feedback on an earlier version of this manuscript.

5.1 Funding statement

This research was supported by the Fish Futures programme funded through a Ministry of Business, Innovation and Employment grant (CAWX2101) with additional funding provided by the Bay of Plenty Regional Council under the Toi huarewa Waimāori - Bay of Plenty Regional Council Chair in Lake and Freshwater Science programme.

5.2 Authors’ contributions

Study conception and design were undertaken by O.V.R., F.J.B., I.A.K.K., R.H., and D.Ö. Fieldwork was conducted by O.V.R. Data analysis was performed by O.V.R., with statistical advice from F.J.B. Local ecological and traditional knowledge was contributed by I.A.K.K. and R.H. Manuscript drafting was led by O.V.R. All authors contributed to manuscript revision, approved the results, and the final manuscript.

5.3 Ethical approval

This study did not involve experimentation on humans or animals. Kōura and fish were captured using standard fisheries sampling methods. Permission to conduct field sampling and handle aquatic fauna was granted by Te Arawa Lakes Trust and Te Komiti Whakahaere. Formal institutional ethical approval was not required.

8 **5.4 Declarations**

9 The corresponding author has declared that none of the authors has any competing interests.

10 **5.5 Data availability statement**

11 The code and derived data supporting the findings of this study are openly available on GitHub
12 (https://github.com/OlivierRaven/Koura_shoreline_habitats) and a rendered version of the
13 analysis is hosted at https://olivierraven.github.io/Koura_shoreline_habitats/. The primary
14 raw dataset and code are archived on Zenodo (<https://doi.org/10.5281/zenodo.19476793>).
15 Peer review documents, including the cover letter and reviewer responses, are available in the
16 manuscript repository in the interest of open and transparent science.

17 **6 Tables**

Table 1: Physical, morphometric, and trophic characteristics of lakes in the Lake Taupo Regional Council database and number of littoral sites sampled per lake. Data include lake name, sampling date, elevation, mixing regime, and trophic state for 2025. Shoreline composition percentages are provided for cliff sections.

Lake name	Sampling date (n sites)	Surface area (km ²)	Perimeter length (km)	Catchment area (km ²)	Mean depth (m)
Rotorua	20/02/2025 (12)	81	45	508.0	11.0
Rotoiti	15/01/2025 (12)	34	61	123.7	31.5
Rotoehu	10/12/2024 (12)	8	40	49.2	8.0
Rotomā	6/11/2024 (12)	11	24	27.8	36.9
Ōkāreka	31/10/2024 (10), 22/01/2025 (2)	3	11	19.6	20.0

Table 2: Overview of environmental and biotic variables measured at littoral sampling sites, including variable descriptions, units, and hypothesised relevance for kōura (*Paranephrops planifrons*). Variables were selected a priori based on known habitat requirements, physiological constraints, and potential biotic interactions influencing kōura occurrence, abundance, and biomass in lake littoral zones.

Variable	Description / Unit	Hypothesised importance for kōura	References
Lake identity	Categorical variable identifying lake	Captures unmeasured lake-specific differences in water chemistry, productivity, and catchment characteristics.	Zuur et al., (2009)
Substrate index	Index based on % cover of bedrock, boulders, cobble, gravel, sand, mud, and organic matter	Important for burrowing and shelter availability. Coarser substrates increase shelter availability through more crevices.	Usio & Townsend, (2000); Kusabs et al., (2015b)
Slope	Slope from shoreline to the 5 m depth contour	Steeper slopes facilitate access to deeper water during daylight refugia and associate with coarser substrates.	Devcich, (1979); Kusabs et al., (2015b)
Riparian vegetation	Percentage cover of vegetation growing in the riparian zone	Contributes detrital inputs, bank stability, and shading at the water's edge.	Parkyn et al., (2002)
Overhanging trees	Percentage cover of trees hanging over the shoreline	Provides direct shading and structural inputs into littoral habitats.	Smith et al., (1996); Vedia et al., (2017)
Wood cover	Cover of wooden logs and tree branches in the sample site	Provides physical structure creating refuge spaces and supports macroinvertebrate prey availability.	Parkyn et al., (2009)

Table 2: Overview of environmental and biotic variables measured at littoral sampling sites, including variable descriptions, units, and hypothesised relevance for kōura (*Paranephrops planifrons*). Variables were selected a priori based on known habitat requirements, physiological constraints, and potential biotic interactions influencing kōura occurrence, abundance, and biomass in lake littoral zones.

Vari- able	Description / Unit	Hypothesised importance for kōura	References
Emer- gent and sub- merged macro- phytes	Percentage cover of macrophytes in the sample site divided over emergent and submerged and native and non-native species	Native macrophytes can provide cover and serve as a food source. Non-native macrophytes may alter movement pathways and modify local habitat and water quality conditions.	Coffey & Clayton, (1988); Kusabs & Quinn, (2009)
Tem- pera- ture	Temperature of surface water in °C	Influences metabolic rate, activity, physiological stress, and habitat suitability.	Devcich, (1979); Hammond et al., (2006); Parkyn & Collier, (2002); Angilletta et al., (2004)
Dis- solved oxygen	Dissolved oxygen concentration in mg L ⁻¹	Essential for respiration; reduced oxygen may constrain activity and habitat use.	Hammond et al., (2006); Broughton et al., (2017)
Spe- cific con- ductiv- ity	Electrical conductivity of the water in µS cm ⁻¹	Reflects overall lake productivity, supporting food availability.	Devcich, (1979)
pH	Acidity or alkalinity of the water	Influences moulting success and exoskeleton strength, affected by acidity or calcium levels.	Olsson et al., (2006)

Table 2: Overview of environmental and biotic variables measured at littoral sampling sites, including variable descriptions, units, and hypothesised relevance for kōura (*Paranephrops planifrons*). Variables were selected a priori based on known habitat requirements, physiological constraints, and potential biotic interactions influencing kōura occurrence, abundance, and biomass in lake littoral zones.

Vari- able	Description / Unit	Hypothesised importance for kōura	References
Fish pres- ence	Presence/absence of selected native and non-native fish species	Fish act as predators, competitors, or indirectly modify habitat structure.	Shave et al., (1994); Barnes, (1996); Usio & Townsend, (2000); Barnes & Hicks, (2003)

¹⁹ Source: [Article Notebook](#)

²⁰ **7 Figures**

²¹ **8 Supplementary information**

Table 3: Distribution of environmental and biotic variables measured at littoral sampling sites across five Te Arawa lakes in the Rotorua region of Aotearoa New Zealand.

Lake	Variable	Unit	n	Mean	Me- dian	Min	Max	CI low	CI high
Rotorua	Substrate index		12	3.66	3.30	2.80	6.30	3.01	4.31
Rotoiti	Substrate index		12	3.92	3.20	2.00	8.00	2.73	5.11
Ro- toehu	Substrate index		12	2.77	2.45	2.00	4.90	2.15	3.39
Rotomā	Substrate index		12	3.50	3.03	2.00	6.20	2.64	4.36
Ōkāreka	Substrate index		12	2.86	2.70	1.00	6.20	1.92	3.79
All lakes	Substrate index		60	3.34	3.00	1.00	8.00	2.98	3.71

Table 3: Distribution of environmental and biotic variables measured at littoral sampling sites across five Te Arawa lakes in the Rotorua region of Aotearoa New Zealand.

Lake	Variable	Unit	n	Mean	Me- dian	Min	Max	CI low	CI high
Rotorua	Slope 5m		12	0.02	0.01	0.01	0.16	0.00	0.05
Rotoiti	Slope 5m		12	0.09	0.07	0.04	0.30	0.04	0.14
Ro- toehu	Slope 5m		12	0.08	0.06	0.02	0.27	0.03	0.12
Rotomā	Slope 5m		12	0.07	0.06	0.01	0.23	0.03	0.11
Ōkāreka	Slope 5m		12	0.11	0.13	0.01	0.20	0.07	0.15
All lakes	Slope 5m		60	0.08	0.06	0.01	0.30	0.06	0.09
Rotorua	Riparian vegetation	%	12	67.50	100.00	0.00	100.00	40.12	94.88
Rotoiti	Riparian vegetation	%	12	92.50	100.00	20.00	100.00	77.88	107.12
Ro- toehu	Riparian vegetation	%	12	67.50	100.00	0.00	100.00	36.95	98.05
Rotomā	Riparian vegetation	%	12	63.33	90.00	0.00	100.00	35.13	91.53
Ōkāreka	Riparian vegetation	%	12	75.83	100.00	0.00	100.00	48.01	103.66
All lakes	Riparian vegetation	%	60	73.33	100.00	0.00	100.00	62.65	84.02
Rotorua	Overhanging trees	%	12	46.67	30.00	0.00	100.00	15.50	77.83
Rotoiti	Overhanging trees	%	12	54.17	65.00	0.00	100.00	23.22	85.11
Ro- toehu	Overhanging trees	%	12	35.83	0.00	0.00	100.00	5.25	66.42
Rotomā	Overhanging trees	%	12	52.50	55.00	0.00	100.00	22.44	82.56
Ōkāreka	Overhanging trees	%	12	52.50	65.00	0.00	100.00	21.71	83.29
All lakes	Overhanging trees	%	60	48.33	30.00	0.00	100.00	36.15	60.52
Rotorua	Wood cover	%	12	8.33	3.50	0.00	30.00	1.98	14.69
Rotoiti	Wood cover	%	12	4.25	1.50	0.00	20.00	0.31	8.19

Table 3: Distribution of environmental and biotic variables measured at littoral sampling sites across five Te Arawa lakes in the Rotorua region of Aotearoa New Zealand.

Lake	Variable	Unit	n	Mean	Me- dian	Min	Max	CI low	CI high
Ro- toehu	Wood cover	%	12	14.92	10.00	3.00	40.00	7.24	22.60
Rotomā	Wood cover	%	12	10.00	10.00	0.00	30.00	4.13	15.87
Ōkāreka	Wood cover	%	12	11.25	7.50	0.00	40.00	2.56	19.94
All lakes	Wood cover	%	60	9.75	5.00	0.00	40.00	6.96	12.54
Rotorua	Emergent macrophytes	%	12	0.00	0.00	0.00	0.00	0.00	0.00
Rotoiti	Emergent macrophytes	%	12	9.50	0.00	0.00	100.00	-8.71	27.71
Ro- toehu	Emergent macrophytes	%	12	8.75	0.00	0.00	35.00	-0.74	18.24
Rotomā	Emergent macrophytes	%	12	10.83	0.00	0.00	80.00	-4.78	26.45
Ōkāreka	Emergent macrophytes	%	12	22.50	0.00	0.00	90.00	-0.35	45.35
All lakes	Emergent macrophytes	%	60	10.32	0.00	0.00	100.00	3.98	16.65
Rotorua	Submerged macrophytes	%	12	0.83	0.00	0.00	10.00	-1.00	2.67
Rotoiti	Submerged macrophytes	%	12	0.83	0.00	0.00	10.00	-1.00	2.67
Ro- toehu	Submerged macrophytes	%	12	51.42	52.50	0.00	100.00	20.51	82.33
Rotomā	Submerged macrophytes	%	12	21.67	10.00	0.00	85.00	2.21	41.12
Ōkāreka	Submerged macrophytes	%	12	10.83	0.00	0.00	100.00	-7.43	29.10
All lakes	Submerged macrophytes	%	60	17.12	0.00	0.00	100.00	8.42	25.81

Table 3: Distribution of environmental and biotic variables measured at littoral sampling sites across five Te Arawa lakes in the Rotorua region of Aotearoa New Zealand.

Lake	Variable	Unit	n	Mean	Me- dian	Min	Max	CI low	CI high
Rotorua	Temperature	Degree C	12	23.15	23.25	21.80	25.30	22.52	23.78
Rotoiti	Temperature	Degree C	12	21.62	21.50	20.60	23.80	21.03	22.22
Ro- toehu	Temperature	Degree C	12	22.11	22.05	21.20	23.00	21.72	22.49
Rotomā	Temperature	Degree C	12	17.02	16.85	16.30	18.10	16.64	17.39
Ōkāreka	Temperature	Degree C	12	16.98	16.30	16.00	20.70	15.93	18.04
All lakes	Temperature	Degree C	60	20.18	21.25	16.00	25.30	19.44	20.91
Rotorua	Dissolved oxygen	mg/l	12	8.49	7.92	5.25	11.05	7.39	9.59
Rotoiti	Dissolved oxygen	mg/l	12	9.59	9.75	6.80	10.64	8.92	10.25
Ro- toehu	Dissolved oxygen	mg/l	12	10.36	10.77	8.85	11.79	9.67	11.05
Rotomā	Dissolved oxygen	mg/l	12	9.72	9.76	8.66	10.31	9.45	9.98
Ōkāreka	Dissolved oxygen	mg/l	12	9.37	9.38	8.83	9.74	9.20	9.53
All lakes	Dissolved oxygen	mg/l	60	9.50	9.62	5.25	11.79	9.20	9.81
Rotorua	Specific conductivity	µS/cm	12	205.36	188.27	156.20	443.01	156.39	254.33
Rotoiti	Specific conductivity	µS/cm	12	163.11	158.70	155.79	196.01	156.22	170.00
Ro- toehu	Specific conductivity	µS/cm	12	458.89	458.22	455.44	464.72	456.91	460.87
Rotomā	Specific conductivity	µS/cm	12	162.90	161.79	156.73	174.96	159.94	165.85
Ōkāreka	Specific conductivity	µS/cm	12	76.34	76.15	75.62	78.59	75.82	76.86

Table 3: Distribution of environmental and biotic variables measured at littoral sampling sites across five Te Arawa lakes in the Rotorua region of Aotearoa New Zealand.

Lake	Variable	Unit	n	Mean	Me- dian	Min	Max	CI low	CI high
All lakes	Specific conductivity	$\mu\text{S}/\text{cm}$	60	213.32	162.55	75.62	464.72	178.41	248.23
Rotorua	pH		12	6.94	7.04	3.78	7.69	6.29	7.60
Rotoiti	pH		12	8.67	8.73	8.08	9.02	8.46	8.87
Ro-toehu	pH		12	8.83	8.91	8.46	9.21	8.68	8.99
Rotomā	pH		12	7.80	7.92	6.44	8.56	7.47	8.13
Ōkāreka	pH		12	8.32	8.29	7.77	9.00	8.15	8.50
All lakes	pH		60	8.11	8.28	3.78	9.21	7.89	8.34
Rotorua	Presence Catfish		12	0.08	0.00	0.00	1.00	0.00	0.38
Rotoiti	Presence Catfish		12	0.42	0.00	0.00	1.00	0.15	0.72
Ro-toehu	Presence Catfish		12	0.00	0.00	0.00	0.00	0.00	0.26
Rotomā	Presence Catfish		12	0.00	0.00	0.00	0.00	0.00	0.26
Ōkāreka	Presence Catfish		12	0.00	0.00	0.00	0.00	0.00	0.26
All lakes	Presence Catfish		60	0.10	0.00	0.00	1.00	0.04	0.21
Rotorua	Presence Eel		12	0.33	0.00	0.00	1.00	0.10	0.65
Rotoiti	Presence Eel		12	0.08	0.00	0.00	1.00	0.00	0.38
Ro-toehu	Presence Eel		12	0.00	0.00	0.00	0.00	0.00	0.26
Rotomā	Presence Eel		12	0.25	0.00	0.00	1.00	0.05	0.57
Ōkāreka	Presence Eel		12	0.00	0.00	0.00	0.00	0.00	0.26
All lakes	Presence Eel		60	0.13	0.00	0.00	1.00	0.06	0.25
Rotorua	Presence Goldfish		12	0.08	0.00	0.00	1.00	0.00	0.38
Rotoiti	Presence Goldfish		12	0.67	1.00	0.00	1.00	0.35	0.90
Ro-toehu	Presence Goldfish		12	1.00	1.00	1.00	1.00	0.74	1.00

Table 3: Distribution of environmental and biotic variables measured at littoral sampling sites across five Te Arawa lakes in the Rotorua region of Aotearoa New Zealand.

Lake	Variable	Unit	n	Mean	Me- dian	Min	Max	CI low	CI high
Rotomā	Presence Goldfish		12	0.50	0.50	0.00	1.00	0.21	0.79
Ōkāreka	Presence Goldfish		12	0.25	0.00	0.00	1.00	0.05	0.57
All lakes	Presence Goldfish		60	0.50	0.50	0.00	1.00	0.37	0.63
Rotorua	Presence Common smelt		12	0.25	0.00	0.00	1.00	0.05	0.57
Rotoiti	Presence Common smelt		12	0.67	1.00	0.00	1.00	0.35	0.90
Ro-toehu	Presence Common smelt		12	0.50	0.50	0.00	1.00	0.21	0.79
Rotomā	Presence Common smelt		12	0.42	0.00	0.00	1.00	0.15	0.72
Ōkāreka	Presence Common smelt		12	0.58	1.00	0.00	1.00	0.28	0.85
All lakes	Presence Common smelt		60	0.48	0.00	0.00	1.00	0.35	0.62
Rotorua	Presence Kōaro		12	0.00	0.00	0.00	0.00	0.00	0.26
Rotoiti	Presence Kōaro		12	0.00	0.00	0.00	0.00	0.00	0.26
Ro-toehu	Presence Kōaro		12	0.00	0.00	0.00	0.00	0.00	0.26
Rotomā	Presence Kōaro		12	0.00	0.00	0.00	0.00	0.00	0.26
Ōkāreka	Presence Kōaro		12	0.67	1.00	0.00	1.00	0.35	0.90
All lakes	Presence Kōaro		60	0.13	0.00	0.00	1.00	0.06	0.25
Rotorua	Presence Trout		12	0.17	0.00	0.00	1.00	0.02	0.48
Rotoiti	Presence Trout		12	0.17	0.00	0.00	1.00	0.02	0.48
Ro-toehu	Presence Trout		12	0.00	0.00	0.00	0.00	0.00	0.26
Rotomā	Presence Trout		12	0.00	0.00	0.00	0.00	0.00	0.26
Ōkāreka	Presence Trout		12	0.17	0.00	0.00	1.00	0.02	0.48

Table 3: Distribution of environmental and biotic variables measured at littoral sampling sites across five Te Arawa lakes in the Rotorua region of Aotearoa New Zealand.

Lake	Variable	Unit	n	Mean	Me- dian	Min	Max	CI low	CI high
All lakes	Presence Trout		60	0.10	0.00	0.00	1.00	0.04	0.21

- 22 Albertson, L. K., & M. D. Daniels, 2018. Crayfish ecosystem engineering effects on riverbed
23 disturbance and topography are mediated by size and behavior. *Freshwater Science* 37: 836–844.
24 <https://doi.org/10.1086/700884>.
- 25 Angilletta, M. J., T. D. Steury, & M. W. Sears, 2004. Temperature, growth rate, and body
26 size in ectotherms: Fitting pieces of a life-history puzzle. *Integrative and Comparative Biology*
27 44: 498–509. <https://doi.org/10.1093/icb/44.6.498>.
- 1 Barnes, G. E., 1996. [The biology and general ecology of the brown bullhead catfish \(*Ameiurus*
2 *nebulosus*\) in Lake Taupo](#). Master of Science thesis, University of Waikato.
- 3 Barnes, G. E., & B. J. Hicks, 2003. Brown bullhead catfish (*Ameiurus nebulosus*) in Lake
4 Taupo. In *Managing invasive freshwater fish in New Zealand: Proceedings of a workshop hosted*
5 *by Department of Conservation, 10-12 may 2001, Hamilton. Department of Conservation.*
- 6 Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, & J.-S.
7 S. White, 2009. Generalized linear mixed models: A practical guide for ecology and evolution.
8 *Trends in Ecology & Evolution* 24: 127–135. <https://doi.org/10.1016/j.tree.2008.10.008>.
- 9 Brooks, M., B. Bolker, K. Kristensen, M. Maechler, A. Magnusson, H. Skaug, A. Nielsen, C.
10 Berg, & K. van Benthem, 2017. [glmmTMB: Generalized linear mixed models using template
11 model builder](#).
- 12 Broughton, R. J., I. D. Marsden, J. V. Hill, & C. N. Glover, 2017. Behavioural, physiological and
13 biochemical responses to aquatic hypoxia in the freshwater crayfish, *Paranephrops zealandicus*.
14 *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 212:
15 72–80. <https://doi.org/10.1016/j.cbpa.2017.07.013>.




Figure 1: Map of the sampling locations with different dominant littoral habitat types classified. a) Overview map of the 5 lakes. Lakes where invasive catfish were detected are indicated. b) Zoom in of Lake Ōkareka. c) Guidance map to show the lakes location (white square) on the North Island of Aotearoa New Zealand.




Figure 2: Kōura presence, CPUE, and BPUE across lakes. The upper panel shows the proportion of sampled sites with kōura present in each lake, with error bars indicating 95% binomial confidence intervals. The middle and lower panels show the distribution of kōura CPUE and BPUE, respectively, across lakes using boxplots (median, interquartile range, and range).




Figure 3: Kōura occupancy GAMM results and model performance. a) Estimated smooth terms for riparian vegetation, substrate index, surface water temperature, and specific conductivity, shown as partial effects with 95% confidence intervals. See Fig. S2 for raw data underlying modelled relationships. b) Receiver operating characteristic (ROC) curve illustrating model discrimination, with the area under the curve (AUC) reported. c) Calibration plot based on five equal-frequency bins, comparing mean predicted occupancy probabilities with observed proportions of occupied sites; the dashed 1:1 line indicates perfect calibration.




Figure 4: Kōura CPUE GAMM results and model performance. a) Estimated smooth terms for surface water temperature, pH, and emergent native macrophytes, shown with partial effects and 95% confidence intervals. See Fig. S2 for raw data underlying modelled relationships. b) Predicted versus observed CPUE from the positive (Gamma) component of the hurdle model, with predictions excluding the random lake effect. The dashed 1:1 line indicates perfect agreement; annotated values report R^2 and RMSE to summarise model fit.




Figure 5: Kōura BPUE GAMM results and model performance. a) Estimated smooth terms for pH and emergent native macrophytes, shown with partial effects and 95% confidence intervals. See Fig. S2 for raw data underlying modelled relationships. b) Predicted versus observed BPUE from the positive (Gamma) component of the hurdle model, with predictions excluding the random lake effect. The dashed 1:1 line indicates perfect agreement; annotated values report R^2 and RMSE to summarise model fit.




Figure 6: Length–weight relationship used to estimate missing kōura weights. Observed (measured) and model-predicted body weights are shown in relation to orbital carapace length (OCL). The log–log regression was fitted to individuals with both length and weight measurements ($n = 240$) and used to estimate missing weights ($n = 81$). Predicted values were back-transformed using a lognormal bias correction factor. The fitted regression equation is shown on the plot.




Figure 7: Raw relationships between all environmental predictors examined and kōura occurrence, CPUE, and BPUE across all 60 surveyed littoral sites. Each panel shows observed values at individual sites ($n = 60$ for occurrence; $n = 33$ for CPUE and BPUE). Points are jittered slightly for occurrence data to reduce overplotting. Predictors highlighted in bold were retained in final GAM models.

- 16 Bunch, A. J., M. S. Allen, & D. C. Gwinn, 2010. Spatial and temporal hypoxia dynamics in dense
17 emergent macrophytes in a Florida lake. *Wetlands* 30: 429–435. [https://doi.org/10.1007/s13157-](https://doi.org/10.1007/s13157-010-0051-9)
18 [010-0051-9](https://doi.org/10.1007/s13157-010-0051-9).
- 19 Capinha, C., E. R. Larson, E. Tricarico, J. D. Olden, & F. Gherardi, 2013. Effects of
20 climate change, invasive species, and disease on the distribution of native European crayfishes.
21 *Conservation Biology* 27: 731–740. <https://doi.org/10.1111/cobi.12043>.
- 22 Chambers, P. A., 1987. Nearshore occurrence of submersed aquatic macrophytes in relation
23 to wave action. *Canadian Journal of Fisheries and Aquatic Sciences* 44: 1666–1669. <https://doi.org/10.1139/f87-204>.
24
- 25 Coffey, B. T., & J. S. Clayton, 1988. Contrasting deep-water macrophyte communities in two
26 highly transparent New Zealand lakes and their possible association with freshwater crayfish,
1 *Paranephrops* spp. *New Zealand Journal of Marine and Freshwater Research* 22: 225–230.
2 <https://doi.org/10.1080/00288330.1988.9516294>.
- 3 Collier, K. J., S. M. Parkyn, & C. F. Rabeni, 1997. Koura: A keystone species. *Water and*
4 *Atmosphere* 5: 18–20.
- 5 Creed, R. P., & J. M. Reed, 2004. Ecosystem engineering by crayfish in a headwater stream
6 community. *Journal of the North American Benthological Society* 23: 224–236. [https://doi.](https://doi.org/10.1899/0887-3593(2004)023%3C0224:EEBCIA%3E2.0.CO;2)
7 [org/10.1899/0887-3593\(2004\)023%3C0224:EEBCIA%3E2.0.CO;2](https://doi.org/10.1899/0887-3593(2004)023%3C0224:EEBCIA%3E2.0.CO;2).
- 8 Danilović, M., I. Maguire, & L. Füreder, 2022. Overlooked keystone species in conservation
9 plans of fluvial ecosystems in Southeast Europe: A review of native freshwater crayfish species.
10 *Knowledge & Management of Aquatic Ecosystems* 423: 21. [https://doi.org/10.1051/kmae/](https://doi.org/10.1051/kmae/2022016)
11 [2022016](https://doi.org/10.1051/kmae/2022016).
- 12 Dedual, M., 2002. Vertical distribution and movements of brown bullhead (*Ameiurus nebulosus*
13 Lesueur 1819) in Motuoapa Bay, southern Lake Taupo, New Zealand. *Hydrobiologia* 483:
14 129–135. https://doi.org/10.1007/978-94-017-0771-8_15.
- 15 Devcich, A. A., 1979. An ecological study of *Paranephrops planifrons* White (Decapoda:
16 Parastacidae) in Lake Rotoiti, North Island. PhD thesis, University of Waikato.

- 17 Dumelle, M., T. Kincaid, A. R. Olsen, & M. Weber, 2023. Spsurvey: Spatial sampling design
18 and analysis in R. *Journal of Statistical Software* 105: <https://doi.org/10.18637/jss.v105.i03>.
- 19 Fielding, A. H., & J. F. Bell, 1997. A review of methods for the assessment of prediction
20 errors in conservation presence/absence models. *Environmental Conservation* 24: 38–49.
21 <https://doi.org/10.1017/S0376892997000088>.
- 22 Ford, T., & T. L. Beiting, 2005. Temperature tolerance in the goldfish, *Carassius auratus*.
23 *Journal of Thermal Biology* 30: 147–152. <https://doi.org/10.1016/j.jtherbio.2004.09.004>.
- 24 Francis, L. B., 2019. [Evaluating the effects of invasive brown bullhead catfish \(*Ameiurus*](#)
25 [nebulosus\)](#) on kōura (freshwater crayfish, *Paranephrops planifrons*) in Lake Rotoiti. Master’s
26 thesis, University of Waikato.
- 27 Froese, R., 2006. Cube law, condition factor and weight-length relationships: History, meta-
1 analysis and recommendations. *Journal of Applied Ichthyology* 22: 241–253. [https://doi.org/](https://doi.org/10.1111/j.1439-0426.2006.00805.x)
2 [10.1111/j.1439-0426.2006.00805.x](https://doi.org/10.1111/j.1439-0426.2006.00805.x).
- 3 Fry, F. E. J., & J. S. Hart, 1948. Cruising speed of goldfish in relation to water temperature.
4 *Journal of the Fisheries Research Board of Canada* 7b: 169–175. [https://doi.org/10.1139/f47-](https://doi.org/10.1139/f47-018)
5 [018](https://doi.org/10.1139/f47-018).
- 6 Grayling, S., 2020. [Bay of Plenty regional pest management plan 2011-2016 annual report for](#)
7 [2019/2020](#). Bay of Plenty Regional Council.
- 8 Greenaway, P., 1985. Calcium balance and moulting in the Crustacea. *Biological Reviews* 60:
9 425–454.
- 10 Hamilton, D., C. McBride, & T. Uraoka, 2005. Lake Rotoiti fieldwork and modelling to support
11 considerations of Ohau channel diversion from Lake Rotoiti. NIWA & University of Waikato.
- 12 Hammond, K. S., J. W. Hollows, C. R. Townsend, & P. M. Lokman, 2006. Effects of
13 temperature and water calcium concentration on growth, survival and moulting of freshwater
14 crayfish, *Paranephrops zealandicus*. *Aquaculture* 251: 271–279. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.aquaculture.2005.05.032)
15 [aquaculture.2005.05.032](https://doi.org/10.1016/j.aquaculture.2005.05.032).

- Harding, J. S., J. E. Clapcott, J. M. Quinn, J. W. Hayes, M. K. Joy, R. G. Storey, H. S. Greig, J. Hay, T. James, M. A. Beech, R. Ozane, A. S. Meredith, & I. K. G. Boothroyd, 2009. Stream habitat assessment protocols for wadeable rivers and streams of New Zealand. School of Biological Sciences, University of Canterbury.
- Haschenburger, J. K., & P. Roest, 2009. Substrate indices as indicators of interstitial pore space in gravel-bed channels. *River Research and Applications* 25: 98–105. <https://doi.org/10.1002/rra.1168>.
- Hesni, M. A., N. Shabanipour, A. Zahmatkesh, & M. M. Toutouni, 2009. Effects of temperature and salinity on survival and moulting of the narrow-clawed crayfish, *Astacus leptodactylus* Eschscholtz, 1823 (Decapoda, Astacidea). *Crustaceana* 82: 1495–1507. <https://doi.org/10.1163/156854009X407191>.
- Hessen, D. O., J. Skurdal, & J. E. Braathen, 2004. Plant exclusion of a herbivore; crayfish population decline caused by an invading waterweed. *Biological Invasions* 6: 133–140. <https://doi.org/10.1023/B:BINV.0000022131.40783.f0>.
- Hiroa, T. R., 1921. Māori food-supplies of Lake Rotorua, with methods of obtaining them, and usages and customs appertaining thereto. *Transactions and Proceedings of the Royal Society of New Zealand* 1868-1961 53: 433–451.
- Hopkins, C. L., 1970. Systematics of the New Zealand freshwater crayfish *Paranephrops* (Crustacea: Decapoda: Parastacidae). *New Zealand Journal of Marine and Freshwater Research* 4: 278–291. <https://doi.org/10.1080/00288330.1970.9515347>.
- Jane, S. F., G. J. A. Hansen, B. M. Kraemer, P. R. Leavitt, J. L. Mincer, R. L. North, R. M. Pilla, J. T. Stetler, C. E. Williamson, R. I. Woolway, & others, 2021. Widespread deoxygenation of temperate lakes. *Nature* 594: 66–70. <https://doi.org/10.1038/s41586-021-03550-y>.
- Johnsen, S. I., & T. Taugbøl, 2008. Add stones, get crayfish - is it that simple? *Freshwater Crayfish* 16: 47–50.
- Jowett, I. G., S. M. Parkyn, & J. Richardson, 2008. Habitat characteristics of crayfish (*Paranephrops planifrons*) in New Zealand streams using generalised additive models (GAMs).

- Hydrobiologia 596: 353–365. <https://doi.org/10.1007/s10750-007-9108-z>.
- Jowett, I. G., & J. Richardson, 1990. Microhabitat preferences of benthic invertebrates in a New Zealand river and the development of in-stream flow-habitat models for *Deleatidium* spp. New Zealand Journal of Marine and Freshwater Research 24: 19–30. <https://doi.org/10.1080/00288330.1990.9516399>.
- Kusabs, I. A. K., I. C. Duggan, A. C. Bruere, P. Scholes, T. W. Te Kurapa, T. M. Grant, & F. J. Burdon, 2026. Invasive catfish linked to the decline of freshwater crayfish in two Aotearoa New Zealand lakes. Biological Invasions 28: <https://doi.org/10.1007/s10530-026-03852-0>.
- Kusabs, I. A., B. J. Hicks, J. M. Quinn, & D. P. Hamilton, 2015a. Sustainable management of freshwater crayfish (kōura, *Paranephrops planifrons*) in Te Arawa (Rotorua) lakes, North Island, New Zealand. Fisheries Research 168: 35–46. <https://doi.org/10.1016/j.fishres.2015.03.015>.
- Kusabs, I. A., & J. M. Quinn, 2009. Use of a traditional Māori harvesting method, the tau kōura, for monitoring kōura (freshwater crayfish, *Paranephrops planifrons*) in Lake Rotoiti, North Island, New Zealand. New Zealand Journal of Marine and Freshwater Research 43: 713–722. <https://doi.org/10.1080/00288330909510036>.
- Kusabs, I. A., J. M. Quinn, & D. P. Hamilton, 2015b. Effects of benthic substrate, nutrient enrichment and predatory fish on freshwater crayfish (kōura, *Paranephrops planifrons*) population characteristics in seven Te Arawa (Rotorua) lakes, North Island, New Zealand. Marine and Freshwater Research 66: 631–643. <https://doi.org/10.1071/MF14148>.
- Kutner, M. H., C. J. Nachtsheim, J. Neter, & W. Li, 2005. Applied linear statistical models. McGraw-Hill Irwin.
- Land, Air, Water Aotearoa (LAWA), 2025. Lakes data for the Bay of Plenty region. <https://www.lawa.org.nz/explore-data/bay-of-plenty-region/lakes>.
- Lee, F., I. A. K. Kusabs, G. L. W. Perry, & C. MacNeil, 2025. Identifying refugia from the synergistic threats of climate change and invasive species. Web Ecology 25: 221–239. <https://doi.org/10.5194/we-25-221-2025>.

- 16 Levin, S. A., 1992. The problem of pattern and scale in ecology: The Robert H. MacArthur
17 award lecture. *Ecology* 73: 1943–1967. <https://doi.org/10.2307/1941447>.
- 18 Lishawa, S. C., A. J. Schrank, B. A. Lawrence, A. M. Monks, & D. A. Albert, 2023. Aquatic
19 connectivity treatments increase fish and macroinvertebrate use of *Typha*-invaded Great Lakes
20 coastal wetlands. *Freshwater Biology* 68: 1462–1477. <https://doi.org/10.1111/fwb.14141>.
- 21 Meerhoff, M., & M. de los Ángeles González-Sagrario, 2022. Habitat complexity in shallow lakes
22 and ponds: Importance, threats, and potential for restoration. *Hydrobiologia* 849: 3737–3760.
23 <https://doi.org/10.1007/s10750-021-04771-y>.
- 24 Momot, W. T., 1995. Redefining the role of crayfish in aquatic ecosystems. *Reviews in Fisheries*
25 *Science* 3: 33–63. <https://doi.org/10.1080/10641269509388566>.
- 26 Nyström, P., P. Stenroth, N. Holmqvist, O. Berglund, P. Larsson, & W. Granéli, 2006. Crayfish
27 in lakes and streams: Individual and population responses to predation, productivity and
1 substratum availability. *Freshwater Biology* 51: 2096–2113. [https://doi.org/10.1111/j.1365-
2 2427.2006.01641.x](https://doi.org/10.1111/j.1365-2427.2006.01641.x).
- 3 Olsson, K., P. Stenroth, P. Nyström, N. Holmqvist, A. R. McIntosh, & M. J. Winterbourn,
4 2006. Does natural acidity mediate interactions between introduced brown trout, native fish,
5 crayfish and other invertebrates in West Coast New Zealand streams? *Biological Conservation*
6 130: 255–267. <https://doi.org/10.1016/j.biocon.2005.12.019>.
- 7 Parkyn, S. M., & K. J. Collier, 2002. Differentiating the effects of diet and temperature on
8 juvenile crayfish (*Paranephrops planifrons*) growth: Leaf detritus versus invertebrate food
9 sources at two diurnally varying temperatures. *Freshwater Crayfish* 13: 371–382.
- 10 Parkyn, S. M., & K. J. Collier, 2004. Interaction of press and pulse disturbance on crayfish
11 populations: Flood impacts in pasture and forest streams. *Hydrobiologia* 527: 113–124.
12 <https://doi.org/10.1023/B:HYDR.0000043189.91134.94>.
- 13 Parkyn, S. M., K. J. Collier, & B. J. Hicks, 2001. New Zealand stream crayfish: Functional
14 omnivores but trophic predators? *Freshwater Biology* 46: 641–652. [https://doi.org/10.1046/j.
15 1365-2427.2001.00702.x](https://doi.org/10.1046/j.1365-2427.2001.00702.x).

- 16 Parkyn, S. M., M. A. Meleason, & R. J. Davies-Colley, 2009. Wood enhances crayfish
17 (*Paranephrops planifrons*) habitat in a forested stream. New Zealand Journal of Marine and
18 Freshwater Research 43: 689–700. <https://doi.org/10.1080/00288330909510034>.
- 19 Quinn, G. P., & M. J. Keough, 2002. Experimental design and data analysis for biologists.
20 Cambridge University Press.
- 21 Quinn, J. M., & C. W. Hickey, 1990. Magnitude of effects of substrate particle size, recent
22 flooding, and catchment development on benthic invertebrates in 88 New Zealand rivers. New
23 Zealand Journal of Marine and Freshwater Research 24: 411–427. [https://doi.org/10.1080/](https://doi.org/10.1080/00288330.1990.9516433)
24 [00288330.1990.9516433](https://doi.org/10.1080/00288330.1990.9516433).
- 25 R Core Team, 2025. [R: A language and environment for statistical computing](#). R Foundation
26 for Statistical Computing, Vienna, Austria.
- 1 Richman, N. I., M. Böhm, S. B. Adams, F. Alvarez, E. A. Bergey, J. J. S. Bunn, Q. Burnham,
2 J. Cordeiro, J. Coughran, K. A. Crandall, & others, 2015. Multiple drivers of decline in the
3 global status of freshwater crayfish (Decapoda: Astacidea). Philosophical Transactions of the
4 Royal Society B: Biological Sciences 370: 20140060. <https://doi.org/10.1098/rstb.2014.0060>.
- 5 Rosewarne, P., A. Piper, R. Wright, & A. Dunn, 2013. Do low-head riverine structures
6 hinder the spread of invasive crayfish? Case study of signal crayfish (*Pacifastacus leniusculus*)
7 movements at a flow gauging weir. Management of Biological Invasions 4: 273–282. <https://doi.org/10.3391/mbi.2013.4.4.02>.
- 9 Rowe, D. K., 1993. Identification of fish responsible for five layers of echoes recorded by
10 high-frequency (200 kHz) echosounding in Lake Rotoiti, North Island, New Zealand. New
11 Zealand Journal of Marine and Freshwater Research 27: 87–100. [https://doi.org/10.1080/](https://doi.org/10.1080/00288330.1993.9516548)
12 [00288330.1993.9516548](https://doi.org/10.1080/00288330.1993.9516548).
- 13 Schoen, E. R., D. A. Beauchamp, A. R. Buettner, & N. C. Overman, 2015. Temperature and
14 depth mediate resource competition and apparent competition between *Mysis diluviana* and
15 kokanee. Ecological Applications 25: 1962–1975. <https://doi.org/10.1890/14-1822.1>.
- 16 Schrank, A. J., & S. C. Lishawa, 2019. Invasive cattail reduces fish diversity and abundance in

- 17 the emergent marsh of a Great Lakes coastal wetland. *Journal of Great Lakes Research* 45:
18 1251–1259. <https://doi.org/10.1016/j.jglr.2019.09.013>.
- 19 Shave, C. R., C. R. Townsend, & T. A. Crowl, 1994. Anti-predator behaviours of a freshwater
20 crayfish (*Paranephrops zealandicus*) to a native and an introduced predator. *New Zealand*
21 *Journal of Ecology* 18: 1–10.
- 22 Shono, H., 2008. Application of the Tweedie distribution to zero-catch data in CPUE analysis.
23 *Fisheries Research* 93: 154–162. <https://doi.org/10.1016/j.fishres.2008.03.006>.
- 24 Smith, G. R. T., M. A. Learner, F. M. Slater, & J. Foster, 1996. Habitat features important
25 for the conservation of the native crayfish *Austropotamobius pallipes* in Britain. *Biological*
26 *Conservation* 75: 239–246. [https://doi.org/10.1016/0006-3207\(95\)00073-9](https://doi.org/10.1016/0006-3207(95)00073-9).
- 1 Statzner, B., O. Peltret, & S. Tomanova, 2003. Crayfish as geomorphic agents and ecosystem
2 engineers: Effect of a biomass gradient on baseflow and flood-induced transport of gravel and
3 sand in experimental streams. *Freshwater Biology* 48: 147–163. [https://doi.org/10.1046/j.1365-](https://doi.org/10.1046/j.1365-2427.2003.00984.x)
746 [2427.2003.00984.x](https://doi.org/10.1046/j.1365-2427.2003.00984.x).
- 747 Stevens, D. L., & A. R. Olsen, 2004. Spatially balanced sampling of natural resources.
748 *Journal of the American Statistical Association* 99: 262–278. [https://doi.org/10.1198/](https://doi.org/10.1198/016214504000000250)
749 [016214504000000250](https://doi.org/10.1198/016214504000000250).
- 750 Strayer, D. L., & S. E. G. Findlay, 2010. Ecology of freshwater shore zones. *Aquatic Sciences*
751 72: 127–163. <https://doi.org/10.1007/s00027-010-0128-9>.
- 752 Taylor, C. A., R. J. DiStefano, E. R. Larson, & J. Stoeckel, 2019. Towards a cohesive strategy for
753 the conservation of the United States’ diverse and highly endemic crayfish fauna. *Hydrobiologia*
754 846: 39–58. <https://doi.org/10.1007/s10750-019-04066-3>.
- 755 Usio, N., & C. R. Townsend, 2000. Distribution of the New Zealand crayfish (*Paranephrops*
756 *zealandicus*) in relation to stream physico-chemistry, predatory fish, and invertebrate prey. *New*
757 *Zealand Journal of Marine and Freshwater Research* 34: 557–567. [https://doi.org/10.1080/](https://doi.org/10.1080/00288330.2000.9516957)
758 [00288330.2000.9516957](https://doi.org/10.1080/00288330.2000.9516957).

- Usio, N., & C. R. Townsend, 2004. Roles of crayfish: Consequences of predation and bioturbation for stream invertebrates. *Ecology* 85: 807–822. <https://doi.org/10.1890/02-0618>.
- Vedia, I., D. Galicia, E. Baquero, J. Oscoz, & R. Miranda, 2017. Environmental factors influencing the distribution and abundance of the introduced signal crayfish in the north of Iberian Peninsula. *Marine and Freshwater Research* 68: 900–908. <https://doi.org/10.1071/MF16020>.
- Vincent, W. F., M. M. Gibbs, & S. J. Dryden, 1984. Accelerated eutrophication in a New Zealand lake: Lake Rotoiti, central North Island. *New Zealand Journal of Marine and Freshwater Research* 18: 431–440. <https://doi.org/10.1080/00288330.1984.9516064>.
- Westernhagen, N. von, D. P. Hamilton, & C. A. Pilditch, 2010. Temporal and spatial variations in phytoplankton productivity in surface waters of a warm-temperate, monomictic lake in New Zealand. *Hydrobiologia* 652: 57–70. <https://doi.org/10.1007/s10750-010-0318-4>.
- Wetzel, R. G., 2001. *Limnology: Lake and river ecosystems*. Gulf Professional Publishing.
- White, J., & K. Irvine, 2003. The use of littoral mesohabitats and their macroinvertebrate assemblages in the ecological assessment of lakes. *Aquatic Conservation: Marine and Freshwater Ecosystems* 13: 331–351. <https://doi.org/10.1002/aqc.586>.
- Wood, S. N., 2004. Stable and efficient multiple smoothing parameter estimation for generalized additive models. *Journal of the American Statistical Association* 99: 673–686. <https://doi.org/10.1198/016214504000000980>.
- Wood, S. N., 2017. [Generalized additive models: An introduction with R](#). Chapman; Hall/CRC.
- Woolway, R. I., S. Sharma, G. A. Weyhenmeyer, A. Debolskiy, M. Golub, D. Mercado-Bettín, M. Perroud, V. Stepanenko, Z. Tan, L. Grant, & others, 2021. Phenological shifts in lake stratification under climate change. *Nature Communications* 12: 2318. <https://doi.org/10.1038/s41467-021-22657-4>.
- Zhang, Y., J. Shen, L. He, J. Feng, L. Chi, & X. Wang, 2024. Challenge to lake ecosystems: Changes in thermal structure triggered by climate change. *Water* 16: 888. <https://doi.org/10.3390/w16080888>.

785 [3390/w16060888](#).

786 Zuur, A. F., E. N. Ieno, N. Walker, A. A. Saveliev, & G. M. Smith, 2009. [Mixed effects models](#)
787 [and extensions in ecology with R](#). Springer.