

INIT

Individually Neurostimulated Inhibition Training for AUD

Andrea Häfliger, Olivier Ulrich, Konstantinos Zervas

2021-12-03

Contents

1	Research Plan: Summary	3
2	Research Plan	4
2.1	State of research	4
2.2	Detailed research plan	4
2.2.1	Objectives and hypotheses	4
2.2.2	Participants	4
2.2.3	Study design, procedures, and measures	4
2.2.4	Data analysis	6
2.2.5	Ethical considerations	6
2.3	Timetable and milestones	6
2.4	Significance of planned research to the scientific community and to potential users	6

1 Research Plan: Summary

The proposed study will evaluate the potential of network-targeted brain stimulation to increase the efficacy of inhibition training protocols in treating alcohol use disorder. By comparing inhibition training accompanied by individually targeted transcranial magnetic stimulation with both sham, conventional TMS, a standalone inhibition training protocol, and a control group receiving the standard of care in AUD treatment, we aim to assess both the feasibility of such a regimen, as well as its effect on treatment outcomes relating to AUD. The inhibition training uses a modified Go/NoGo task to improve patients' ability to inhibit pre-potent responses toward alcoholic stimuli and has been shown to reduce alcohol intake (Houben et al., 2012). Performance differences between AUD patients and healthy controls in regular Go/NoGo tasks using alcohol related stimuli has been shown to depend on a network of neuronal circuitry commonly thought of being involved in inhibitory processes (Czapla et al., 2017; Luijten et al., 2014; Simmonds et al., 2008; Volkow and Baler, 2014). Network-targeted TMS has been suggested to be able to modulate neuronal activity in deeper lying brain regions, overcoming a critical hurdle of non-invasive brain stimulation (Momi et al., 2020). By using critical regions in inhibitory processes to build a functional connectivity map of inhibition we can identify individual cortical regions in each participant that can serve as gateways for TMS to target the underlying inhibitory network and thus increase inhibitory performance. This allows us to evaluate whether this facilitated inhibitory control can improve the effects of inhibition training for AUD.

2 Research Plan

2.1 State of research

2.2 Detailed research plan

2.2.1 Objectives and hypotheses

To further the understanding of the neuronal circuitry involved in inhibitory processes related to alcohol specific stimuli we intend to investigate functional connections between various subcortical regions known to be implicated in the act of inhibitory control, as well as search for further connections linking these areas to cortical regions that may be used as targets for non-invasive brain stimulation. We intend to use these findings to identify patient specific cortical gateway regions to stimulate during established inhibition training session in order to investigate the stimulation's potential to bolster the improvement in inhibitory control performance evoked by these trainings.

2.2.2 Participants

2.2.3 Study design, procedures, and measures

To allow for a thorough later analysis of the trainings' effects, we will initially conduct a baseline measure of both inhibitory control as well as the individuals' reactions to alcoholic stimuli during an fMRI scan. Inihbitory control is measured by a modified Go/NoGo task using alcohol related stimuli as no-go signals. These kinds of Go/NoGo task find widespread use in the assesment of inhibitory control in a context of AUD and provide a robust measure to assess an individual's ability to inhibit an automatised response to alcohol related information (Ames et al., 2014; Bowley et al., 2013; Rose

and Duka, 2008; Simmonds et al., 2008). Unlike the Go/NoGo task used in the actual inhibition training (Houben et al., 2012), this task uses the alcoholic stimuli, i.e. images depicting alcoholic beverages, as NoGo signals, which stand in contrast to pictures of non-alcoholic beverages that serve as Go signals. The participants are thus instructed to press a button whenever a Go signal appears, but not when a NoGo signal does. The stimuli are presented in a randomised order and a four to one ratio of Go and NoGo signals, respectively. This ratio has been shown to ensure the formation of a pre-potent response tendency towards Go signals (Ames et al., 2014). Since the task will be performed during an fMRI scan the timings of both the trials and the inter trial duration will be jittered.

Each participant’s neurophysiological reaction to alcoholic stimuli will be measured using a simple cue reactivity task consisting of alcoholic stimuli.

These baseline measures will be conducted during an fMRI scan, following a T1 weighted structural scan to infer anatomical information of each participant. Based on the fMRI data collected during both the cue reactivity task and the Go/NoGo task, BOLD signals will be analysed to assess individual neurophysiological activity. For the Go/NoGo task, BOLD activity during successfully inhibited responses (correctly rejected NoGo trials) and correctly answered Go trials are compared, allowing the identification of regions involved in inhibitory control. Contrasts in the cue reactivity task will be formed between the fixation trials and the alcohol stimulus trials, thus providing information on the location and extent of the individual physiological response.

The functional data obtained during the Go/NoGo task will be used to conduct connectivity analyses between inhibitory regions and a series of candidate regions closer to the surface of the brain. These analyses are intended to identify potential gateway regions, which can be used as targets of non-invasive brain stimulation and would propagate

the stimulation's effects to the lower laying inhibitory regions that cannot be targeted directly. The aim is to find the most positively correlated nodes of a functional connectivity map calculated from individual seed regions. The seeds will be chosen based on the BOLD activity during the inhibitory control task.

Potential gateway regions include the premotor cortex, the dorsolateral prefrontal cortex, and the orbitofrontal cortex. Based on these connectivity data for each participant, an individualised stimulation locust will be decided upon to be used during stimulation enhanced inhibition training.

2.2.4 Data analysis

2.2.5 Ethical considerations

2.3 Timetable and milestones

2.4 Significance of planned research to the scientific community and to potential users

References

- Ames, S. L., Wong, S. W., Bechara, A., Cappelli, C., Dust, M., Grenard, J. L., & Stacy, A. W. (2014). Neural correlates of a Go/NoGo task with alcohol stimuli in light and heavy young drinkers. *Behavioural Brain Research*, 274, 382–389. <https://doi.org/10.1016/j.bbr.2014.08.039>
- Bowley, C., Faricy, C., Hegarty, B., J. Johnstone, S., L. Smith, J., J. Kelly, P., & A. Rushby, J. (2013). The effects of inhibitory control training on alcohol consumption, implicit alcohol-related cognitions and brain electrical activity. *Inter-*

- national Journal of Psychophysiology*, 89(3), 342–348. <https://doi.org/10.1016/j.ijpsycho.2013.04.011>
- Czapla, M., Baeuchl, C., Simon, J. J., Richter, B., Kluge, M., Friederich, H.-C., Mann, K., Herpertz, S. C., & Loeber, S. (2017). Do alcohol-dependent patients show different neural activation during response inhibition than healthy controls in an alcohol-related fMRI go/no-go-task? *Psychopharmacology*, 234(6), 1001–1015. <https://doi.org/10.1007/s00213-017-4541-9>
- Houben, K., Havermans, R. C., Nederkoorn, C., & Jansen, A. (2012). Beer à no-go: Learning to stop responding to alcohol cues reduces alcohol intake via reduced affective associations rather than increased response inhibition: How stopping responding to alcohol reduces alcohol use. *Addiction*, 107(7), 1280–1287. <https://doi.org/10.1111/j.1360-0443.2012.03827.x>
- Luijten, M., Machielsen, M., Veltman, D., Hester, R., de Haan, L., & Franken, I. (2014). Systematic review of ERP and fMRI studies investigating inhibitory control and error processing in people. *Journal of Psychiatry & Neuroscience*, 39(3), 149–169. <https://doi.org/10.1503/jpn.130052>
- Momi, D., Neri, F., Coiro, G., Smeralda, C., Veniero, D., Sprugnoli, G., Rossi, A., Pascual-Leone, A., Rossi, S., & Santarnecchi, E. (2020). Cognitive Enhancement via Network-Targeted Cortico-cortical Associative Brain Stimulation. *Cerebral Cortex*, 30(3), 1516–1527. <https://doi.org/10.1093/cercor/bhz182>
- Rose, A. K., & Duka, T. (2008). Effects of alcohol on inhibitory processes. *Behavioural Pharmacology*, 19(4), 284–291. <https://doi.org/10.1097/FBP.0b013e328308f1b2>
- Simmonds, D. J., Pekar, J. J., & Mostofsky, S. H. (2008). Meta-analysis of Go/No-go tasks demonstrating that fMRI activation associated with response inhibition is

task-dependent. *Neuropsychologia*, 46(1), 224–232. <https://doi.org/10.1016/j.neuropsychologia.2007.07.015>

Volkow, N., & Baler, R. (2014). Addiction science: Uncovering neurobiological complexity. *Neuropharmacology*, 76, 235–249. <https://doi.org/10.1016/j.neuropharm.2013.05.007>