# Feedback — Assignment 3

Help

You submitted this quiz on **Sun 18 May 2014 6:24 PM PDT**. You got a score of **7.00** out of **7.00**.

#### **Question 1**

At noted in the lectures, in single-variable systems, rate-balance plots can be used to determine whether bistability is present.

Several examples are given in the lecture notes. Download the Matlab script ratebalance.m from here. This script plots these curves for two simple cases:


- 1) A constant forward rate
- 2) An ultrasensitive autocatalytic feedback that causes the forward rate to be highly nonlinear.


The variable of interest A\* (Astar in the code) is assumed to be normalized to the quantity [A]<sub>TOTAL</sub>, so this ranges from 0 to 1. For several values of the stimulus S, the program plots forward and backward rates, as well as the steady-states where the forward and backward rates cross.

Based on what you have learnt about autocatalytic feedback loops, which of the following plots represents a case where bistability presents? (Red line shows the backward rate and blue line shows the forward rate)


Your Answer Score Explanation








1.8 -1.6 -





## **Question 2**

Bistability can be produced with ultrasensitive feedback as in the example. It is not possible with linear feedback and a first order reverse rate. However, bistability can be produced if linear autocatalytic feedback is combined with a "back reaction" that becomes partially saturated when the level of product (A\* in this case) increases.

Consider the linear feedback plus saturating back reaction:

$$FR = (k_{+}[S] + k_{f}[A^{*}])([A]_{TOTAL} - [A^{*}])$$

$$ER = k_{minus} \left(\frac{[A^{*}]}{[A^{*}] + K_{mb}}\right)$$

$$ER = k_{minus} \left(\frac{[A^{*}]}{[A^{*}] + K_{mb}}\right)$$

Modify ratebalance.m to simulate this condition. You can assume that the stimulus S is zero. Use the following constants: kminus = 5;  $K_{mb}$  = 0.1, kplus=2, kf=30.

Test several different values of  $K_{mb}$  and determine which values of  $K_{mb}$  produce bistability. You will have to determine the "interesting" range of  $K_{mb}$  values to examine. Plot your results as a bifurcation diagram:  $K_{mb}$  on the abscissa, steady-state values of  $[A^*]$  on the ordinate. Lines 61-69 in the program are written to be able to accommodate the situation in which one or multiple steady-states are possible.

Which of the following statements is correct about  $K_{mb}$  and bistability in this system? (Assume: kminus = 5, kplus=2, kf=30, S=1)

Your Answer		Score	Explanation
The only condition in which bistability present is when $\mathbf{K}_{\mathbf{mb}}$ >1.			
There is no bistability in the system when 0.1> $\mathbf{K}_{mb}$ .			
●When K <sub>mb</sub> >3 , there is no bistability in the system.	<b>~</b>	1.00	
When K <sub>mb</sub> >0.1 bistability is present.			
Total		1.00 /	
		1.00	

#### **Question 3**

In the above case assume that  $\mathbf{K}_{mb}$  is fixed and equals to 1, which of the following statements is correct? (kminus = 5, kplus=2, kf=30) (Remeber that S can not be a negative number)

Your Answer		Score	Explanation
The system shows bistability when S>3.			
The system is always mono-stable regardless of value of S.	<b>~</b>	1.00	
The system shows bistability when $0.1 < S < 1$ .			
The system shows bistability when S>1.			
Total		1.00 / 1.00	

## **Question 4**

What if a system has both ultrasensitive autocatalytic feedback and a back reaction that becomes saturated? Is bistability in such a system likely to be less robust or more robust compared with a system that contains only one of these features? You do not have to explicitly simulate this condition, but you should be able to make an argument based on the shapes of the forward and backwards reaction rates. If you do choose to simulate this, it is probably easiest to modify the code so that it simulates several values of stimulus [S] but only single values of other parameters.

Using the following parameter sets which statement is correct when comparing a system with ultrasensitive autocatalytic feedback loop, a system with linear feedback plus saturating back reaction and a system which has both of this conditions? (kminus = 5;  $K_{mb}$  = 0.5, kplus=0.5, kf=30, Kmf = 0.5, h = 4)

Your Answer		Score	Explanation
•Ultrasensitive autocatalytic feedback loop with a back reaction that becomes saturated shows bistability in a longer range of values of parameter S.	<b>~</b>	1.00	
Ultrasensitive autocatalytic feedback loop shows bistability in a longer range of values of parameter S.			
Linear feedback plus saturating back reaction shows bistability in a longer range of values of parameter S.			

Oultrasensitive autocatalytic feedback loop, linear feedback loop plus saturating back reaction, and a system with both of them show bistability in the same range of values of parameter S.

Total 1.00 / 1.00

#### **Question 5**

We now implement a simple two-variable model of the E. coli lac operon. The differential equations governing this system are as follows:

$$\frac{dl}{dt} = \beta l_{ext} LacY - \gamma l$$

$$\frac{dLacY}{dt} = \delta + p \frac{l^4}{l^4 + l_0^4} - \sigma LacY$$

$$\beta = 1, \gamma = 1, \delta = 0.2, l_0 = 4, p = 4, \sigma = 1$$

The variable  $I_{\text{ext}}$ , representing extracellular lactose, is the variable that can vary and potentially cause the bacteria to change states. Plot the nullclines of this model for  $I_{\text{ext}}$  =2.5. Because of the  $I_{\text{ext}}$  term appearing in the second equation, it is easiest to treat LacY as the dependent variable (ordinate) and I as the independent variable (abscissa). Which of the following statements is correct?

Your Answer		Score	Explanation
Nullclines intersect in three points which one of them is an unstable steady states.	<b>~</b>	1.00	
Above the LacY nullcline, LacY and / are increasing.			
Nullclines intersect in three points which all of them are stable steady states.			

Nullclines intersect in two points which both of them are unstable steady states.

Total

1.00 /
1.00

## **Question 6**

Modify the program repression.m (You can download it from here) so that this model now simulates the temporal evolution of I and LacY. To reach steady-state, a total simulation time of 20 seconds is reasonable, and the solution converges with a time step of 0.01 if Euler's method is used. For  $I_{ext} = 2.5$ , plot the time evolution of LacY and I for the following initial conditions:

- 1) / =8, LacY=3;
- 2) /=3, LacY=1.3;
- 3) /=3, LacY=1.2;
- 4) /=2, LacY=1;

Which of the following statements is correct?

Your Answer		Score	Explanation
•Initial values of LacY and / determine if they will reach their lower or higher stable steady state levels.	<b>~</b>	1.00	
has always lower stable steady state levels compared to LacY.			
CLacY has two stable steady states and I has one stable steady state.			
Changing the initial condition of LacY and / can remove the bistability from the system.			
Total		1.00 /	
		1.00	

## **Question 7**

Vary  $I_{\text{ext}}$  over the range 1 to 7 and compute the temporal evolution of I and LacY. For each simulation, store the final value of LacY. Run two sets of simulations: one in which the initial conditions are I = 8, LacY=3, and a second in which the initial conditions are I = 2, LacY=1. Generate a bifurcation diagram of how the final value of LacY varies as a function of  $I_{\text{ext}}$ . Plot the results of the two sets of simulations in different colors.

Which of the following statements is correct?

Your Answer		Score	Explanation
When extracellular lactose is less than 1, the system is bistable.			
●When 2 < I <sub>ext</sub> < 4 , the system is bistable.	~	1.00	
Changing the initial conditions of / and LacY can change the bifurcation points.			
For lext > 6 the system is bistable.			
Total		1.00 /	
		1.00	