

CRITICAL PAPER REVIEW

INTRINSIC REACTIVITY OF Ni, Pd AND Pt SURFACES IN DRY REFORMING AND COMPETITIVE REACTIONS:
INSIGHTS FROM FIRST PRINCIPLES CALCULATIONS AND MICROKINETIC MODELING SIMULATIONS




Authored by: Lucas Foppa, Marius-Christian Silaghi, Kim Larmier, Aleix Comas-Vives




Review by: Om Mihani
Course: CL 325

ABSTRACT (THE WHAT?)

1. Activity and intrinsic mechanisms of DRM on various catalysts
2. Analysis done using Density functional theory and Microkinetic calculations
3. Reactions:



THE MECHANISM (THE WHAT?)



NEED (THE WHY?)

- Both Economical and ecological need
- Ecological: Substitute for SRM (Water usage)
- Economical: Controllable ratio of CO and H₂O
- Ecological: Consumes CO₂
- Economical: Can be used to extract energy from gas hydrates

METHOD (THE HOW?)

- Density Functional Theory
 - quantum mechanical modelling method
 - used to calculate the electronic structure of atoms, molecules, and materials
 - works by solving the quantum mechanical equations
 - Simpler than solving the Schrödinger equations
- Microkinetic Modelling
 - Calculations to guess the rates of different steps based on DFT calculations

THE GOOD PART

- An excellent step by step analysis of various possible elementary steps in the mechanism
- For each step, there is a comparison based on thermodynamic calculations between three catalysts: Ni, Pd, Pt (all from a family)
- DFT calculations were compared with experimental data and were found to match well within acceptable limits
- All possible side reactions along with Coking were also analysed
- Effects of pressure also analysed
- In section 3.5, H^* stability was analysed thermodynamically, and a physical explanation was also given. This can be observed throughout the paper

THE DRAWBACKS

- No usable conclusions
- DFT is a strong method and not very computationally expensive but is strongly dependent on the functionals used. This can allow errors
- Defects in catalyst and adsorbate-adsorbate interactions were ignored which might not always be insignificant
- Stability of an intermediate is directly correlated to its adsorption enthalpy which may not be a correct interpretation
- Thermal effects and other sides not analysed
- Continued on next slide

MICROKINETIC MODELLING

- Model assumes catalyst surface remains constant throughout the reaction which is incorrect, owing to Coking
- It is mentioned that the accurate model is computationally expensive hence approximate models were used
- It is great as it:
 - Provides insight into mechanisms
 - Aids in catalyst design
 - Accounts for Temp and Pressure
- But has disadvantages:
 - Sensitive to assumptions
 - Computationally expensive
 - Can't capture effect of heat and mass transfer (accounted for by adsorption studies)

WRITING STYLE AND OTHER COMMENTS

- Clear and concise
- Final summary is missing
- All interpretations backed by data
- Still has scope of other studies
- An academic paper, not industrially usable

Citations

- Lucas Foppa, Marius-Christian Silaghi, Kim Larmier, Aleix Comas-Vives,
● <https://doi.org/10.1016/j.jcat.2016.02.030>.
- 2. Hohenberg, P. and Kohn, W. (1964) Inhomogeneous Electron Gas. Physical Review B, 136, 864-871.
● <http://dx.doi.org/10.1103/PhysRev.136.B864>
- 3. Kohn, W. and Sham, L.J. (1965) Self-Consistent Equations Including Exchange and Correlation Effects.
● Physical Review, 140, A1133-A1138. <http://dx.doi.org/10.1103/PhysRev.140.A1133>
- 4. Vlachos, D. G. Kinetics of Heterogeneous Catalytic Reactions; Princeton University Press, 2017.
- 5. Campbell, C. T. Microkinetics: From Heterogeneous Catalysis to Solid-State Chemistry; Wiley-VCH
● Verlag GmbH & Co. KGaA, 2014

THANK YOU