

Assignment-based Subjective Questions &

Answers:

-Om Mihir R Bhoopalam

1) From your analysis of the categorical variables from the dataset, what could you infer about their effect on the dependent variable?

Ans. Following are the effects for categorical variables on the dependent variable:


- a. Fall has the highest median, which is expected as weather conditions are most optimal to ride bike followed by summer.
- b. Median bike rents are increasing year on as year 2019 has a higher median than 2018, it might be due the fact that bike rentals are getting popular and people are becoming more aware about environment.
- c. Overall spread in the month plot is reflection of season plot as fall months have higher median.
- d. People rent more on non - holidays compared to holidays, so reason might be they prefer to spend time with family and use personal vehicle instead of bike rentals.
- e. Overall median across all days is same but spread for Saturday and Wednesday is bigger may be evident that those who have plans for Saturday might not rent bikes as it a non-working day.
- f. Working and non-working days have almost the same median although spread is bigger for non-working days as people might have plans and do not want to rent bikes because of that
- g. Clear weather is most optimal for bike renting, as temperate is optimal, humidity is less, and temperature is less.

2) Why is it important to use drop_first=True during dummy variable creation?


Ans. The drop_first = True is important as it helps in reducing the extra column created during creation of dummy variable. Dropping the column is important because the importance or value of that left over variable can be found by remaining variables. So to avoid redundancy we are dropping a column. This helps in the column to become linearly independent.

3) Looking at the pair-plot among the numerical variables, which one has the highest correlation with the target variable?

Ans. Temp and atemp are the numerical variables which are showing the highest correlation with the target variable.



4) How did you validate the assumptions of Linear Regression after building the model on the training set?



Ans. Residuals distribution should follow normal distribution and centered around 0.(mean = 0). We validate this assumption about residuals by plotting a distplot of residuals and see if residuals are following normal distribution or not. The above diagram shows that the residuals are distributed about mean = 0.

5) Based on the final model, which are the top 3 features contributing significantly towards explaining the demand of the shared bikes?

Ans. Based on the final model , the top 3 features contributing significantly towards explaining the demand of the shared bikes are:

- yr
- temp
- weather

General Subjective Questions & Answers:

1) Explain the linear regression algorithm in detail.

Ans. Linear Regression Algorithm is a machine learning algorithm based on supervised learning where the predicted output is continuous and has a constant slope. It's used to predict values within a continuous range, rather than trying to classify them into categories. It is a part of regression analysis. Linear regression shows the linear relationship between the independent variable (X-axis) and the dependent variable (Y-axis), consequently called linear regression. If there is a single input variable (x), such linear regression is called simple linear regression. And if there is more than one input variable, such linear regression is called multiple linear regression.

Simple Linear Regression - Simple linear regression uses traditional slope-intercept form, where m and b are the variables our algorithm will try to "learn" to produce the most accurate predictions. x represents our input data and y represents our prediction.

$$Y=mx+b$$

where,

y = dependent variable

x = independent variable

m = intercept of the line

b = linear regression coefficient

When the value of x (independent variable) increases, the value of y (dependent variable) is likewise increasing. Regression method tries to find the best fit line which shows the relationship between the dependent variable and predictors with least error.

Assumptions of Simple Linear Regression –

There are four assumptions associated with a linear regression model:

- a. **Linearity:** The relationship between X and the mean of Y is linear.
- b. **Homoscedasticity:** The variance of residual is the same for any value of X.
- c. **Independence:** Observations are independent of each other.
- d. **Normality:** For any fixed value of X, Y is normally distributed.

Multiple Linear Regression - A more complex, multi-variable linear equation might look like this, where w represents the coefficients, or weights, our model will try to learn.

$$f(x,y,z)=w_1x + w_2y + w_3z$$

The variables x,y,z represent the attributes, or distinct pieces of information, we have about each observation.

Assumptions of Multiple Linear Regression –

Multiple linear regression analysis makes five key assumptions:

- a. **Linear relationship:** There exists a linear relationship between each predictor variable and the response variable.
- b. **No Multicollinearity:** None of the predictor variables are highly correlated with each other.
- c. **Independence:** The observations are independent.
- d. **Homoscedasticity:** The residuals have constant variance at every point in the linear model.
- e. **Multivariate Normality:** The residuals of the model are normally distributed.


2) Explain the Anscombe's quartet in detail.

Ans. Anscombe's quartet was constructed in 1973 by statistician Francis Anscombe to illustrate the importance of plotting data before you analyze it and build your model. These four data sets have nearly the same statistical observations, which provide the same information (involving variance and mean) for each x and y point in all four data sets. However, when you plot these data sets, they look very different from one another. Anscombe's quartet intended to counter the impression among statisticians that "numerical calculations are exact, but graphs are rough."

These four plots can be defined as follows:

Anscombe's quartet							
I		II		III		IV	
x	y	x	y	x	y	x	y
10.0	8.04	10.0	9.14	10.0	7.46	8.0	6.58
8.0	6.95	8.0	8.14	8.0	6.77	8.0	5.76
13.0	7.58	13.0	8.74	13.0	12.74	8.0	7.71
9.0	8.81	9.0	8.77	9.0	7.11	8.0	8.84
11.0	8.33	11.0	9.26	11.0	7.81	8.0	8.47
14.0	9.96	14.0	8.10	14.0	8.84	8.0	7.04
6.0	7.24	6.0	6.13	6.0	6.08	8.0	5.25
4.0	4.26	4.0	3.10	4.0	5.39	19.0	12.50
12.0	10.84	12.0	9.13	12.0	8.15	8.0	5.56
7.0	4.82	7.0	7.26	7.0	6.42	8.0	7.91
5.0	5.68	5.0	4.74	5.0	5.73	8.0	6.89

When these models are plotted on a scatter plot, all datasets generates a different kind of plot that is not interpretable by any regression algorithm which is fooled by these peculiarities and can be seen as follows:



The four datasets can be described as:

- a. The first scatter plot (top left) appears to be a simple linear relationship,
- b. The second graph (top right); cannot fit the linear regression model because the data is non-linear
- c. In the third graph (bottom left), the modelled relationship is linear, but should have a different regression line. It shows the outliers involved in the dataset which cannot be handled by linear regression model
- d. Finally, the fourth graph (bottom right) shows the outliers involved in the dataset which cannot be handled by linear regression model. It shows an example when one high-leverage point is enough to produce a high correlation coefficient, even though the other data points do not indicate any relationship between the variables. It shows the outliers involved in the dataset which cannot be handled by linear regression model.

3. What is Pearson's R?

Ans. The Pearson correlation method is the most common method used for numerical variables. It assigns a value between – 1 and 1, where 0 is no correlation, 1 is total positive correlation, and – 1 is total negative correlation. Pearson correlation coefficient (PCC), also referred to as Pearson's r, the Pearson product moment correlation coefficient (PPMCC), or the bivariate correlation, is a measure of linear correlation between two sets of data. It is the covariance of two variables, divided by the product of their standard deviations.

Pearson's R Formula is as follows:

$$r = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}}$$

Here,

1. r = correlation coefficient
2. xi = values of the x-variable in a sample
3. \bar{x} = mean of the values of the x-variable
4. yi = values of the y-variable in a sample
5. \bar{y} = mean of the values of the y-variable

4. What is scaling? Why is scaling performed? What is the difference between normalized scaling and standardized scaling?

Ans. Scaling is a method used to normalize the range of independent variables or features of data. In data processing, it is also known as data normalization and is generally performed during the data preprocessing step. In simpler terms, in machine learning algorithms we need to bring all features in the same standing, so that one significant number doesn't impact the model just because of their large magnitude. This is called scaling or Feature scaling.

Feature scaling in machine learning is one of the most critical steps during the pre-processing of data before creating a machine learning model. Scaling can make a difference between a weak machine learning model and a better one. Most of the times, collected data set contains features highly varying in magnitudes, units and range. If scaling is not done then algorithm only takes magnitude in account and not units which results in an incorrect model. To solve this issue, we have to do scaling to bring all the variables to the same level of magnitude.

It is important to note that scaling just affects the coefficients and none of the other parameters like t-statistic, F-statistic, p-values, R-squared, etc.

Sl. No	Normalization	Standardization
1	Feature scaling method to bring the data into common range such as [0, 1], [-1, 1], etc.	Feature scaling method bring the data with mean 0 and unit variance
2	Scikit-learn provides MinMaxScaler, MaxAbsScaler and RobustScaler methods for normalization	Scikit-learn provides StandardScaler for standardization
3	MinMaxScaler and MaxAbsScaler are sensitive to outliers whereas RobustScaler is more robust to outliers	Standardization is less sensitive to outliers compared to MinMaxScaler and MaxAbsScaler
4	Useful when we don't know about the distribution of features and there are no or little outliers - MinMaxScaler: if features don't follow normal distribution and if there are no or less outliers - MaxAbsScaler: if the data is sparse - RobustScaler: if the data contains outliers	Useful when we know features are normally distributed (Gaussian distribution)

5. You might have observed that sometimes the value of VIF is infinite. Why does this happen?

Ans. Variance Inflation Factor (VIF) is an index that provides a measure of how much the variance of an estimated regression coefficient increases due to collinearity. It is calculated by taking the ratio of the variance of all a given model's betas divide by the variance of a single beta if it were fit alone. The higher the VIF value, the greater the correlation of the variable with other variables. Values of more than 4 or 5 are sometimes regarded as being moderate to high, with values of 10 or more being regarded as very high. If there is perfect correlation, then **VIF = infinity**. An infinite VIF value means that the variable is exactly linear combination of other variable. If the independent variable can be explained perfectly by other independent variables, then it will have perfect correlation and its R-squared value will be equal to 1. So, $VIF = 1/(1-1)$ which gives $VIF = 1/0$ which results in "infinity".

6. What is a Q-Q plot? Explain the use and importance of a Q-Q plot in linear regression.

Ans. Quantile - Quantile (Q-Q) plot, is a graphical tool to help us assess if a set of data plausibly came from some theoretical distribution such as a Normal, Exponential or Uniform distribution. Also, it helps to determine if two data sets come from populations with a common distribution. A Q-Q plot is used to compare the shapes of distributions, providing a graphical view of how properties such as location, scale, and skewness are similar or different in the two distributions. The power of Q-Q plots lies in their ability to summarize any distribution visually.

The advantages of the Q-Q plot are:

- a. The sample sizes do not need to be equal.
- b. Many distributional aspects can be simultaneously tested.

Q-Q plot is very useful to determine:

- a. If two populations are of the same distribution
- b. If residuals follow a normal distribution. Having a normal error term is an assumption in regression and we can verify if it's met using this.
- c. Skewness of distribution.