

# Filtering in Frequency Domain

Lecture 1 (Introduction and Maths Primer)


Prepared by: Neha Fotedar

# Why we switch to Frequency Domain?





# Fourier Transform





# Applications



- Reading Text – Captcha
- Transferring books to electronic copies
- Number plate recognition
- Automating the input of handwritten forms




Time domain




Frequency domain


























Image: 'cameraman'


*Copyright owned by MIT. Used with permission.*

<https://au.mathworks.com/help/images/examples/deblurring-images-using-the-blind-deconvolution-algorithm.html>


A




A




A




B



C



D








Image: 'cameraman'

*Copyright owned by MIT. Used with permission.*

<https://au.mathworks.com/help/images/examples/deblurring-images-using-the-blind-deconvolution-algorithm.html>




Image: 'cameraman'

*Copyright owned by MIT. Used with permission.*

<https://au.mathworks.com/help/images/examples/deblurring-images-using-the-blind-deconvolution-algorithm.html>



a b  
c d

**FIGURE 4.64**

- (a) Sampled newspaper image showing a moiré pattern.
- (b) Spectrum.
- (c) Butterworth notch reject filter multiplied by the Fourier transform.
- (d) Filtered image.

a b  
c d


**FIGURE 4.64**

(a) Sampled newspaper image showing a moiré pattern.

(b) Spectrum.

(c) Butterworth notch reject filter multiplied by the Fourier transform.

(d) Filtered image.



a b  
c d


**FIGURE 4.64**

(a) Sampled newspaper image showing a moiré pattern.

(b) Spectrum.

(c) Butterworth notch reject filter multiplied by the Fourier transform.

(d) Filtered image.



a b  
c d


**FIGURE 4.64**

(a) Sampled newspaper image showing a moiré pattern.

(b) Spectrum.

(c) Butterworth notch reject filter multiplied by the Fourier transform.

(d) Filtered image.





a b  
c d


**FIGURE 4.64**

(a) Sampled newspaper image showing a moiré pattern.

(b) Spectrum.

(c) Butterworth notch reject filter multiplied by the Fourier transform.

(d) Filtered image.





**FIGURE 4.65**

(a)  $674 \times 674$

image of the  
Saturn rings  
showing nearly  
periodic  
interference.

(b) Spectrum: The  
bursts of energy  
in the vertical axis  
near the origin  
correspond to the  
interference

pattern. (c) A  
vertical notch  
reject filter.

(d) Result of  
filtering. The thin  
black border in  
(c) was added for  
clarity; it is not  
part of the data.

(Original image  
courtesy  
of Dr. Robert  
A. West,  
NASA/JPL.)


a	b
c	d

**FIGURE 4.65**

(a)  $674 \times 674$  image of the Saturn rings showing nearly periodic interference.

(b) Spectrum: The bursts of energy in the vertical axis near the origin correspond to the interference pattern. (c) A vertical notch reject filter.

(d) Result of filtering. The thin black border in (c) was added for clarity; it is not part of the data. (Original image courtesy of Dr. Robert A. West, NASA/JPL.)






a	b
c	d

**FIGURE 4.65**

(a)  $674 \times 674$  image of the Saturn rings showing nearly periodic interference.

(b) Spectrum: The bursts of energy in the vertical axis near the origin correspond to the interference pattern. (c) A vertical notch reject filter.


(d) Result of filtering. The thin black border in (c) was added for clarity; it is not part of the data. (Original image courtesy of Dr. Robert A. West, NASA/JPL.)



a	b
c	d

**FIGURE 4.65**


(a)  $674 \times 674$  image of the Saturn rings showing nearly periodic interference.  
 (b) Spectrum: The bursts of energy in the vertical axis near the origin correspond to the interference pattern.  
 (c) A vertical notch reject filter.  
 (d) Result of filtering. The thin black border in (c) was added for clarity; it is not part of the data.  
 (Original image courtesy of Dr. Robert A. West, NASA/JPL.)



a	b
c	d

**FIGURE 4.65**

(a)  $674 \times 674$  image of the Saturn rings showing nearly periodic interference. (b) Spectrum: The bursts of energy in the vertical axis near the origin correspond to the interference pattern. (c) A vertical notch reject filter. (d) Result of filtering. The thin black border in (c) was added for clarity; it is not part of the data. (Original image courtesy of Dr. Robert A. West, NASA/JPL.)



a b

**FIGURE 4.66**

(a) Result  
(spectrum) of  
applying a notch  
pass filter to  
the DFT of  
Fig. 4.65(a).  
(b) Spatial  
pattern obtained  
by computing the  
IDFT of (a).


---

a b

**FIGURE 4.66**

(a) Result  
(spectrum) of  
applying a notch  
pass filter to  
the DFT of  
Fig. 4.65(a).  
(b) Spatial  
pattern obtained  
by computing the  
IDFT of (a).

---




a b

**FIGURE 4.66**

(a) Result  
(spectrum) of  
applying a notch  
pass filter to  
the DFT of  
Fig. 4.65(a).  
(b) Spatial  
pattern obtained  
by computing the  
IDFT of (a).

---



# Jean Baptiste Joseph Fourier

---



(1768-1830)


Any Periodic Function can be rewritten as a Weighted Sum  
of Infinite Sinusoids of Different Frequencies.



# Sinusoid

---

$$f(x) = A \sin(2\pi u x + \varphi)$$



$A$ : Amplitude


$T$ : Period

$\varphi$ : Phase


$u$ : Frequency ( $1/T$ )

# Fourier Series

---




Square Wave  
(Period  $2\pi$ )




# Fourier Series





---




Sum of First  
8 Sinusoids



# Frequency Representation of Signal



# Fourier Transform (FT)




Represents a signal  $f(x)$  in terms of Amplitudes and Phases of its Constituent Sinusoids.

$$f(x) \longrightarrow \boxed{\text{FT}} \longrightarrow F(u)$$

# Inverse Fourier Transform (IFT)

---



Computes the signal  $f(x)$  from the Amplitudes and Phases of its Constituent Sinusoids.


$$f(x) \leftarrow \boxed{\text{IFT}} \leftarrow F(u)$$

# Maths Primer(Preliminary Concept)

- A complex number  $C$ , is defined as

$$C = R + jI$$

$$C^* = R - jI$$



- Sometimes, it is useful to represent complex numbers in polar coordinates

$$C = |C|(\cos \theta + j \sin \theta) \quad |C| = \sqrt{(R^2 + I^2)} \quad \theta = \arctan(I/R)$$

$$e^{j\theta} = \cos \theta + j \sin \theta$$

$$C = |C|e^{j\theta}$$



# Fourier Series

# Fourier Series

- $f(t)$  of a continuous variable  $t$  that is a periodic function with period  $T$  can be expressed as the sum of sines and cosines multiplied by appropriate coefficients. This sum known as a *Fourier Series*, has the form

# Fourier Series

- $f(t)$  of a continuous variable  $t$  that is a periodic function with period  $T$  can be expressed as the sum of sines and cosines multiplied by appropriate coefficients. This sum known as a *Fourier Series*, has the form

$$f(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{T}t}$$

# Fourier Series

- $f(t)$  of a continuous variable  $t$  that is a periodic function with period  $T$  can be expressed as the sum of sines and cosines multiplied by appropriate coefficients. This sum known as a *Fourier Series*, has the form

$$f(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{T}t}$$

where

$$c_n = \frac{1}{T} \int_{-T/2}^{T/2} f(t) e^{-j\frac{2\pi n}{T}t} dt \quad \text{for } n = 0, \pm 1, \pm 2, \dots$$

# Impulses

# Impulses

- A *unit impulse* of a continuous variable  $t$  located at  $t = 0$ , denoted by  $\delta(t)$ , is defined as
- And is constrained also to satisfy the identity



# Impulses

- A *unit impulse* of a continuous variable  $t$  located at  $t = 0$ , denoted by  $\delta(t)$ , is defined as

$$\delta(t) = \begin{cases} \infty & \text{if } t = 0 \\ 0 & \text{if } t \neq 0 \end{cases}$$

- And is constrained also to satisfy the identity

# Impulses

- A *unit impulse* of a continuous variable  $t$  located at  $t = 0$ , denoted by  $\delta(t)$ , is defined as

$$\delta(t) = \begin{cases} \infty & \text{if } t = 0 \\ 0 & \text{if } t \neq 0 \end{cases}$$

- And is constrained also to satisfy the identity

# Impulses

- A *unit impulse* of a continuous variable  $t$  located at  $t = 0$ , denoted by  $\delta(t)$ , is defined as

$$\delta(t) = \begin{cases} \infty & \text{if } t = 0 \\ 0 & \text{if } t \neq 0 \end{cases}$$

- And is constrained also to satisfy the identity

$$\int_{-\infty}^{\infty} \delta(t) dt = 1$$

# Sifting Property of Impulses

# Sifting Property of Impulses

- Impulse has a sifting property with respect to integration, provided that  $f(t)$  is continuous at  $t = 0$
- A more general statement involves an impulse located at an arbitrary point  $t_0$  denoted by  $\delta(t - t_0)$ ,

# Sifting Property of Impulses

- Impulse has a sifting property with respect to integration, provided that  $f(t)$  is continuous at  $t = 0$

$$\int_{-\infty}^{\infty} f(t) \delta(t) dt = f(0)$$

- A more general statement involves an impulse located at an arbitrary point  $t_0$  denoted by  $\delta(t - t_0)$ ,

# Sifting Property of Impulses

- Impulse has a sifting property with respect to integration, provided that  $f(t)$  is continuous at  $t = 0$

$$\int_{-\infty}^{\infty} f(t) \delta(t) dt = f(0)$$

- A more general statement involves an impulse located at an arbitrary point  $t_0$  denoted by  $\delta(t - t_0)$ ,



# Sifting Property of Impulses

- Impulse has a sifting property with respect to integration, provided that  $f(t)$  is continuous at  $t = 0$

$$\int_{-\infty}^{\infty} f(t) \delta(t) dt = f(0)$$

- A more general statement involves an impulse located at an arbitrary point  $t_0$  denoted by  $\delta(t - t_0)$ ,

$$\int_{-\infty}^{\infty} f(t) \delta(t - t_0) dt = f(t_0)$$

# Unit Discrete Impulse

# Unit Discrete Impulse

- Unit discrete impulse located at  $x = 0$ ,
- And is constrained to satisfy the property
- The sifting property of the discrete variable has the form
- More generally using a discrete impulse located at  $x = x_0$

# Unit Discrete Impulse

- Unit discrete impulse located at  $x = 0$ ,

$$\delta(x) = \begin{cases} 1 & x = 0 \\ 0 & x \neq 0 \end{cases}$$

- And is constrained to satisfy the property
- The sifting property of the discrete variable has the form
- More generally using a discrete impulse located at  $x = x_0$

# Unit Discrete Impulse

- Unit discrete impulse located at  $x = 0$ ,

$$\delta(x) = \begin{cases} 1 & x = 0 \\ 0 & x \neq 0 \end{cases}$$

- And is constrained to satisfy the property
- The sifting property of the discrete variable has the form
- More generally using a discrete impulse located at  $x = x_0$

# Unit Discrete Impulse

- Unit discrete impulse located at  $x = 0$ ,

$$\delta(x) = \begin{cases} 1 & x = 0 \\ 0 & x \neq 0 \end{cases}$$

- And is constrained to satisfy the property

$$\sum_{x=-\infty}^{\infty} \delta(x) = 1$$

- The sifting property of the discrete variable has the form
- More generally using a discrete impulse located at  $x = x_0$

# Unit Discrete Impulse

- Unit discrete impulse located at  $x = 0$ ,

$$\delta(x) = \begin{cases} 1 & x = 0 \\ 0 & x \neq 0 \end{cases}$$

- And is constrained to satisfy the property

$$\sum_{x=-\infty}^{\infty} \delta(x) = 1$$

- The sifting property of the discrete variable has the form
- More generally using a discrete impulse located at  $x = x_0$



# Unit Discrete Impulse

- Unit discrete impulse located at  $x = 0$ ,

$$\delta(x) = \begin{cases} 1 & x = 0 \\ 0 & x \neq 0 \end{cases}$$

- And is constrained to satisfy the property

$$\sum_{x=-\infty}^{\infty} \delta(x) = 1$$

- The sifting property of the discrete variable has the form

$$\sum_{x=-\infty}^{\infty} f(x) \delta(x) = f(0)$$

- More generally using a discrete impulse located at  $x = x_0$

# Unit Discrete Impulse

- Unit discrete impulse located at  $x = 0$ ,

$$\delta(x) = \begin{cases} 1 & x = 0 \\ 0 & x \neq 0 \end{cases}$$

- And is constrained to satisfy the property

$$\sum_{x=-\infty}^{\infty} \delta(x) = 1$$

- The sifting property of the discrete variable has the form

$$\sum_{x=-\infty}^{\infty} f(x) \delta(x) = f(0)$$

- More generally using a discrete impulse located at  $x = x_0$

# Unit Discrete Impulse

- Unit discrete impulse located at  $x = 0$ ,

$$\delta(x) = \begin{cases} 1 & x = 0 \\ 0 & x \neq 0 \end{cases}$$

- And is constrained to satisfy the property

$$\sum_{x=-\infty}^{\infty} \delta(x) = 1$$

- The sifting property of the discrete variable has the form

$$\sum_{x=-\infty}^{\infty} f(x) \delta(x) = f(0)$$

- More generally using a discrete impulse located at  $x = x_0$

$$\sum_{x=-\infty}^{\infty} f(x) \delta(x - x_0) = f(x_0)$$

# Unit Discrete Impulse

- Unit discrete impulse located at  $x = 0$ ,

$$\delta(x) = \begin{cases} 1 & x = 0 \\ 0 & x \neq 0 \end{cases}$$

- And is constrained to satisfy the property


$$\sum_{x=-\infty}^{\infty} \delta(x) = 1$$

- The sifting property of the discrete variable has the form

$$\sum_{x=-\infty}^{\infty} f(x) \delta(x) = f(0)$$

- More generally using a discrete impulse located at  $x = x_0$

$$\sum_{x=-\infty}^{\infty} f(x) \delta(x - x_0) = f(x_0)$$



**FIGURE 4.2**

A unit discrete impulse located at  $x = x_0$ . Variable  $x$  is discrete, and  $\delta$  is 0 everywhere except at  $x = x_0$ .

# Impulse Train

# Impulse Train

- Sum of infinitely many periodic impulses  $\Delta T$  units apart:

# Impulse Train


- Sum of infinitely many periodic impulses  $\Delta T$  units apart:

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} \delta(t - n\Delta T)$$

# Impulse Train

- Sum of infinitely many periodic impulses  $\Delta T$  units apart:

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} \delta(t - n\Delta T)$$






# Impulse Train

- Sum of infinitely many periodic impulses  $\Delta T$  units apart:

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} \delta(t - n\Delta T)$$



**FIGURE 4.3** An impulse train.

# Fourier Transform of functions of one continuous variable

# Fourier Transform of functions of one continuous variable

- Fourier transform of a continuous function  $f(t)$  of a continuous variable  $t$  :
- Inverse Fourier Transform

# Fourier Transform of functions of one continuous variable

- Fourier transform of a continuous function  $f(t)$  of a continuous variable  $t$  :

$$F(\mu) = \mathfrak{F}\{f(t)\} = \int_{-\infty}^{\infty} f(t) e^{-j2\pi\mu t} dt$$

- Inverse Fourier Transform

# Fourier Transform of functions of one continuous variable

- Fourier transform of a continuous function  $f(t)$  of a continuous variable  $t$  :

$$F(\mu) = \mathfrak{F}\{f(t)\} = \int_{-\infty}^{\infty} f(t) e^{-j2\pi\mu t} dt$$

- Inverse Fourier Transform

# Fourier Transform of functions of one continuous variable

- Fourier transform of a continuous function  $f(t)$  of a continuous variable  $t$  :

$$F(\mu) = \mathfrak{F}\{f(t)\} = \int_{-\infty}^{\infty} f(t) e^{-j2\pi\mu t} dt$$

- Inverse Fourier Transform

$$\mathfrak{F}^{-1}\{F(\mu)\} = f(t) = \int_{-\infty}^{\infty} F(\mu) e^{j2\pi\mu t} d\mu$$

# Fourier Transform of functions of one continuous variable

- Fourier transform of a continuous function  $f(t)$  of a continuous variable  $t$  :

$$F(\mu) = \mathfrak{F}\{f(t)\} = \int_{-\infty}^{\infty} f(t) e^{-j2\pi\mu t} dt$$


- Inverse Fourier Transform  $= \int_{-\infty}^{\infty} f(t) [\cos(2\pi\mu t) - j\sin(2\pi\mu t)] dt$

$$\mathfrak{F}^{-1}\{F(\mu)\} = f(t) = \int_{-\infty}^{\infty} F(\mu) e^{j2\pi\mu t} d\mu$$

# Example

- Example 1: Find Fourier Transform of function shown in figure below

$$\begin{aligned} F(\mu) &= \int_{-\infty}^{\infty} f(t) e^{-j2\pi\mu t} dt = \int_{-W/2}^{W/2} A e^{-j2\pi\mu t} dt \\ &= \frac{-A}{j2\pi\mu} \left[ e^{-j2\pi\mu t} \right]_{-W/2}^{W/2} = \frac{-A}{j2\pi\mu} \left[ e^{-j\pi\mu W} - e^{j\pi\mu W} \right] \\ &= \frac{A}{j2\pi\mu} \left[ e^{j\pi\mu W} - e^{-j\pi\mu W} \right] \\ &= AW \frac{\sin(\pi\mu W)}{(\pi\mu W)} \quad \sin \theta = (e^{j\theta} - e^{-j\theta})/2j. \end{aligned}$$







- The result in the last step of the preceding expression is known as *sinc* function:
- Where  $\text{sinc}(0) = 1$  and  $\text{sinc}(m) = 0$  for all other integer values of  $m$ .

- The result in the last step of the preceding expression is known as *sinc* function:


$$\text{sinc}(m) = \frac{\sin(\pi m)}{(\pi m)}$$

- Where  $\text{sinc}(0) = 1$  and  $\text{sinc}(m) = 0$  for all other integer values of  $m$ .

- The result in the last step of the preceding expression is known as *sinc* function:

$$\text{sinc}(m) = \frac{\sin(\pi m)}{(\pi m)}$$

- Where  $\text{sinc}(0) = 1$  and  $\text{sinc}(m) = 0$  for all other integer values of  $m$ .




a b c

**FIGURE 4.4** (a) A simple function; (b) its Fourier transform; and (c) the spectrum. All functions extend to infinity in both directions.

a b c


**FIGURE 4.4** (a) A simple function; (b) its Fourier transform; and (c) the spectrum. All functions extend to infinity in both directions.



a b c


**FIGURE 4.4** (a) A simple function; (b) its Fourier transform; and (c) the spectrum. All functions extend to infinity in both directions.

The locations of the zeros of both  $F(\mu)$  and  $|F(\mu)|$  are *inversely* proportional to the width,  $W$ , of the “box” function



**FIGURE 4.4** (a) A simple function; (b) its Fourier transform; and (c) the spectrum. All functions extend to infinity in both directions.






a b c

**FIGURE 4.4** (a) A simple function; (b) its Fourier transform; and (c) the spectrum. All functions extend to infinity in both directions.

The function extends to infinity for both positive and negative values of  $\mu$



a b c

**FIGURE 4.4** (a) A simple function; (b) its Fourier transform; and (c) the spectrum. All functions extend to infinity in both directions.

- Example 2: Fourier Transform of a unit impulse located at the origin

- Example 2: Fourier Transform of a unit impulse located at the origin

$$F(\mu) = \int_{-\infty}^{\infty} \delta(t) e^{-j2\pi\mu t} dt$$

- Example 2: Fourier Transform of a unit impulse located at the origin

$$\begin{aligned} F(\mu) &= \int_{-\infty}^{\infty} \delta(t) e^{-j2\pi\mu t} dt \\ &= \int_{-\infty}^{\infty} e^{-j2\pi\mu t} \delta(t) dt \end{aligned}$$

- Example 2: Fourier Transform of a unit impulse located at the origin

$$\begin{aligned} F(\mu) &= \int_{-\infty}^{\infty} \delta(t) e^{-j2\pi\mu t} dt \\ &= \int_{-\infty}^{\infty} e^{-j2\pi\mu t} \delta(t) dt \\ &= e^{-j2\pi\mu 0} = e^0 \end{aligned}$$

- Example 2: Fourier Transform of a unit impulse located at the origin

$$\begin{aligned} F(\mu) &= \int_{-\infty}^{\infty} \delta(t) e^{-j2\pi\mu t} dt \\ &= \int_{-\infty}^{\infty} e^{-j2\pi\mu t} \delta(t) dt \\ &= e^{-j2\pi\mu 0} = e^0 \\ &= 1 \end{aligned}$$

- Example 2: Fourier Transform of a unit impulse located at the origin

$$F(\mu) = \int_{-\infty}^{\infty} \delta(t) e^{-j2\pi\mu t} dt$$

$$= \int_{-\infty}^{\infty} e^{-j2\pi\mu t} \delta(t) dt$$

Sifting property of impulse function

$$= e^{-j2\pi\mu 0} = e^0$$

$$= 1$$



- Example 3: Fourier Transform of a unit impulse located at  $t = t_0$

- Example 3: Fourier Transform of a unit impulse located at  $t = t_0$

$$F(\mu) = \int_{-\infty}^{\infty} \delta(t - t_0) e^{-j2\pi\mu t} dt$$

- Example 3: Fourier Transform of a unit impulse located at  $t = t_0$

$$\begin{aligned} F(\mu) &= \int_{-\infty}^{\infty} \delta(t - t_0) e^{-j2\pi\mu t} dt \\ &= \int_{-\infty}^{\infty} e^{-j2\pi\mu t} \delta(t - t_0) dt \end{aligned}$$

- Example 3: Fourier Transform of a unit impulse located at  $t = t_0$

$$\begin{aligned} F(\mu) &= \int_{-\infty}^{\infty} \delta(t - t_0) e^{-j2\pi\mu t} dt \\ &= \int_{-\infty}^{\infty} e^{-j2\pi\mu t} \delta(t - t_0) dt \\ &= e^{-j2\pi\mu t_0} \end{aligned}$$

- Example 3: Fourier Transform of a unit impulse located at  $t = t_0$

$$\begin{aligned} F(\mu) &= \int_{-\infty}^{\infty} \delta(t - t_0) e^{-j2\pi\mu t} dt \\ &= \int_{-\infty}^{\infty} e^{-j2\pi\mu t} \delta(t - t_0) dt \\ &= e^{-j2\pi\mu t_0} \\ &= \cos(2\pi\mu t_0) - j \sin(2\pi\mu t_0) \end{aligned}$$



$$F(\mu) = \delta(\mu - \mu_0)$$

$$IFT [ F(\mu) ] = \int_{-\infty}^{\infty} \delta(\mu - \mu_0) e^{j2\pi\mu t} d\mu = f(t)$$

$$f(t) = e^{j2\pi\mu_0 t}$$

Therefore,  $FT (e^{j2\pi\mu_0 t}) = \delta(\mu - \mu_0)$

$$F(\mu) = \delta(\mu - \mu_0)$$

$$IFT [ F(\mu) ] = \int_{-\infty}^{\infty} \delta(\mu - \mu_0) e^{j2\pi\mu t} d\mu = f(t)$$

$$f(t) = e^{j2\pi\mu_0 t}$$

Therefore,  $FT (e^{j2\pi\mu_0 t}) = \delta(\mu - \mu_0)$



$$F(\mu) = \delta(\mu - \mu_0)$$

$$IFT [ F(\mu) ] = \int_{-\infty}^{\infty} \delta(\mu - \mu_0) e^{j2\pi\mu t} d\mu = f(t)$$

$$f(t) = e^{j2\pi\mu_0 t}$$

Therefore,  $FT (e^{j2\pi\mu_0 t}) = \delta(\mu - \mu_0)$

$$F(\mu) = \delta(\mu - \mu_0)$$

$$IFT [ F(\mu) ] = \int_{-\infty}^{\infty} \delta(\mu - \mu_0) e^{j2\pi\mu t} d\mu = f(t)$$

$$f(t) = e^{j2\pi\mu_0 t}$$

Therefore,  $FT (e^{j2\pi\mu_0 t}) = \delta(\mu - \mu_0)$

$$F(\mu) = \delta(\mu - \mu_0)$$

$$IFT [ F(\mu) ] = \int_{-\infty}^{\infty} \delta(\mu - \mu_0) e^{j2\pi\mu t} d\mu = f(t)$$

$$f(t) = e^{j2\pi\mu_0 t}$$

Therefore,  $FT (e^{j2\pi\mu_0 t}) = \delta(\mu - \mu_0)$



$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{\Delta T}t}$$

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{\Delta T}t}$$

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} \delta(t - n\Delta T)$$

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{\Delta T}t}$$

$$f(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{T}t}$$

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} \delta(t - n\Delta T)$$

- Example 4: Fourier Transform  $S(\mu)$  of an impulse train with period  $\Delta T$

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{\Delta T}t}$$

$$f(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{T}t}$$

where

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} \delta(t - n\Delta T)$$



- Example 4: Fourier Transform  $S(\mu)$  of an impulse train with period  $\Delta T$

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{\Delta T}t}$$

$$f(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{T}t}$$

where

$$c_n = \frac{1}{\Delta T} \int_{-\Delta T/2}^{\Delta T/2} s_{\Delta T}(t) e^{-j\frac{2\pi n}{\Delta T}t} dt$$

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} \delta(t - n\Delta T)$$


- Example 4: Fourier Transform  $S(\mu)$  of an impulse train with period  $\Delta T$

$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{\Delta T}t}$$

$$f(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{T}t}$$

where

$$c_n = \frac{1}{\Delta T} \int_{-\Delta T/2}^{\Delta T/2} s_{\Delta T}(t) e^{-j\frac{2\pi n}{\Delta T}t} dt$$



$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} \delta(t - n\Delta T)$$

- Example 4: Fourier Transform  $S(\mu)$  of an impulse train with period  $\Delta T$


$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{\Delta T}t}$$

$$f(t) = \sum_{n=-\infty}^{\infty} c_n e^{j\frac{2\pi n}{T}t}$$

where

$$c_n = \frac{1}{\Delta T} \int_{-\Delta T/2}^{\Delta T/2} s_{\Delta T}(t) e^{-j\frac{2\pi n}{\Delta T}t} dt$$

$$c_n = \frac{1}{T} \int_{-T/2}^{T/2} f(t) e^{-j\frac{2\pi n}{T}t} dt$$



$$s_{\Delta T}(t) = \sum_{n=-\infty}^{\infty} \delta(t - n\Delta T)$$



$$c_n = \frac{1}{\Delta T} \int_{-\Delta T/2}^{\Delta T/2} \delta(t) e^{-j\frac{2\pi n}{\Delta T}t} dt$$

$$\begin{aligned}
 c_n &= \frac{1}{\Delta T} \int_{-\Delta T/2}^{\Delta T/2} \delta(t) e^{-j\frac{2\pi n}{\Delta T}t} dt \\
 &= \frac{1}{\Delta T} e^0 \\
 &= \frac{1}{\Delta T}
 \end{aligned}$$

$$\begin{aligned}
 c_n &= \frac{1}{\Delta T} \int_{-\Delta T/2}^{\Delta T/2} \delta(t) e^{-j\frac{2\pi n}{\Delta T}t} dt \\
 &= \frac{1}{\Delta T} e^0 \\
 &= \frac{1}{\Delta T}
 \end{aligned}$$

The Fourier series expansion then becomes

$$s_{\Delta T}(t) = \frac{1}{\Delta T} \sum_{n=-\infty}^{\infty} e^{j\frac{2\pi n}{\Delta T}t}$$

So,  $S(\mu)$ , the Fourier transform of the periodic impulse train  $s_{\Delta T}(t)$ , is



So,  $S(\mu)$ , the Fourier transform of the periodic impulse train  $s_{\Delta T}(t)$ , is

$$S(\mu) = \mathfrak{F}\{s_{\Delta T}(t)\}$$

So,  $S(\mu)$ , the Fourier transform of the periodic impulse train  $s_{\Delta T}(t)$ , is

$$\begin{aligned} S(\mu) &= \mathfrak{F}\{s_{\Delta T}(t)\} \\ &= \mathfrak{F}\left\{\frac{1}{\Delta T} \sum_{n=-\infty}^{\infty} e^{j\frac{2\pi n}{\Delta T}t}\right\} \end{aligned}$$

So,  $S(\mu)$ , the Fourier transform of the periodic impulse train  $s_{\Delta T}(t)$ , is

$$\begin{aligned} S(\mu) &= \mathfrak{F}\{s_{\Delta T}(t)\} \\ &= \mathfrak{F}\left\{\frac{1}{\Delta T} \sum_{n=-\infty}^{\infty} e^{j\frac{2\pi n}{\Delta T}t}\right\} \\ &= \frac{1}{\Delta T} \mathfrak{F}\left\{\sum_{n=-\infty}^{\infty} e^{j\frac{2\pi n}{\Delta T}t}\right\} \end{aligned}$$

So,  $S(\mu)$ , the Fourier transform of the periodic impulse train  $s_{\Delta T}(t)$ , is

$$\begin{aligned} S(\mu) &= \mathfrak{F}\{s_{\Delta T}(t)\} \\ &= \mathfrak{F}\left\{\frac{1}{\Delta T} \sum_{n=-\infty}^{\infty} e^{j\frac{2\pi n}{\Delta T}t}\right\} \\ &= \frac{1}{\Delta T} \mathfrak{F}\left\{\sum_{n=-\infty}^{\infty} e^{j\frac{2\pi n}{\Delta T}t}\right\} \end{aligned}$$

$$\text{FT } (e^{j2\pi\mu_0 t}) = \delta(\mu - \mu_0)$$

So,  $S(\mu)$ , the Fourier transform of the periodic impulse train  $s_{\Delta T}(t)$ , is

$$\begin{aligned} S(\mu) &= \mathfrak{F}\{s_{\Delta T}(t)\} \\ &= \mathfrak{F}\left\{\frac{1}{\Delta T} \sum_{n=-\infty}^{\infty} e^{j\frac{2\pi n}{\Delta T}t}\right\} \\ &= \frac{1}{\Delta T} \mathfrak{F}\left\{\sum_{n=-\infty}^{\infty} e^{j\frac{2\pi n}{\Delta T}t}\right\} \\ &= \frac{1}{\Delta T} \sum_{n=-\infty}^{\infty} \delta\left(\mu - \frac{n}{\Delta T}\right) \end{aligned}$$

$$\text{FT } (e^{j2\pi\mu_0 t}) = \delta(\mu - \mu_0)$$

$$S(\mu) = \mathfrak{F}\left\{s_{\Delta T}(t)\right\} = \frac{1}{\Delta T} \sum_{n=-\infty}^{\infty} \delta\left(\mu - \frac{n}{\Delta T}\right)$$

$$S(\mu) = \mathfrak{F}\{s_{\Delta T}(t)\} = \frac{1}{\Delta T} \sum_{n=-\infty}^{\infty} \delta\left(\mu - \frac{n}{\Delta T}\right)$$

This fundamental result tells us that Fourier transform of an impulse train with period  $\Delta T$  is also an impulse train, whose period is  $1/\Delta T$