

Université d'Ottawa  
Faculté de génie

University of Ottawa  
Faculty of Engineering



École de science informatique et  
de génie électrique (SIGE)

School of Electrical Engineering  
and Computer Science (EECS)

CSI2110 (Fall 2021)  
**Assignment 8 (3%) – 12 points**

Due: Thursday Nov 18, 11:59PM

Late assignment policy: 1min-24hs late are accepted with 30% off; no assignments accepted after that.

Note: It is copyrighted content. You are not allowed to post any of content in public.

**Question 1. [7 points]**

Consider the following Hash table where insertions are done using the hash function

$$h(k) = 3*k + 1 \bmod 11.$$

Insert the following keys 10, 21, 3, 14, 18, 7 in an empty Hash table with 11 spots, **mark the positions (underline/square/circle a key on the position)** that are probed for each insertion (in the given table show the Hash table after each insertion). After all insertions, what is the **average number of probes** (comparisons) needed to search for an existing key in this table? In each part, collisions are solved by each of the following 3 methods specified in the part.

Hint: Here is an example when inserting different keys 1, 3 and 8 with linear probing. Each row shows the table after the insertion of each key. For each insertion we underline the positions that were probed; since every element was inserted in their home address the number of probes for each element is 1 and the average number of probes is  $(1+1+1)/3 = 1$ .

0	1	2	3	4	5	6	7	8	9	10
				<u>1</u>						
				1						<u>3</u>
			<u>8</u>	1						3

1. [2 point] Linear probing

0	1	2	3	4	5	6	7	8	9	10	
									<u>10</u>		1
									<u>10</u>	21	2
<u>3</u>									<u>10</u>	<u>21</u>	2



3	14								10	21	3
3	14	18							10	21	3
3	14	18	7						10	21	4

Average number of probes =  $(1+2+2+3+3+4)/6 = 2.5$  probes

2. [2 points] Quadrating probing

0	1	2	3	4	5	6	7	8	9	10	
									10		1
									10	21	2
3									10	21	3
3		14							10	21	3
3	18	14							10	21	2
3	18	14	7						10	21	3

Average number of probes =  $(1+2+2+3+3+3)/6 = 2.17$  probes

3. [3 points] Double hashing with secondary hash function:  $d(k) = 4 - (k \bmod 3)$ .  
Double hashing is done by  $h_j(k) = [h(k) + j \cdot d(k)] \bmod 11$ .

0	1	2	3	4	5	6	7	8	9	10
									10	
		21							10	
		21							10	3
	14	21							10	3
18	14	21							10	3
18	14	21	7						10	3

k	$h(k)$	$d(k)$	Probes
10	9	3	9
21	2	4	9 2
3	10	4	10
14	1	2	10 1
18	0	4	0
7	3	3	0 3

Average number of probes =  $(1+2+1+2+1+2)/6 = 1.5$  probes

**Question 2. [5 points]** Consider a hash table that is constructed with double hashing:

$$h(k) = [h(k) + j \cdot d(k)] \bmod 11$$

$j \cdot d(k)$ ?



The primary hash function is  $h(k) = (2k + 1) \bmod 11$  and the secondary hash function is  $d(k) = (k \text{ div } 2) + 1$  where  $\text{div}$  is integer division (e.g.  $10 \bmod 3 = 1$  and  $10 \text{ div } 3 = 3$ ).

1. [2 point] Insert the keys 13 and then 11 in the following hash table and describe the process indicating for each key the positions probed.

- ① cal the  $h(k)$   
 ② if occupied cal  $d(k)$   
 ③ Then cal the double hashing, if occupied repeat from step ②

0	1	2	3	4	5	6	7	8	9	10
					2	19	3			
					2	19	3			13
	11				2	19	3			13

K	$h_1(K)$	$d(K)$	Probe
13	10	8	5.10
11	1	6	1

2. [1 point] Delete the key 2 in the hash table below and describe the process.

0	1	2	3	4	5	6	7	8	9	10
	11				13		2			
	11				13		avail			

- Probe until found (5)  
 ① Search Key 2 → 5  
 ② Delete & replace with avail so we can tell that it was an occupied cell

3. [1 point] Search the key 11 in the hash table and describe the process.

0	1	2	3	4	5	6	7	8	9	10
16	AVAILABLE	AVAILABLE			2		3	11		

$$h(11) = (2 \cdot 11 + 1) \bmod 11 = 1$$

$$= \text{available (keep prob)}$$

$$d(11) = (11 \text{ div } 2) + 1$$

$$= 6$$

$$h_1(11) = (1 + 1 \cdot 6) \bmod 11$$

$$= 7 \rightarrow 3 \text{ (not 11 keep probing)}$$

$$h_2(11) = (1 + 2 \cdot 6) \bmod 11 = 13 \rightarrow \text{avail}$$

$$\text{Keep probing}$$

4. [1 point] Insert the key 25 in the hash table, describe the process.

0	1	2	3	4	5	6	7	8	9	10
16	AVAILABLE	AVAILABLE		25	2		3	11	4	

→  $h_3(25) = (7 + 3 \cdot 6) \bmod 11$   
 $= 2$ , occupied, keep probing  
 $h_4(25) = (7 + 5 \cdot 6) \bmod 11$   
 $= 4$ , not occupied  
 then insert at index 4  
 5 probes

$h(25) = (51) \bmod 11 = 7$ , occupied, find  $d(k)$  & keep probing  
 $d(25) = 13$   
 $h_1(25) = (7 + 13) \bmod 11$   
 $= 9$ , occupied, keep probing  
 $h_2(25) = (7 + 26) \bmod 11$   
 $= 0$ , occupied, keep probing