

Proportion of bloodstream infection due to methicillin-resistant Staphylococcus aureus (MRSA) (%)

FILTERS

EXPORT DATA in CSV format:
Right-click here & Save link



Last updated: 2025-10-22

Indicator	Proportion of bloodstream infection due to methicillin-resistant Staphylococcus aureus (MRSA) (%) ¹							
Location	2023	2022	2021	2020	2019	2018	2017	2016
Algeria	46							
Argentina	35.7	32.8	34.3	38.4	42	42.5	45	
Australia	15.8	15	16.9	17.7	18.5			
Austria	4.3	3.9	3.1	4.5	5.3	6.4	5.9	
Bahrain	53.3	42.1		34.4	40	35.5		
Bangladesh	78	88.5						
Belgium	7	4.6	4.2	6.8				
Benin	62.8	52.6						
Bhutan	5			11.7	20.8			
Bosnia and Herzegovina	28.4	26.5	20.8	23.2	10.6	16.3	25.8	18
Brazil	27.5	31.6	7	24.4	20.6	11.1		
Brunei Darussalam	5.3	12.2	14.6	12.5	12.3			
Burkina Faso	22	30.4	51.2	68.8				
Cote d'Ivoire	100			71.5				
Cambodia	54.2	81.3		68.1		48.8		
Cameroon	62.5	76	45.5					
Canada	18.3	21						
China, Hong Kong Special Administrative Region	45.7	45.8						
Colombia	47.8	45.2	45.2					
Croatia	27.4	53.8	66.7	50	24.9	26.4	28.5	
Cyprus	53	55.9	42.9	49.5	100	40.2	31.2	
Czechia	28.5	7.6	9.4	9.3	12.5	13.6	13.2	
Democratic Republic of the Congo	90.4	88.5	100					