

Name: _____

Math 40, Section _____

HW03 - Linear Systems, Span, Linear Independence, Matrix Operations

February 2, 2017

Section 2.2 Numbers: 14 , 18, 20, 22, 23*, 30 42

Section 2.3 Numbers: 8 , 16, 28, 42

Chapter 2 Review Numbers: 1f, 1h, 1j

3.1 Numbers: 8, 16, 18

2.2.14 Use elementary row operations to reduce the given matrix to (a) row echelon form and (b) reduced row echelon form

$$[M] = \begin{bmatrix} -2 & -4 & 7 \\ -3 & -6 & 10 \\ 1 & 2 & -3 \end{bmatrix}$$

■

2.2.18 Show that the given matrices are row equivalent and find a sequence of elementary row operations that will convert A into B

$$A = \begin{bmatrix} 2 & 0 & -1 \\ 1 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 3 & 1 & -1 \\ 3 & 5 & 1 \\ 2 & 2 & 0 \end{bmatrix}$$

■

2.2.20 What is the net effect of performing the following sequence of elementary row operations on a matrix (with at least two rows)?

$$R_1 + R_2, R_1 - R_2, R_2 + R_1, -R_1$$

■

2.2.22 Consider the matrix $A = \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}$. Show that any of the three types of elementary row operations can be used to create a leading 1 at the top of the first column. Which do you prefer and why?

■

2.2.23 What is the rank of each of the matrices in Exercises 2, 4, 6, and 8

$$2. \begin{bmatrix} 7 & 0 & 1 & 0 \\ 0 & 1 & -1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad 4. \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$6. \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \quad 8. \begin{bmatrix} 2 & 1 & 3 & 5 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

■

2.2.30 Use either Gaussian or Gauss-Jordan elimination to solve the system

$$-x_1 + 3x_2 - 2x_3 + 4x_4 = 0$$

$$2x_1 - 6x_2 + x_3 - 2x_4 = -3$$

$$x_1 + 3x_2 + 4x_3 - 8x_4 = 2$$

■

2.2.42 Determine what values of k have, for the following system, (a) no solution, (b) a unique solution, and (c) infinitely many solutions

$$x - 2y + 3z = 2$$

$$x + y + z = k$$

$$2x - y + 4z = k^2$$

■

2.3.8 Determine if the vector \mathbf{b} is in the span of the columns of the matrix A ;

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 5 & 6 & 7 \\ 9 & 10 & 11 \end{bmatrix} \quad \mathbf{b} = \begin{bmatrix} 4 \\ 8 \\ 12 \end{bmatrix}$$

■

| |
|---------------|
| 2.3.16 |
|---------------|

■

| |
|--------|
| 2.3.28 |
|--------|

■

| |
|--------|
| 2.3.42 |
|--------|

■







| |
|--------------|
| 3.1.8 |
|--------------|

■

| |
|---------------|
| 3.1.16 |
|---------------|

■

| |
|---------------|
| 3.1.18 |
|---------------|

■