

1: (5 points) Consider the general, linear, second-order, constant-coefficient, homogeneous differential equation $ay'' + by' + cy = 0$. Assuming $a \neq 0$, if we divide the whole equation by a , we arrive at

$$y'' + \alpha y' + \beta y = 0.$$

where $\alpha = b/a$ and $\beta = c/a$. Assume that α and β are real numbers.

- (a) For which values of α and β will we get solutions that oscillate (i.e., the roots of the characteristic equation are complex)? For which values of α and β will we get solutions that don't oscillate? For which values of α and β will we get repeated roots of the characteristic equation?
- (b) You saw before on homework 4 what values of λ must be to get solutions that decay to zero as $t \rightarrow \infty$. Now answer for what values of α and β will we get solutions that decay to zero as $t \rightarrow \infty$? For which values of α and β will we get solutions that grow exponentially (oscillatory or not) as $t \rightarrow \infty$?
- (c) Draw a plane with α and β as the two axes. Summarize the information in parts (a) and (b) graphically by showing which regions of the plane correspond to different types of behaviors.
- (d) Watch this video: <http://vimeo.com/63064216>. If the behavior of the door could be described by a linear, second-order, constant-coefficient, homogeneous ODE, what must be true about the parameters α and β ?

2: (5 points) Consider the IVP $y'' - 4y' + 4y = 2x$ with $y(0) = 5, y'(0) = 3$.

- (a) Use the method of variation of parameters find the solution to this IVP.
- (b) Now use the method of undetermined coefficients to find the solution to this IVP. (You do not have to redo the part where find the unknown constants using the initial conditions you can reuse your work from the previous part.)

3: (5 points) Consider the DE $y'' + 3y' + 2y = e^{-2x}$.

- (a) Use the method of variation of parameters find the general solution to this DE.
- (b) Now use the method of undetermined coefficients to find the general solution.
- (c) If your two solutions from part (a) and part (b) are different, reconcile them.

4: (10 points; 5 points for (a) and 5 points for (b)) For each IVP below, think about whether undetermined coefficients or variation of parameters would be more desirable to use. Solve using whichever method you think is best.

(a) $y'' - 2y' + y = \frac{e^t}{1+t^2}$ with $y(0) = y'(0) = 0$

(b) $y'' + y = 20e^{2t}$ with $y(0) = y'(0) = 0$

5: (5 points) Use the Wronskian to determine whether the following functions y_1 and y_2 are linearly independent or linearly dependent on the interval $t \in (0, 1)$.

(a) $y_1(t) = e^{5t}, y_2(t) = e^{-3t}$

(b) $y_1(t) = t^2, y_2(t) = t^4$

(c) $y_1(t) = \cos(t)\sin(t), y_2(t) = \sin 2t$

(d) $y_1(t) = e^{\alpha t}\cos(\beta t), y_2(t) = e^{\alpha t}\sin(\beta t)$ (α, β constants, $\beta \neq 0$)

6: (5 points) Consider the equation

$$y'' + p(t)y' + q(t)y = 0, t \in I$$

where $p(t)$ and $q(t)$ are continuous on I . Let y_1 and y_2 be solutions to this equation.

- (a) Prove that if y_1 and y_2 are zero at the same point in I , then they cannot be a *fundamental solution set* (i.e. basis for the set of solutions) on I .
- (b) Prove that if y_1 and y_2 have maxima or minima at the same point in I , then they cannot be a *fundamental solution set* on I .

7: (5 points) Meet with your project groups sometime before class Friday to discuss what you want to model. Be prepared to discuss your model on Friday. Each member of your group must sign your homework to verify that you met. Here are important guidelines you can follow when collaborating: <https://drive.google.com/drive/u/0/folders/17p6TQosy900yKBUCWqM1vws-c0AmyCex>