The Æther-flow Interpretation of Relativity

Higher-Derivative Quartic Branch Unit-Lapse Weighted/Homogeneous Asymptotic Closure Test

Abstract

The fixed-completion unit-lapse weighted/homogeneous route has now passed the local-closure, coercive-bootstrap, and decay/integrability stages. The next honest question is therefore no longer whether a compatible decay package can be written down, but whether that same package closes the actual nonlinear asymptotic argument in the same fixed completion $S_{\min,4}$, in the same unit-lapse spatial-harmonic gauge, and in the same weighted/homogeneous class, or whether the first genuine asymptotic obstruction appears only there.

This note performs exactly that same-class test. It first fixes explicit asymptotic variables for the three propagating sectors already isolated by the previous note:

$$U^{\pm}_{\mathrm{TL}} \equiv -|\nabla| G_\chi \pm i|\nabla|\Theta$$

for the low-frequency trace–longitudinal wave pair,

$$U^{\pm}_{\mathrm{Ein}} \equiv -2\mathbf{P}_{\mathrm{hi}}\mathbb{K} \pm i|\nabla|\mathbf{P}_{\mathrm{hi}} h$$

for the high-frequency Einstein wave block, and

$$u_\sigma \equiv \frac{\pi_\sigma}{m_S} + i\frac{\sqrt{\lambda_4}}{M_*}\Delta_\delta \sigma$$

for the quartic scalar Schrödinger channel. In the unit-lapse formulation these satisfy transport-dressed half-wave or Schrödinger equations whose nonlinear source terms are exactly the quadratic and lower-order remainders already proved to lie in L^1_t.

The decisive point is then Duhamel closure. Using the same weighted/homogeneous auxiliary estimates and the same remainder bound

$$\mathfrak{M}(t) \lesssim \varepsilon^2 (1+t)^{-2},$$

the Duhamel terms improve the decay bootstrap rather than saturating it:

$$\mathfrak{D}_{\mathrm{TL}}(t) + \mathfrak{D}_{\mathrm{Ein,hi}}(t) \lesssim \varepsilon(1+t)^{-1}, \qquad \mathfrak{D}_\sigma(t) \lesssim \varepsilon(1+t)^{-3/2}.$$

At the same time the coercive energy obeys an integrable-growth inequality

$$\frac{d}{dt}\mathfrak{F}_N^{\mathrm{ul,wh}}(t) \le C_N \mathfrak{M}(t)\mathfrak{F}_N^{\mathrm{ul,wh}}(t),$$

so the weighted/homogeneous energy remains globally small and the same-class bootstrap closes.

The asymptotic verdict is therefore positive but slightly sharper than "ordinary decay." The perturbation does decay to the flat background in the same weighted/homogeneous norms, and

no first genuine asymptotic obstruction appears. However the exact profile is best described by a *mild modified asymptotic profile*: the limiting radiation is measured after conjugation by the transport-dressed unit-lapse linear evolution rather than by the flat free wave/Schrödinger groups in fixed coordinates. That modification is generated only by the already-recorded t^{-1} unit-lapse shift transport and does not force any deeper completion or auxiliary redesign. At present disciplined scope, the same-class asymptotic/nonlinear-stability closure succeeds.

1. Introduction

The fixed-completion unit-lapse route has now been narrowed repeatedly without yet forcing a deeper redesign. The old maximal-gauge lapse obstruction was removed at equation level by the gauge-first reformulation. The direct unweighted (K, χ, β) closure package then failed for a different reason, and the two smallest unweighted repairs also failed. The weighted/homogeneous reformulation changed the picture again: the longitudinal slot closes locally in a derivative-level class, the corrected weighted/homogeneous energy closes a $T \sim \varepsilon^{-1}$ coercive bootstrap, and the dedicated decay/integrability note shows that the corresponding quadratic and lower-order nonlinear remainder is L^1_t in that same class.

That leaves one genuinely narrower but decisive burden. The decay note still treated the linear-compatible decay package as a compatible bootstrap regime. The immediate next question is whether that same package actually closes the nonlinear asymptotic argument in the same class, or whether the first genuine obstruction only appears once the Duhamel terms and asymptotic profiles themselves are analyzed.

Framework and claim boundary. Æther / Æther-flow framework denotes the broader Æther / Æther-flow framework built on the claims that the Æther is the underlying four-dimensional substrate of reality, the Æther-flow is its intrinsic ordered motion, observed three-dimensional space is the local experiential slice of that deeper substrate, S-time is the experienced order of change arising from matter, light, and the Æther-flow, and observed expansion is the three-dimensional appearance of deeper four-dimensional ordered motion. Within that framework, Æther-flow Interpretation of Relativity denotes the exact-closure theory in which the effective gravitational dynamics are adopted to be exactly Einsteinian with universal matter coupling. In the active sequence, *adoption* means use of that established relativistic dynamics without claiming substrate derivation, while *derivation* is reserved for a first-principles recovery from explicit substrate variables.

The present note stays within that boundary. It does not change the explicit completion $S_{\mathrm{min},4}$. It does not change the unit-lapse spatial-harmonic gauge. It does not introduce a new longitudinal norm, a deeper auxiliary architecture, or a different nonlinear completion. It asks only whether the already-recorded weighted/homogeneous decay/integrability package closes the actual same-class asymptotic bootstrap and, if it does, what the asymptotic verdict is.

2. Asymptotic Bootstrap Regime and Profile Variables

Work in the same unit-lapse spatial-harmonic gauge

$$N \equiv 1, \qquad V^i(\gamma) = 0, \tag{1}$$

with reduced unknowns

$$(\gamma_{ij}, \Sigma_{ij}, K; \beta^i; \sigma, \pi_\sigma; \psi, \pi_\psi) \tag{2}$$

and slice-wise auxiliary variables (Q^I, χ, W_i^T).

Set

$$h_{ij} \equiv \gamma_{ij} - \delta_{ij}, \qquad \vartheta \equiv \frac{1}{2}\delta^{ij} h_{ij}, \qquad \mathbb{K}_{ij} \equiv \Sigma_{ij} + \frac{1}{3}\delta_{ij} K. \tag{3}$$

As in the previous weighted/homogeneous notes, define

$$F_\chi \equiv K - \mathcal{N}_\chi[\gamma, \Sigma, K, \beta, \sigma, \pi_\sigma, Q, \chi, W^T, \psi, \pi_\psi], \tag{4}$$

and the low-frequency trace–longitudinal variables

$$\Theta \equiv \mathbf{P}_{\mathrm{lo}} \vartheta, \qquad G_\chi \equiv |\nabla|^{-1} \mathbf{P}_{\mathrm{lo}} F_\chi. \tag{5}$$

For the quartic scalar channel, keep

$$c_4 \equiv \frac{\sqrt{\lambda_4}}{m_S M_*}, \qquad u_\sigma \equiv \frac{\pi_\sigma}{m_S} + i \frac{\sqrt{\lambda_4}}{M_*} \Delta_\delta \sigma. \tag{6}$$

Definition 1 (Asymptotic profile variables)**.** *Define the same-class asymptotic variables*

$$U_{\mathrm{TL}}^{\pm} \equiv -|\nabla| G_\chi \pm i |\nabla| \Theta, \tag{7}$$

$$U_{\mathrm{Ein}}^{\pm} \equiv -2\mathbf{P}_{\mathrm{hi}} \mathbb{K} \pm i |\nabla| \mathbf{P}_{\mathrm{hi}} h, \tag{8}$$

$$u_\sigma \equiv \frac{\pi_\sigma}{m_S} + i \frac{\sqrt{\lambda_4}}{M_*} \Delta_\delta \sigma. \tag{9}$$

Remark 1. *Definition 1 does not change the function class. It only diagonalizes the three propagating sectors already isolated in the previous note: the low-frequency trace–longitudinal wave pair, the high-frequency Einstein wave block, and the quartic scalar Schrödinger channel.*

Definition 2 (Same-class asymptotic bootstrap interval)**.** *Fix $N \geq 6$, $0 < \varepsilon \ll 1$, and $C_* \geq 1$. An interval $[0, T]$ is said to lie in the* same-class weighted/homogeneous asymptotic bootstrap regime *if:*

1. *the strict positivity assumptions*

$$\widehat{M}_{IJ} \xi^I \xi^J \geq m_{Q,\min}^2 |\xi|^2, \qquad \kappa_\theta \geq \kappa_{\theta,\min} > 0, \qquad \kappa_\Omega \geq \kappa_{\Omega,\min} > 0 \tag{10}$$

   hold on every slice

2. *the corrected weighted/homogeneous energy satisfies*

$$\sup_{0 \leq t \leq T} \mathfrak{F}_N^{\mathrm{ul,wh}}(t) \leq 4\varepsilon^2 \tag{11}$$

3. *the decay norms of the previous note obey*

$$\mathfrak{D}_{\mathrm{TL}}(t) \leq 4C_* \varepsilon (1+t)^{-1}, \tag{12}$$

$$\mathfrak{D}_{\mathrm{Ein,hi}}(t) \leq 4C_* \varepsilon (1+t)^{-1}, \tag{13}$$

$$\mathfrak{D}_\sigma(t) \leq 4C_* \varepsilon (1+t)^{-3/2} \tag{14}$$

for all $t \in [0, T]$

4. *the low-order dispersive seed size*

$$\mathfrak{S}_0 \equiv \sum_{\pm} \|U^{\pm}_{\mathrm{TL}}(0)\|_{W^{2,1}\cap H^{N-2}} + \sum_{\pm} \|U^{\pm}_{\mathrm{Ein}}(0)\|_{W^{2,1}\cap H^{N-2}} + \|u_\sigma(0)\|_{W^{3,1}\cap H^{N-2}} \tag{15}$$

satisfies

$$\mathfrak{S}_0 \le \varepsilon. \tag{16}$$

Remark 2. *Item* (16) *is not a new closure norm for the longitudinal sector and not a redesign of the weighted/homogeneous class. It is only the standard low-order dispersive seed needed to launch the linear decay rates already identified by the previous note. The same weighted/homogeneous energy and the same decay norms remain the actual bootstrap quantities.*

Definition 3 (Quadratic time-integrable coefficient)**.** *Let*

$$\begin{aligned}\mathfrak{M}(t) \equiv\ & \mathfrak{D}_{\mathrm{TL}}(t)^2 + \mathfrak{D}_{\mathrm{Ein,hi}}(t)^2 + \mathfrak{D}_\sigma(t)^2 \\ & + \mathfrak{D}_{\mathrm{aux}}(t)\Big(\mathfrak{D}_{\mathrm{TL}}(t) + \mathfrak{D}_{\mathrm{Ein,hi}}(t) + \mathfrak{D}_\sigma(t)\Big),\end{aligned} \tag{17}$$

where

$$\mathfrak{D}_{\mathrm{aux}}(t) \equiv \|\beta(t)\|_{W^{2,\infty}} + \|Q(t)\|_{W^{2,\infty}} + \|\nabla\chi(t)\|_{W^{1,\infty}} + \|W^T(t)\|_{W^{2,\infty}}. \tag{18}$$

Proposition 1 (Auxiliary decay inherited from the same class)**.** *There exists* $C_N > 0$ *such that every solution in the asymptotic bootstrap regime satisfies*

$$\mathfrak{D}_{\mathrm{aux}}(t) \le C_N\Big(\mathfrak{D}_{\mathrm{TL}}(t) + \mathfrak{D}_{\mathrm{Ein,hi}}(t) + \mathfrak{D}_\sigma(t)\Big) \tag{19}$$

for all $t \in [0, T]$*. In particular,*

$$\mathfrak{D}_{\mathrm{aux}}(t) \le C_N C_* \varepsilon (1+t)^{-1}. \tag{20}$$

Proof. This is exactly the auxiliary decay inheritance already proved in the weighted/homogeneous decay/integrability note. The shift, Q-sector, and transverse ordered-flow estimates are unchanged, while the low-frequency longitudinal part is controlled by G_χ together with derivative-level elliptic closure. □

Corollary 1 (Time-integrability of the nonlinear coefficient)**.** *On every asymptotic bootstrap interval,*

$$\mathfrak{M}(t) \le C_N C_*^2 \varepsilon^2 (1+t)^{-2}, \qquad \int_0^T \mathfrak{M}(s)\, ds \le C_N C_*^2 \varepsilon^2. \tag{21}$$

Proof. Combine (12)–(14) with Proposition 1. This is the same L^1_t-remainder package already recorded in the previous note. □

3. Transport-Dressed Linear Blocks

The asymptotic variables of Definition 1 diagonalize the principal propagating sectors modulo the already-recorded quadratic and lower-order remainder.

Proposition 2 (Principal equations for the asymptotic variables)**.** *On every asymptotic bootstrap interval, the variables $U^{\pm}_{\mathrm{TL}}$, $U^{\pm}_{\mathrm{Ein}}$, and u_σ satisfy*

$$(\partial_t - \mathcal{L}_\beta)U^{\pm}_{\mathrm{TL}} = \pm i|\nabla|\, U^{\pm}_{\mathrm{TL}} + \mathcal{N}^{\pm}_{\mathrm{TL}}, \tag{22}$$

$$(\partial_t - \mathcal{L}_\beta)U^{\pm}_{\mathrm{Ein}} = \pm i|\nabla|\, U^{\pm}_{\mathrm{Ein}} + \mathcal{N}^{\pm}_{\mathrm{Ein}}, \tag{23}$$

$$(\partial_t - \mathcal{L}_\beta)u_\sigma = ic_4\Delta_\delta u_\sigma + \mathcal{N}_\sigma. \tag{24}$$

Moreover,

$$\sum_{\pm} \|\mathcal{N}^{\pm}_{\mathrm{TL}}(t)\|_{H^{N-3}} + \sum_{\pm} \|\mathcal{N}^{\pm}_{\mathrm{Ein}}(t)\|_{H^{N-3}} + \|\mathcal{N}_\sigma(t)\|_{H^{N-3}} \le C_N\, \mathfrak{M}(t) \tag{25}$$

for all $t \in [0, T]$.

Proof. For the low-frequency pair, the previous note already proved

$$(\partial_t - \mathcal{L}_\beta)\Theta = -|\nabla|G_\chi + \mathbf{P}_{\mathrm{lo}}\mathcal{Q}_\vartheta, \qquad (\partial_t - \mathcal{L}_\beta)G_\chi = |\nabla|\Theta + |\nabla|^{-1}\mathbf{P}_{\mathrm{lo}}\mathcal{R}_\chi.$$

Applying the linear combinations in (7) yields (22), with $\mathcal{N}^{\pm}_{\mathrm{TL}}$ made of $\mathbf{P}_{\mathrm{lo}}\mathcal{Q}_\vartheta$ and $|\nabla|^{-1}\mathbf{P}_{\mathrm{lo}}\mathcal{R}_\chi$.

For the high-frequency Einstein block, the previous note already isolated the flat principal wave system for $\mathbf{P}_{\mathrm{hi}}h$. Since $\partial_t h = -2\mathbb{K}+$ transport and lower-order terms, the half-wave variables $U^{\pm}_{\mathrm{Ein}}$ satisfy (23) after collecting the same Ricci, transport, and auxiliary substitutions into $\mathcal{N}^{\pm}_{\mathrm{Ein}}$.

For the scalar channel, Proposition 4 of the decay note rewrites the quartic scalar block as a Schrödinger equation for u_σ up to the same lower-order remainder terms, giving (24).

Finally, the previous decay/integrability note proves that every quadratic and lower-order source term left after extracting the principal wave and Schrödinger blocks is bounded by the same time-integrable coefficient $\mathfrak{M}(t)$. That is exactly (25). □

Remark 3. *Proposition 2 identifies the exact asymptotic issue. No new obstruction enters through the principal operators themselves. The only possible same-class failure would now have to come from the Duhamel terms generated by $\mathcal{N}^{\pm}_{\mathrm{TL}}$, $\mathcal{N}^{\pm}_{\mathrm{Ein}}$, and $\mathcal{N}_\sigma$.*

Definition 4 (Transport-dressed linear evolution families)**.** *Let*

$$\mathcal{U}^{\beta}_{\mathrm{TL},\pm}(t,s), \qquad \mathcal{U}^{\beta}_{\mathrm{Ein},\pm}(t,s), \qquad \mathcal{U}^{\beta}_{\sigma}(t,s)$$

denote the evolution families for the homogeneous principal equations

$$(\partial_t - \mathcal{L}_\beta)U = \pm i|\nabla|U, \tag{26}$$

$$(\partial_t - \mathcal{L}_\beta)u = ic_4\Delta_\delta u. \tag{27}$$

Proposition 3 (Linear dispersive bounds for the transport-dressed principal blocks)**.** *There exists $C_N > 0$ such that on every asymptotic bootstrap interval and for $0 \le s \le t \le T$,*

$$\|\mathcal{U}^{\beta}_{\mathrm{TL},\pm}(t,s)f\|_{W^{1,\infty}} \le C_N(1+t-s)^{-1}\|f\|_{W^{2,1}\cap H^{N-2}}, \tag{28}$$

$$\|\mathcal{U}^{\beta}_{\mathrm{Ein},\pm}(t,s)f\|_{W^{2,\infty}} \le C_N(1+t-s)^{-1}\|f\|_{W^{2,1}\cap H^{N-2}}, \tag{29}$$

$$\|\mathcal{U}^{\beta}_{\sigma}(t,s)f\|_{W^{1,\infty}} \le C_N(1+t-s)^{-3/2}\|f\|_{W^{3,1}\cap H^{N-2}}. \tag{30}$$

Proof. At the flat exact-closure background, these are exactly the standard three-dimensional wave and Schrödinger dispersive estimates already identified in the previous note. On the present bootstrap interval the coefficients remain small in the same weighted/homogeneous class, and the only nonintegrable coefficient contribution is the already-recorded unit-lapse transport by β. Standard perturbative parametrix estimates for small variable-coefficient wave and Schrödinger operators in spatial-harmonic coordinates therefore preserve the same decay rates for the transport-dressed evolution families $\mathcal{U}^\beta$. □

Proposition 4 (Duhamel representations)**.** *For every $t \in [0, T]$,*

$$U^{\pm}_{\mathrm{TL}}(t) = \mathcal{U}^{\beta}_{\mathrm{TL},\pm}(t,0)U^{\pm}_{\mathrm{TL}}(0) + \int_0^t \mathcal{U}^{\beta}_{\mathrm{TL},\pm}(t,s)\mathcal{N}^{\pm}_{\mathrm{TL}}(s)\,ds, \tag{31}$$

$$U^{\pm}_{\mathrm{Ein}}(t) = \mathcal{U}^{\beta}_{\mathrm{Ein},\pm}(t,0)U^{\pm}_{\mathrm{Ein}}(0) + \int_0^t \mathcal{U}^{\beta}_{\mathrm{Ein},\pm}(t,s)\mathcal{N}^{\pm}_{\mathrm{Ein}}(s)\,ds, \tag{32}$$

$$u_\sigma(t) = \mathcal{U}^{\beta}_{\sigma}(t,0)u_\sigma(0) + \int_0^t \mathcal{U}^{\beta}_{\sigma}(t,s)\mathcal{N}_\sigma(s)\,ds. \tag{33}$$

Proof. This is the usual variation-of-constants formula for the nonautonomous principal systems of Proposition 2. □

4. Duhamel Closure of the Same-Class Decay Bootstrap

The remaining issue is now exactly whether the Duhamel terms in Proposition 4 preserve the same weighted/homogeneous decay rates.

Proposition 5 (Duhamel terms close in the same decay norms)**.** *There exists $C_N > 0$ such that every solution in the asymptotic bootstrap regime satisfies*

$$\mathfrak{D}_{\mathrm{TL}}(t) \le C_N\varepsilon(1+t)^{-1} + C_N\int_0^t (1+t-s)^{-1}\mathfrak{M}(s)\,ds, \tag{34}$$

$$\mathfrak{D}_{\mathrm{Ein,hi}}(t) \le C_N\varepsilon(1+t)^{-1} + C_N\int_0^t (1+t-s)^{-1}\mathfrak{M}(s)\,ds, \tag{35}$$

$$\mathfrak{D}_{\sigma}(t) \le C_N\varepsilon(1+t)^{-3/2} + C_N\int_0^t (1+t-s)^{-3/2}\mathfrak{M}(s)\,ds \tag{36}$$

for all $t \in [0, T]$.

Proof. Apply the dispersive estimates of Proposition 3 to the homogeneous terms in Proposition 4. The initial-data contributions are bounded by $C_N\mathfrak{S}_0$, hence by $C_N\varepsilon$.

For the inhomogeneous terms, use (25) together with the same dispersive estimates. Since Θ, G_χ, $\mathbf{P}_{\mathrm{hi}}h$, $\mathbf{P}_{\mathrm{hi}}\mathbb{K}$, π_σ, and $D^2\sigma$ are recovered from $U^{\pm}_{\mathrm{TL}}$, $U^{\pm}_{\mathrm{Ein}}$, and u_σ by bounded Fourier multipliers, the corresponding decay norms satisfy (34)–(36). □

Proposition 6 (Convolution improvement)**.** *There exists $C_N > 0$ such that on every asymptotic*

bootstrap interval,

$$\int_0^t (1+t-s)^{-1}\mathfrak{M}(s)\,ds \le C_N C_*^2 \varepsilon^2 (1+t)^{-1}, \tag{37}$$

$$\int_0^t (1+t-s)^{-3/2}\mathfrak{M}(s)\,ds \le C_N C_*^2 \varepsilon^2 (1+t)^{-3/2} \tag{38}$$

for all $t \in [0,T]$.

Proof. By Corollary 1, $\mathfrak{M}(s) \le C_N C_*^2 \varepsilon^2 (1+s)^{-2}$. Therefore

$$\int_0^t (1+t-s)^{-1}\mathfrak{M}(s)\,ds \lesssim C_N C_*^2 \varepsilon^2 \int_0^t (1+t-s)^{-1}(1+s)^{-2}\,ds.$$

Split the integral into $[0,t/2]$ and $[t/2,t]$. On the first subinterval, $(1+t-s)^{-1} \lesssim (1+t)^{-1}$ and $\int_0^{t/2}(1+s)^{-2}\,ds \lesssim 1$. On the second subinterval, $(1+s)^{-2} \lesssim (1+t)^{-2}$, so

$$\int_{t/2}^t (1+t-s)^{-1}(1+s)^{-2}\,ds \lesssim (1+t)^{-2}\int_0^{t/2}(1+r)^{-1}\,dr \lesssim (1+t)^{-1}.$$

This proves (37). The Schrödinger estimate (38) is easier because the kernel $(1+t-s)^{-3/2}$ is integrable near $s=t$. □

Theorem 1 (Improved same-class decay bounds)**.** *Fix* $N \ge 6$. *There exist* $C_* \ge 1$ *and* $\varepsilon_\sharp > 0$ *such that the following holds. If a smooth solution on* $[0,T]$ *lies in the asymptotic bootstrap regime of Definition 2 with* $0 < \varepsilon \le \varepsilon_\sharp$, *then*

$$\mathfrak{D}_{\mathrm{TL}}(t) \le 2C_*\varepsilon(1+t)^{-1}, \tag{39}$$

$$\mathfrak{D}_{\mathrm{Ein,hi}}(t) \le 2C_*\varepsilon(1+t)^{-1}, \tag{40}$$

$$\mathfrak{D}_{\sigma}(t) \le 2C_*\varepsilon(1+t)^{-3/2} \tag{41}$$

for all $t \in [0,T]$. *Consequently*

$$\mathfrak{D}_{\mathrm{aux}}(t) \le C_N\varepsilon(1+t)^{-1}. \tag{42}$$

Proof. Insert Proposition 6 into Proposition 5. This gives

$$\mathfrak{D}_{\mathrm{TL}}(t) + \mathfrak{D}_{\mathrm{Ein,hi}}(t) \le C_N\varepsilon(1+t)^{-1} + C_N C_*^2\varepsilon^2(1+t)^{-1},$$

and

$$\mathfrak{D}_{\sigma}(t) \le C_N\varepsilon(1+t)^{-3/2} + C_N C_*^2\varepsilon^2(1+t)^{-3/2}.$$

Choose C_* so that $C_N \le C_*$, and then choose $\varepsilon_\sharp$ so that $C_N C_*^2 \varepsilon_\sharp \le C_*$. This yields (39)–(41). Equation (42) follows from Proposition 1. □

Remark 4. *Theorem 1 is the actual nonlinear closure step missing from the previous note. The recorded decay package is no longer just a compatible bootstrap assumption. It improves itself under the same weighted/homogeneous remainder bounds and therefore closes as a genuine nonlinear dispersive bootstrap.*

5. Global Weighted/Homogeneous Asymptotic Closure

The same-class decay improvement must now be combined with the weighted/homogeneous energy estimate to obtain global continuation.

Proposition 7 (Integrable-growth energy inequality)**.** *There exists $C_N > 0$ such that on every asymptotic bootstrap interval,*

$$\frac{d}{dt}\mathfrak{F}_N^{\mathrm{ul,wh}}(t) \le C_N \mathfrak{M}(t)\mathfrak{F}_N^{\mathrm{ul,wh}}(t) \tag{43}$$

for all $t \in [0, T]$.

Proof. The weighted/homogeneous coercive-bootstrap note already established the quadratic differential inequality

$$\frac{d}{dt}\mathfrak{F}_N^{\mathrm{ul,wh}}(t) \le C_N \big(\mathfrak{F}_N^{\mathrm{ul,wh}}(t)\big)^{3/2}.$$

The decay/integrability note sharpened the same remainder structure by showing that, after extraction of the principal wave and Schrödinger blocks, the differentiated quadratic remainder is pointwise bounded by $\mathfrak{M}(t)\mathfrak{F}_N^{\mathrm{ul,wh}}(t)$. That is exactly (43). □

Corollary 2 (Uniform global weighted/homogeneous energy bound)**.** *There exists $\varepsilon_\flat > 0$ such that if $0 < \varepsilon \le \varepsilon_\flat$ and the solution lies in the asymptotic bootstrap regime on $[0, T]$, then*

$$\sup_{0\le t\le T} \mathfrak{F}_N^{\mathrm{ul,wh}}(t) \le 2\varepsilon^2. \tag{44}$$

Proof. Integrating (43) and using Corollary 1 gives

$$\mathfrak{F}_N^{\mathrm{ul,wh}}(t) \le \mathfrak{F}_N^{\mathrm{ul,wh}}(0) \exp\Big(C_N \int_0^t \mathfrak{M}(s)\, ds\Big) \le \varepsilon^2 \exp\Big(C_N C_*^2 \varepsilon^2\Big).$$

For ε sufficiently small, the exponential factor is at most 2. □

Theorem 2 (Same-class asymptotic bootstrap closes globally)**.** *Fix $N \ge 6$. There exist $\varepsilon_{\mathrm{asy}} > 0$ and $C_* \ge 1$ such that the following holds. Let compatible asymptotically flat initial data for the fixed-completion unit-lapse spatial-harmonic system satisfy the hypotheses of the weighted/homogeneous local theorem together with*

$$\mathfrak{F}_N^{\mathrm{ul,wh}}(0) \le \varepsilon^2, \qquad \mathfrak{S}_0 \le \varepsilon, \qquad 0 < \varepsilon \le \varepsilon_{\mathrm{asy}}. \tag{45}$$

Then the corresponding solution exists for all $t \ge 0$ and satisfies

$$\sup_{t\ge 0} \mathfrak{F}_N^{\mathrm{ul,wh}}(t) \le 2\varepsilon^2, \tag{46}$$

$$\mathfrak{D}_{\mathrm{TL}}(t) + \mathfrak{D}_{\mathrm{Ein,hi}}(t) \le 2C_*\varepsilon(1+t)^{-1}, \tag{47}$$

$$\mathfrak{D}_\sigma(t) \le 2C_*\varepsilon(1+t)^{-3/2}, \tag{48}$$

$$\mathfrak{D}_{\mathrm{aux}}(t) \le C_N\varepsilon(1+t)^{-1} \tag{49}$$

for all $t \ge 0$.

Proof. By the weighted/homogeneous local theorem, a nonempty short-time interval of existence is available. Because the solution is continuous in the closed weighted/homogeneous norms and $(1+t)^{-1}, (1+t)^{-3/2}$ are $O(1)$ at $t=0$, the asymptotic bootstrap assumptions hold on a sufficiently small interval after increasing C_* if necessary.

Let T_* be the supremum of times for which the asymptotic bootstrap regime holds. On every interval $[0,T] \subset [0,T_*)$, Corollary 2 improves the energy bootstrap (11), and Theorem 1 improves the decay bootstrap (12)–(14). Hence the bootstrap assumptions improve strictly and therefore extend past T_* unless $T_* = \infty$.

The weighted/homogeneous local continuation theorem then gives global existence because the closed weighted/homogeneous norm remains finite and the strict positivity regime persists under smallness. This proves (46)–(49). □

Corollary 3 (No first genuine asymptotic obstruction in the same class)**.** *At the present recorded scope of the fixed completion $S_{\min,4}$, the same weighted/homogeneous unit-lapse class does* not *encounter its first genuine obstruction at the asymptotic-bootstrap stage.*

Proof. Theorem 2 closes the nonlinear asymptotic bootstrap globally in the same class. □

6. Asymptotic Profiles and Verdict

The previous section closes the nonlinear bootstrap. What remains is to decide the correct asymptotic description.

Definition 5 (Transport-dressed asymptotic profiles)**.** *Define the profile variables*

$$\mathcal{A}^{\pm}_{\mathrm{TL}}(t) \equiv \mathcal{U}^{\beta}_{\mathrm{TL},\pm}(0,t) U^{\pm}_{\mathrm{TL}}(t), \tag{50}$$

$$\mathcal{A}^{\pm}_{\mathrm{Ein}}(t) \equiv \mathcal{U}^{\beta}_{\mathrm{Ein},\pm}(0,t) U^{\pm}_{\mathrm{Ein}}(t), \tag{51}$$

$$\mathcal{A}_{\sigma}(t) \equiv \mathcal{U}^{\beta}_{\sigma}(0,t) u_{\sigma}(t). \tag{52}$$

Proposition 8 (Cauchy property of the asymptotic profiles)**.** *There exist limits*

$$\mathcal{A}^{\pm}_{\mathrm{TL},\infty}, \qquad \mathcal{A}^{\pm}_{\mathrm{Ein},\infty}, \qquad \mathcal{A}_{\sigma,\infty}$$

in H^{N-3} such that

$$\|\mathcal{A}^{\pm}_{\mathrm{TL}}(t) - \mathcal{A}^{\pm}_{\mathrm{TL},\infty}\|_{H^{N-3}} \le C_N \varepsilon^2 (1+t)^{-1}, \tag{53}$$

$$\|\mathcal{A}^{\pm}_{\mathrm{Ein}}(t) - \mathcal{A}^{\pm}_{\mathrm{Ein},\infty}\|_{H^{N-3}} \le C_N \varepsilon^2 (1+t)^{-1}, \tag{54}$$

$$\|\mathcal{A}_{\sigma}(t) - \mathcal{A}_{\sigma,\infty}\|_{H^{N-3}} \le C_N \varepsilon^2 (1+t)^{-1} \tag{55}$$

for all $t \ge 0$.

Proof. Differentiate (50)–(52) in time and use Proposition 2. Since $\mathcal{U}^{\beta}(0,t)$ solves the homogeneous adjoint transport-dressed linear problem,

$$\partial_t \mathcal{A}^{\pm}_{\mathrm{TL}}(t) = \mathcal{U}^{\beta}_{\mathrm{TL},\pm}(0,t) \mathcal{N}^{\pm}_{\mathrm{TL}}(t),$$

and similarly for the Einstein and scalar profiles. The evolution families are bounded on H^{N-3}, while (25) and Corollary 1 imply

$$\|\partial_t \mathcal{A}^{\pm}_{\mathrm{TL}}(t)\|_{H^{N-3}} + \|\partial_t \mathcal{A}^{\pm}_{\mathrm{Ein}}(t)\|_{H^{N-3}} + \|\partial_t \mathcal{A}_{\sigma}(t)\|_{H^{N-3}} \le C_N \mathfrak{M}(t) \le C_N \varepsilon^2 (1+t)^{-2}.$$

Hence the profiles are Cauchy in H^{N-3} and converge. Integrating from t to ∞ gives the rates (53)–(55). □

Remark 5. *Proposition 8 is exactly where the L^1_t nonlinear remainder is used in its final asymptotic role. The Duhamel defect is not merely integrable enough to preserve the decay rates. It is integrable enough to produce actual limiting same-class profiles.*

Theorem 3 (Asymptotic verdict for the fixed weighted/homogeneous unit-lapse class)**.** *Under the hypotheses of Theorem 2, the same-class asymptotic closure has the following verdict:*

1. *the perturbation remains globally small and decays to the flat exact-closure background in the same weighted/homogeneous norms*
2. *the low-frequency trace–longitudinal wave pair and the high-frequency Einstein block admit limiting transport-dressed wave profiles $\mathcal{A}^{\pm}_{\mathrm{TL},\infty}$ and $\mathcal{A}^{\pm}_{\mathrm{Ein},\infty}$*
3. *the quartic scalar channel admits a limiting transport-dressed Schrödinger profile $\mathcal{A}_{\sigma,\infty}$*
4. *the correct asymptotic alternative is therefore* a mild modified asymptotic profile*, not a first genuine obstruction*
5. *the modification is generated only by the already-recorded t^{-1} unit-lapse transport part $\mathcal{L}_\beta$; no new long-range nonlinear phase, no nonintegrable Duhamel defect, and no same-class norm failure appear*
6. *consequently a deeper completion change or auxiliary redesign is* not *forced at this asymptotic stage.*

Proof. Items (1)–(3) are exactly Theorem 2 together with Proposition 8. Since $\mathfrak{D}_{\mathrm{TL}}(t)$, $\mathfrak{D}_{\mathrm{Ein,hi}}(t)$, and $\mathfrak{D}_{\sigma}(t)$ all vanish as $t \to \infty$, the perturbation decays to the flat background in the same weighted/homogeneous norms.

What prevents the stronger statement of literal unmodified free scattering in the original fixed spatial coordinates is the transport term $\mathcal{L}_\beta$, whose coefficient decays only at the wave rate t^{-1}. That coefficient is too large to be ignored completely in the profile description, but it has already been incorporated into the principal evolution families $\mathcal{U}^\beta$. After that transport dressing, the nonlinear remainder is L^1_t and the profiles converge. Therefore the correct verdict is a mild modified asymptotic profile rather than an obstruction. □

Corollary 4 (Not a redesign yet)**.** *The immediate next step after the present note is not a deeper completion change, not a different gauge, and not a different weighted class. At current disciplined scope, the same-class asymptotic/nonlinear-stability closure succeeds.*

Proof. This is the direct content of Theorem 3. □

7. What This Step Establishes

The present manuscript establishes the following.

1. It turns the previously recorded linear-compatible decay package into a closed nonlinear bootstrap on the same fixed completion $S_{\min,4}$, in the same unit-lapse spatial-harmonic gauge, and in the same weighted/homogeneous class.

2. It fixes explicit asymptotic variables for the low-frequency trace–longitudinal wave pair, the high-frequency Einstein wave block, and the quartic scalar Schrödinger channel.

3. It proves that the Duhamel terms close in the same weighted/homogeneous norms because the nonlinear remainder coefficient $\mathfrak{M}(t)$ is L^1_t.

4. It upgrades the weighted/homogeneous energy estimate from a $T \sim \varepsilon^{-1}$ coercive bootstrap to a global same-class bound.

5. It produces limiting transport-dressed asymptotic profiles for all three propagating sectors.

6. It decides the asymptotic verdict: the correct outcome is a mild modified asymptotic profile, not a first true asymptotic obstruction.

It does not establish the following.

1. It does not claim a first-principles substrate derivation of Einsteinian gravity beyond the already recorded adoption line.

2. It does not remove the strict positivity assumptions $\kappa_\theta > 0$, $\kappa_\Omega > 0$, and positive Q-sector coercivity from the current branch.

3. It does not show that every conceivable weaker completion or weaker auxiliary package would enjoy the same asymptotic closure.

4. It does not say that the transport-dressed modified profile is unique across gauges or that the same result would automatically persist under a different nonlinear completion.

8. Conclusion

The weighted/homogeneous unit-lapse route has now reached the actual asymptotic decision point rather than merely naming it. The previous note showed that the nonlinear remainder is time-integrable. The present note uses that fact in the only place where it can finally decide the branch: the Duhamel terms for the low-frequency trace–longitudinal wave pair, the high-frequency Einstein wave block, and the quartic scalar Schrödinger channel.

The verdict is favorable and narrower than a redesign. In the same fixed completion $S_{\min,4}$, in the same unit-lapse spatial-harmonic gauge, and in the same weighted/homogeneous class, the nonlinear decay bootstrap closes, the weighted/homogeneous energy stays globally small, and the three propagating sectors admit limiting asymptotic profiles. No first genuine asymptotic obstruction appears there.

The asymptotic description is not best stated as literal unmodified free scattering in fixed coordinates. Because the unit-lapse shift transport persists at the wave rate, the correct profile is mildly modified by the transport-dressed linear evolution. But that is a same-class asymptotic

profile, not a failure. At current disciplined scope, the right conclusion is exactly the one the previous notes pointed toward: not a redesign yet. The same-class asymptotic/nonlinear-stability closure manuscript is the correct next step, and it closes positively.