The Æther-flow Interpretation of Relativity

Flow Geometry

Abstract

This paper gives the next positive extension of the active Æther-flow Interpretation of Relativity sequence by formalizing Æther-flow through congruence-based relativistic geometry. The scope is explicit. The manuscript does not alter the effective law of the theory: Æther-flow Interpretation of Relativity still adopts Einsteinian gravity with universal matter coupling and a single operational metric. Its purpose is interpretive deepening inside exact closure.

The central move is to represent Æther-flow by a future-directed unit timelike congruence u^μ on the exact relativistic spacetime and to read its kinematics through the standard decomposition of $\nabla_\mu u_\nu$ into expansion, shear, vorticity, and acceleration. This yields a disciplined flow dictionary. Expansion describes local volume change and cosmological growth of the observer congruence. Acceleration governs static gravitational support and redshift structure. Vorticity captures swirl, rotation, and frame dragging. Shear captures anisotropic distortion, tidal structure, and the strain content associated with gravitational waves. Local special relativity is recovered as the local inertial limit of the same flow geometry.

The resulting claim is precise. In Æther-flow Interpretation of Relativity, Æther-flow is not a second low-energy gravitational field competing with the metric. It is the congruence-based physical interpretation of how the exact relativistic geometry may be read by embedded observers. This sharpened interpretation makes the ontology more valuable without denying SR or GR and without pretending that a first-principles substrate derivation has already been completed.

1. Introduction

The active Æther-flow Interpretation of Relativity sequence already fixes three points. First, the Æther is the underlying four-dimensional substrate of reality and the Æther-flow is its intrinsic ordered motion. Second, the exact effective gravitational law of the theory is general relativity with universal matter coupling. Third, the effective theory is mathematically healthy and recovers the observed relativistic structure exactly. What remains is to make the flow language more rigorous so that it strengthens the interpretation of the theory rather than weakening it through loose metaphor.

That is the task of the present manuscript. The goal is not to replace the metric by a second low-energy object. Nor is the goal to claim a completed microphysical derivation of gravity from substrate variables. The goal is to identify the correct relativistic location of the Æther-flow idea inside exact closure.

Framework and claim boundary. Æther / Æther-flow framework denotes the broader Æther / Æther-flow framework built on the claims that the Æther is the underlying four-dimensional substrate of reality, the Æther-flow is its intrinsic ordered motion, observed three-dimensional space is the local experiential slice of that deeper substrate, S-time is the experienced order of change arising

from matter, light, and the Æther-flow, and observed expansion is the three-dimensional appearance of deeper four-dimensional ordered motion. Within that framework, Æther-flow Interpretation of Relativity denotes the exact-closure theory in which the effective gravitational dynamics are adopted to be exactly Einsteinian with universal matter coupling. In the active sequence, *adoption* means use of that established relativistic dynamics without claiming substrate derivation, while *derivation* is reserved for a first-principles recovery from explicit substrate variables.

The key discipline is this. A viable flow interpretation of GR cannot identify all gravitational phenomena with one naive image such as a literal vortex. Static spherical gravity exists without vorticity. Cosmological expansion exists without local swirl. Frame dragging does involve rotational structure, but it is not the whole content of gravitation. The correct relativistic move is therefore to interpret Æther-flow through the full kinematics of a timelike congruence rather than through one over-literal metaphor.

2. Flow Geometry in Æther-flow Interpretation of Relativity

Definition 1 (Flow geometry in Æther-flow Interpretation of Relativity)**.** *Flow geometry in Æther-flow Interpretation of Relativity is the pair $(g_{\mu\nu}, u^\mu)$, where $g_{\mu\nu}$ is the exact GR metric adopted by the theory and u^μ is a future-directed unit timelike congruence satisfying*

$$u^\mu u_\mu = -c^2. \tag{1}$$

The congruence provides the observer-based kinematic dictionary by which the Æther-flow ontology is read inside the exact relativistic geometry.

Remark 1. *The congruence is not a second field equation added to GR. It is an admissible observer field defined on the exact metric spacetime. Different physical situations may call for different natural congruences, but they all live inside the same metric geometry.*

This point fixes the scientific status of the manuscript. The paper is not proposing a bimetric theory, a preferred-frame matter coupling, or an extra propagating vector sector. It is specifying how the Æther-flow language may be made mathematically disciplined while the effective law remains exactly Einsteinian.

Proposition 1 (Single-law interpretation)**.** *In Æther-flow Interpretation of Relativity, the entire flow-geometry interpretation is subordinate to one effective law,*

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}, \tag{2}$$

with universal matter coupling to $g_{\mu\nu}$. The congruence u^μ adds interpretation, not a second low-energy dynamics.

3. Observer Rest Space and Local Decomposition

Given a unit timelike congruence u^μ, define the observer rest-space projector by

$$h_{\mu\nu} = g_{\mu\nu} + \frac{1}{c^2} u_\mu u_\nu. \tag{3}$$

Then

$$h_{\mu\nu}u^\nu = 0, \qquad h^\mu{}_\alpha h^\alpha{}_\nu = h^\mu{}_\nu, \tag{4}$$

so $h_{\mu\nu}$ defines the spatial geometry measured by observers whose four-velocity is u^μ.

Any vector X^μ decomposes uniquely into temporal and spatial parts relative to the congruence:

$$X^\mu = -\frac{u_\nu X^\nu}{c^2}u^\mu + h^\mu{}_\nu X^\nu. \tag{5}$$

Accordingly, any timelike worldline tangent w^μ may be written as

$$w^\mu = \Gamma\left(u^\mu + v^\mu\right), \qquad u_\mu v^\mu = 0, \tag{6}$$

where v^μ is the observer-measured spatial velocity and

$$\Gamma = -\frac{u_\mu w^\mu}{c^2} = \frac{1}{\sqrt{1 - v^2/c^2}}, \qquad v^2 = h_{\mu\nu}v^\mu v^\nu. \tag{7}$$

Proposition 2 (Local relativistic split)**.** *The pair $(u^\mu, h_{\mu\nu})$ gives the exact observer-based decomposition of relativistic physics in Æther-flow Interpretation of Relativity: u^μ fixes the local time direction, $h_{\mu\nu}$ fixes the local rest space, and* (7) *yields the ordinary Lorentz factor of relative motion.*

This is already enough to see why the Æther-flow language need not threaten SR. The congruence identifies a local observer family, while the local metric split still gives the ordinary relativistic relation between measured speed and Lorentz factor.

4. Kinematic Decomposition of the Æther-flow

Define the spatially projected derivative

$$\mathcal{D}_\mu \equiv h_\mu{}^\nu \nabla_\nu. \tag{8}$$

The fundamental kinematic quantities of the congruence are then:

$$a_\mu \equiv u^\nu \nabla_\nu u_\mu, \tag{9}$$

$$\theta \equiv \nabla_\mu u^\mu, \tag{10}$$

$$\sigma_{\mu\nu} \equiv h_{(\mu}{}^\alpha h_{\nu)}{}^\beta \nabla_\alpha u_\beta - \frac{1}{3}\theta h_{\mu\nu}, \tag{11}$$

$$\omega_{\mu\nu} \equiv h_{[\mu}{}^\alpha h_{\nu]}{}^\beta \nabla_\alpha u_\beta. \tag{12}$$

These satisfy

$$\sigma_{\mu\nu}u^\nu = 0, \qquad \omega_{\mu\nu}u^\nu = 0, \qquad \sigma^\mu{}_\mu = 0. \tag{13}$$

With these definitions, the covariant derivative of the congruence decomposes as

$$\nabla_\mu u_\nu = \frac{1}{3}\theta h_{\mu\nu} + \sigma_{\mu\nu} + \omega_{\mu\nu} - \frac{1}{c^2}u_\mu a_\nu. \tag{14}$$

Proposition 3 (Flow-sector map)**.** *In the exact Æther-flow Interpretation of Relativity line, the congruence kinematics provide the disciplined flow interpretation:*

1. *θ describes local volume expansion or contraction of the observer congruence.*
2. *a_μ describes the proper acceleration structure of non-geodesic observer families and governs static redshift support.*
3. *$\omega_{\mu\nu}$ describes swirl, twist, and frame-dragging structure.*
4. *$\sigma_{\mu\nu}$ describes anisotropic distortion, tidal deformation, and wave-induced strain.*

This proposition is the key conceptual payoff of the paper. It shows how the flow language becomes mathematically useful once it is distributed across the full congruence kinematics rather than collapsed into one slogan.

4.1. Raychaudhuri Control

The evolution of the expansion is governed by the Raychaudhuri equation

$$u^\mu \nabla_\mu \theta = -\frac{1}{3}\theta^2 - \sigma_{\mu\nu}\sigma^{\mu\nu} + \omega_{\mu\nu}\omega^{\mu\nu} - \nabla_\mu a^\mu - \frac{1}{c^2} R_{\mu\nu} u^\mu u^\nu. \tag{15}$$

This formula is important because it prevents the flow interpretation from becoming vague. Focusing, defocusing, local support against free fall, and the effect of matter on congruence evolution are all carried by definite geometric invariants.

4.2. Frobenius Criterion

The vorticity sector is equally controlled. The congruence is hypersurface orthogonal if and only if

$$\omega_{\mu\nu} = 0. \tag{16}$$

Thus swirl is not a metaphorical add-on. It has a sharp geometric meaning: failure of the observer flow lines to be orthogonal to a family of spatial hypersurfaces.

5. Expansion as Ordered Volume Change

The ontology of observed expansion is sharpened immediately by the expansion scalar θ. For any infinitesimal rest-space volume element V carried by the congruence,

$$\frac{1}{V}\frac{dV}{d\tau} = \theta. \tag{17}$$

This is the exact relativistic meaning of local ordered growth or contraction of the observer flow.

In a homogeneous and isotropic cosmological geometry with comoving congruence u^μ, one has

$$a_\mu = 0, \qquad \sigma_{\mu\nu} = 0, \qquad \omega_{\mu\nu} = 0, \qquad \theta = 3H, \tag{18}$$

where H is the Hubble parameter. The observed expansion of cosmology is therefore represented, in the congruence language, by nonzero θ rather than by motion into an external container.

Remark 2. *This is exactly the kind of clarification the Æther / Æther-flow ontology needs. The deeper ordered motion is not a literal river flowing into empty space. Its observer-accessible expansion content is encoded by congruence expansion inside the exact relativistic geometry.*

6. Static Gravity and Redshift as Acceleration Structure

The next step is equally important for physical interpretation. Static gravity is not fundamentally a vorticity effect. It is encoded, for the natural static observer congruence, by acceleration structure together with tidal curvature.

Consider a static line element

$$ds^2 = -N^2(\mathbf{x})\, c^2 dt^2 + \gamma_{ij}(\mathbf{x})\, dx^i dx^j, \tag{19}$$

and choose the static observer congruence

$$u^\mu = N^{-1} \delta^\mu{}_t. \tag{20}$$

For this congruence,

$$\theta = 0, \qquad \sigma_{\mu\nu} = 0, \qquad \omega_{\mu\nu} = 0, \tag{21}$$

while the acceleration is

$$a_\mu = \mathcal{D}_\mu \ln N. \tag{22}$$

Thus the static observers are generally not geodesic even though the spacetime may be stationary and non-rotating. The proper acceleration needed to hold station in the gravitational field is encoded directly in the gradient of the lapse.

In the weak-field limit,

$$N(\mathbf{x}) = 1 + \frac{\Phi(\mathbf{x})}{c^2} + \mathcal{O}(c^{-4}), \tag{23}$$

so

$$a_i = \frac{1}{c^2} \partial_i \Phi + \mathcal{O}(c^{-4}), \tag{24}$$

or equivalently

$$c^2 a_i = \partial_i \Phi + \mathcal{O}(c^{-2}). \tag{25}$$

This is the relativistic statement that the proper support acceleration of the static observer family reproduces the Newtonian gravitational potential gradient.

The same lapse factor controls gravitational redshift. For stationary observers with $dx^i = 0$,

$$d\tau = N(\mathbf{x})\, dt, \tag{26}$$

so for light emitted at A and received at B,

$$\frac{\nu_B}{\nu_A} = \frac{N(\mathbf{x}_A)}{N(\mathbf{x}_B)}. \tag{27}$$

In the weak-field regime this becomes

$$\frac{\nu_B - \nu_A}{\nu_A} = \frac{\Phi_A - \Phi_B}{c^2} + \mathcal{O}(c^{-4}). \tag{28}$$

Proposition 4 (Static-gravity interpretation)**.** *For the natural static observer congruence, the flow-geometry interpretation of gravity is carried primarily by congruence acceleration a_μ and curvature-induced tidal structure, not by vorticity.*

This proposition is essential. It shows why the language of a universal vortex cannot serve as the whole theory. Static gravitational fields are physically real even when $\omega_{\mu\nu} = 0$.

7. Swirl, Rotation, and Frame Dragging

The vortex intuition does have a legitimate relativistic home, but only in the rotational sector. When $\omega_{\mu\nu} \neq 0$, the congruence has local twist or swirl. This is the sector in which rotational gravity and frame dragging belong.

In a stationary but non-static geometry with nontrivial time-space metric components g_{0i}, the congruence of observers held at fixed spatial coordinates generally has nonzero vorticity. In the weak-field regime, $\omega_{\mu\nu}$ is controlled by the antisymmetric spatial derivatives of those time-space components. Thus the rotational part of the metric carries a genuine swirl content in the congruence description.

Proposition 5 (Rotational sector)**.** *Within the Æther-flow interpretation of Æther-flow Interpretation of Relativity, vorticity $\omega_{\mu\nu}$ is the correct mathematical carrier of swirl, rotational dragging, and local failure of hypersurface orthogonality.*

This is the disciplined way to keep the user's physical intuition without distorting the mathematics. Swirl is real, but it is not the whole of gravity. It is the rotational sector of the flow geometry.

The same point may be stated operationally. In the presence of frame dragging, gyroscope axes, local inertial directions, and orbit precession respond to the rotational structure of spacetime. In the congruence picture, that response is read through the vorticity sector rather than through the static acceleration sector.

8. Shear, Tidal Gravity, and Wave Strain

The tracefree symmetric part of the flow gradient is the shear tensor $\sigma_{\mu\nu}$. It governs anisotropic distortion of an observer bundle and therefore gives the correct home for tidal stretching and squeezing.

The curvature content measured by the observer field is encoded in the electric part of the Weyl tensor,

$$E_{\mu\nu} \equiv \frac{1}{c^2} C_{\mu\alpha\nu\beta} u^\alpha u^\beta, \tag{29}$$

which is spatial, symmetric, and tracefree in vacuum. For nearby freely falling observers separated by a spatial vector ξ^μ, the vacuum geodesic-deviation law reduces to

$$\frac{D^2 \xi^\mu}{D\tau^2} = -E^\mu{}_\nu \xi^\nu. \tag{30}$$

Thus the tidal content of gravity is registered directly as differential stretching and compression in the observer rest space.

Proposition 6 (Tidal sector)**.** *In the Æther-flow interpretation of Æther-flow Interpretation of Relativity, shear $\sigma_{\mu\nu}$ and the tidal tensor $E_{\mu\nu}$ together provide the correct mathematical reading of anisotropic gravitational distortion.*

This is again important for conceptual clarity. What people loosely call the bending of space and time is not one undifferentiated effect. In the flow-geometry description, part of it appears as static acceleration, part as rotational vorticity, and part as tidal shear driven by curvature.

The gravitational-wave sector fits naturally into the same picture. In transverse-traceless form around local Minkowski space,

$$ds^2 = -c^2 dt^2 + \left(\delta_{ij} + h_{ij}^{\mathrm{TT}}\right) dx^i dx^j, \qquad \partial_i h_{ij}^{\mathrm{TT}} = 0, \qquad h_{ii}^{\mathrm{TT}} = 0. \tag{31}$$

The wave does not create a second low-energy propagation law. It is already part of the exact GR metric sector. For freely falling detector congruences, it appears as oscillatory tidal deformation and therefore as time-dependent shear and geodesic deviation in the observer rest space.

9. Local Special Relativity as the Local Inertial Limit

The local relation to SR can now be stated sharply. At every event p, one may choose local inertial coordinates such that

$$g_{\mu\nu}(p) = \eta_{\mu\nu}, \qquad \partial_\alpha g_{\mu\nu}(p) = 0. \tag{32}$$

If the observer congruence is chosen so that

$$u^\mu(p) = (c, 0, 0, 0) \tag{33}$$

in those coordinates, then the rest-space projector reduces at p to the ordinary Euclidean spatial projector and the relative-motion decomposition (6) becomes the usual SR split.

Corollary 1 (Local SR recovery)**.** *For any event p and any unit timelike observer $u^\mu(p)$, the flow geometry of Æther-flow Interpretation of Relativity reduces locally to standard special-relativistic kinematics with Lorentz factor*

$$\Gamma = \frac{1}{\sqrt{1 - v^2/c^2}}. \tag{34}$$

This is the precise sense in which the Æther / Æther-flow interpretation does not deny SR. The theory does not add a measurable preferred-frame correction to local inertial physics in the exact-closure theory. Instead, it provides a deeper reading of why embedded observers always recover local Minkowski order inside the exact relativistic geometry.

10. What This Manuscript Fixes

The flow-geometry manuscript fixes the following positive structure for the active Æther-flow Interpretation of Relativity line.

1. It identifies Æther-flow with a congruence-based interpretation of the exact relativistic geometry rather than with a second low-energy gravitational field.
2. It provides the exact observer rest-space split $(u^\mu, h_{\mu\nu})$ appropriate to that interpretation.
3. It assigns distinct physical roles to expansion, acceleration, vorticity, and shear.
4. It shows that static gravity, redshift, swirl, frame dragging, tidal distortion, gravitational-wave strain, and local SR can all be read within one geometric dictionary.

5. It clarifies that vortex language is at most a statement about the vorticity sector and cannot be the whole account of gravity.

The manuscript does not change the claim boundary already fixed elsewhere in the active sequence. Einsteinian gravity is still adopted exactly in Æther-flow Interpretation of Relativity. A first-principles substrate derivation is still open. The present contribution is therefore interpretive sharpening through formal geometry, not a new low-energy dynamics.

11. Conclusion

The clean way to make Æther and Æther-flow more valuable inside Æther-flow Interpretation of Relativity is not to oppose GR or SR. It is to read the exact relativistic geometry more deeply. The congruence-based formalism provides exactly that move.

In this language, the Æther-flow is not one crude image but a structured relativistic object. Expansion θ captures ordered growth and cosmological volume change. Acceleration a_μ captures static gravitational support and redshift structure. Vorticity $\omega_{\mu\nu}$ captures swirl, rotation, and frame dragging. Shear $\sigma_{\mu\nu}$ captures anisotropic tidal distortion and wave strain. Local special relativity appears as the local inertial limit of the same geometry. The result is a stronger scientific interpretation of the ontology because it says exactly where each piece of the flow idea lives inside the accepted relativistic formalism.

That is the appropriate status of the manuscript. It does not claim that GR has already been derived from substrate microphysics. It claims that, once exact closure is adopted, the Æther / Æther-flow ontology can be connected to the measured relativistic world through a disciplined and comprehensive flow geometry.