

CMSC6950 Report: MAGICC, a Python wrapper for the simple climate model

Omid Tarkhaneh

June 2021

Abstract

This report demonstrates three computational tasks using a python wrapper for the simple climate model called MAGICC. In this regard, we employed some defined scenarios in MAGICC to perform computational tasks in python. The achieved results are visualized, and the results are analyzed based on the plots resulted from the computational tasks.

1 Introduction

Pymagicc is a Python interface for MAGICC, a reduced-complexity climatic carbon cycle model written in Fortran [2]. Pymagicc is a wrapper for the MAGICC binary², which works on Windows and is licensed under a Creative Commons Attribution NonCommercial-ShareAlike 3.0 Unported License³ [1]. MAGICC (Model for the Assessment of Greenhouse Gas Induced Climate Change) is commonly used in climate policy evaluations to predict future emissions trajectories, such as in the Intergovernmental Panel on Climate Change's Fifth Assessment Report (IPCC 2014) [1]. MAGICC is used in many Integrated Assessment Models (IAMs) to model the physical components of climate change. It's also been used to simulate the Coupled Model Intercomparison Projects' complicated atmosphere-ocean general circulation models (AOGCM).

This report uses some scenarios defined in the MAGICC called Representative Concentration Pathway (RCP). The IPCC's Representative Concentration Pathway (RCP) is a greenhouse gas concentration (rather than emis-

sions) trajectory. For the IPCC’s fifth Assessment Report (AR5) in 2014, four paths were employed for climate modeling and research [3]. The many climate futures described in the pathways are all considered feasible depending on the amount of greenhouse gases (GHG) emitted in the coming years. The RCPs are named after a possible range of radiative forcing levels in the year 2100 (formerly RCP2.6, RCP4.5, RCP6, and RCP8.5). In what follows, we first define the computational tasks and then discuss the task results and analyze the visualized results. Finally, the conclusion will be drawn.

2 Predicting the surface temperature

Different employed scenarios in MAGICC have some features and samples. There are various features in the dataset like region, scenario, variable, and so forth.

We have collected the results of all scenarios (RCP26, RCP45, RCP60, and RCP85) in a CSV file for this task. Before collecting this data into a CSV file, different computational tasks have been performed, such as dropping duplicate values, Nan values, etc. In addition, one feature of dataset is related to the region, and for predicting the surface temperature, we have limited the dataset to only count the world region despite some other regions in the dataset. Moreover, we have filtered the variable to surface temperature. In the dataset, variable attribute contains different values related to the greenhouse gases, surface temperature, emissions, etc. Given the achieved dataset after performing some computational task on it, we tried to visualize the obtained data utilized to predict mean surface temperature. In figure 1, we have depicted results of the data for all the employed scenarios for surface temperature in the world.




Figure 1: Mean surface temperature based on employed scenarios.

Figure 1 depicted the surface temperature from 1760 to 2100. The depicted figure shows that there is an increasing trend in all the scenarios, although there are some fluctuations in some years. Since 2000, we have a much more increasing trend where the surface temperature experienced the highest value in scenario RCP85.

3 Predicting the atmospheric concentration (CO₂)

This section evaluates the atmospheric concentration of CO₂ and utilized different scenarios of RCP26, RCP45, RCP60, and RCP85. To do so, first, we have performed some computational tasks using a "run" method in the pymagicc library in python. The mentioned method does some computation such as parsing the raw dataset. Then, we will filter the dataset based on region and variable attributes where we have used "world" for the region and "atmospheric concentration—CO₂" as the variable in the dataset. After

creating our dataset, we have visualized the data to evaluate the amount of atmospheric concentration between 1760 to 2100 in the world. Figure 2 depicted the obtained data and shows the atmospheric concentration in different years.




Figure 2: Atmospheric concentration based on employed scenarios.

Figure 2 shows the atmospheric concentration of CO₂ from 1760 to 2100 in the world. According to the figure, from 1760 to 1960, there is not a considerable increment in terms of atmospheric concentration of CO₂; however, since 2000, especially the world experience a huge concentration where its highest value is more dominant in scenario RCP85. To the best of our knowledge, this sharp increment is due to the industrial improvement which is created by mankind. After the industrial revolution, generating greenhouse gases increased, and this is one main reason for increasing the surface temperature too.

4 Predicting Emissions of CO2, Fossil, and Industrial

In this section, we worked on a raw dataset where we have not utilized the run method in pymagic library in python; rather, we have tried to handle the data and manipulate it ourselves. The preprocessing that is done include dropping Nan values and dropping duplicate samples, filtering dataset based on region and variable, dropping some of the features which are not needed to make the projection, and so on. After doing preprocessing, the dataset was visualized, and it shows an interesting diagram of emission of fossil and industrial as well as CO2 in the world. This time we tried to depict the data from 2000 to 2100. Figure 3 shows the results.




Figure 3: Emissions of CO2, Fossil, and Industrial based on employed scenarios.

In Figure 3 there are different trends for each scenarios. Scenario RCP85 shows an increase in the emission of the selected parameters(fossils and industrial, and CO2); however, scenarios such as RCP26 and RCP45 show a

decrease in the emission of the selected parameters. The interesting thing is that from 2050 to 2100, there is no such increase in emissions of the selected metrics, and even in some scenarios it this emission is decreased.

5 Conclusion

In this report, we have visualized some data based on some scenarios utilizing pymagic library in python. According to the results, it is showing that we will see an increasing trend in terms of atmospheric concentration and emission of greenhouse gases in the future. It is concluded that this increment is due to the industrial practice of humankind generating much more greenhouse gases and increasing the surface temperature. One exciting thing was that from 2050 to 2100, we might not experience a sharp increase, and we may even see a decrease base on some scenarios.

References

- [1] Gieseke, Robert, Sven N. Willner, and Matthias Mengel. "Pymagicc: A Python wrapper for the simple climate model MAGICC." *Journal of Open Source Software* 3, no. 22 (2018): 516.
- [2] Meinshausen, Malte, Sarah CB Raper, and Tom ML Wigley. "Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6–Part 1: Model description and calibration." *Atmospheric Chemistry and Physics* 11, no. 4 (2011): 1417-1456.
- [3] Moss, Richard H., Mustafa Babiker, Sander Brinkman, Eduardo Calvo, Tim Carter, James A. Edmonds, Ismail Elgizouli et al. "Towards new scenarios for analysis of emissions, climate change, impacts, and response strategies." (2008).