





Iron and Folic Acid

Pediatric IFA=20mg elemental Iron

Adult IFA tab= 60 mg elemental Iron

Duration: To be given for 14 days in anemic child and then reassess

Frequency: Once a day





Age of the child	Dose	Frequency
<4months	On doctors advise	
4 months-24 months (Wt 6-12 Kg)	1 tab 	Once a day* 
2 yrs-5yrs (12-19 kg)	2 tabs 	Once a day* 
*Can be increased on doctor's advise Side Effects: Constipation In case of diarrhea take doctors advise In case of pain abdomen tablet to be consumed after food		

Paracetamol

1 tablet=500mg

Duration: To be given for 3 days only

Frequency: Maximum four times a day at an interval of six hours





Age	Dose of tablet	Frequency
2 months -3 years (Wt 4-14 kgs)	$\frac{1}{4}$ tab (One fourth) 	Max 4 times a day 
3yrs -5yrs (Wt 14-19 kgs)	$\frac{1}{2}$ tab (Half) 	Max 4 times a day 

5ml or (1 tsp) Syrup=125 mg/5ml (Each1ml contains 25 mg of paracetamol)

Per kg dose of paracetamol=10-15 mg/kg/dose

Duration: To be given for 3 days only

Frequency: Maximum four times a day at an interval of six hours

Age	Dose of Syrup		Frequency
	In ml	In teaspoon	
New Born<3kg	1ml	$\frac{1}{4}$ tsp (One fourth) 	Max 4 times a day 
>1 year (>3kg-8kg)	2.5ml	$\frac{1}{2}$ tsp (Half) 	
1-3 yrs (>8-14 kgs)	5 ml	1 tsp (One) 	
>3 yrs >14 kgs	7.5 ml	1 $\frac{1}{2}$ tsp (One and Half) 