

IL-13 induces translocation of NF- κ B in cultured human bronchial smooth muscle cells

Kumiko Goto, Yoshihiko Chiba *, Miwa Misawa

Department of Pharmacology, School of Pharmacy, Hoshi University, 2-4-41 Ebara, Shinagawa-ku, Tokyo 142-8501, Japan

ARTICLE INFO

Article history:

Received 22 October 2008

Received in revised form 1 December 2008

Accepted 29 December 2008

Keywords:

IL-13

NF- κ B

Bronchial asthma

Human bronchial smooth muscle cell

ABSTRACT

Background and purpose: Interleukin-13 (IL-13), a major Th2 cytokine, plays an important role in bronchial asthma, including mucus production, inflammation and airway hyperresponsiveness. Although IL-13 through its binding to IL-4 receptor α (IL-4R α /IL-13R α 1) uses the canonical signal transducer and activator of transcription 6 (STAT6)-signaling pathway to mediate these tissue responses, recent studies have demonstrated that other signaling pathways may also be involved in. In the present study, whether IL-13 induces an activation of nuclear factor (NF)- κ B, inflammatory transcription factor, was investigated in human bronchial smooth muscle cells (hBSMCs). **Methods:** Nuclear proteins were extracted from cultured hBSMCs treated with tumor necrosis factor (TNF)- α (10 ng/mL) or IL-13 (100 ng/mL), and assayed for activated NF- κ B and STAT6 by Western blotting. **Result:** Treatments with TNF- α and IL-13 induced a translocation of NF- κ B to nuclei in hBSMCs. In addition, coincubation with BMS-345541 (0.3 μ M), an inhibitor of NF- κ B (I κ B) kinase (IKK) inhibitor, markedly inhibited the translocation of NF- κ B. **Conclusion:** Our results suggest for the first time that IL-13 activates NF- κ B in hBSMCs.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

T helper 2 (Th2) cytokines such as interleukin-4 (IL-4), IL-5 and IL-13 are important mediators of bronchial asthma, a complex chronic disease of the airways, which is characterized by airway inflammation, airway remodeling and airway hyperresponsiveness (AHR). In murine models of the asthmatic response, IL-13 has been shown to be both necessary and sufficient for the generation of asthma-like deleterious tissue alterations [1–4]. The mechanisms of these responses have not been fully defined. However, because IL-13 induces a greater number of genes in airway smooth muscle as compared with other lung cell types, including smooth muscle myosin light chain (MLC), phospholipase A₂ and IL-13 receptor α 1 (IL-13R α 1), airway smooth muscle cells have the capacity to respond to direct stimulation by IL-13 [5].

IL-13 binds to a polymeric receptor that contains IL-4R α and IL-13R α 1 subunits. The majority of the studies of this receptor have focused on its ability to induce signal transducer and activator of transcription 6 (STAT6) tyrosine phosphorylation and subsequent signaling. In murine models, IL-13 has been shown to mediate many of its asthma-relevant responses through the activation of the STAT6 pathway in bronchial epithelial cells [6]. However, other signal transduction pathways also contribute to the pathogenesis of asthma-like Th2 responses including airway hyperresponsive-

ness and inflammation [7,8]. Extracellular signal-regulated kinase (ERK) 1/2 mitogen activated protein kinase (MAPK) activation is required for optimal IL-13 stimulation of specific chemokines, matrix metalloproteinases (MMPs), and protease inhibitors in the murine lung [9].

The nuclear transcription factor NF- κ B, typical of p50 and p65 heterodimer, regulates over 150 genes involved in immune and inflammatory responses. Current evidence suggests an essential role in NF- κ B overactivation for increased expression of many inflammatory genes and for airway inflammation in asthma [9,10]. One well established mechanism of NF- κ B suppression is the export of nuclear NF- κ B by inhibitor of NF- κ B (I κ B) α , thereby rapidly repressing NF- κ B activation [11]. In general, NF- κ B has been known to be activated through phosphorylation and degradation of I κ B α by tumor necrosis factor (TNF)- α . However, it is reported that NF- κ B is activated by not only TNF- α but also IL-1 β , UV-light and stress and has an important role in the immune response including inflammation [12–14].

We hypothesized that NF- κ B plays a critical role in the pathogenesis of IL-13-induced signal transduction. To test this hypothesis, we investigated whether IL-13 induces an activation of NF- κ B in human bronchial smooth muscle cells (hBSMCs). As the results, both TNF- α and IL-13 induced activations of NF- κ B and STAT6, respectively. Interestingly, IL-13 also induced an activation of NF- κ B, which was demonstrated by the NF- κ B translocation to nuclei. These phenomena were significantly inhibited by combination with BMS-345541, I κ B kinase (IKK) inhibitor. Therefore, our results

* Corresponding author. Fax: +81 3 5498 5787.

E-mail address: chiba@hoshi.ac.jp (Y. Chiba).

suggest that both the IL-13 signaling and TNF- α signaling regulate the activation of NF- κ B in hBSMCs.

2. Materials and methods

2.1. Chemicals

All biochemicals were of analytical grade and were purchased from commercial suppliers: rhTNF- α (Peprotech, Paris, France), rhIL-13 (Peprotech, Paris, France) and BMS-345541 (Sigma, MO, USA).

2.2. Cell culture

Human bronchial smooth muscle cells (hBSMCs, Cambrex, MD, USA) were maintained in SmBM (Cambrex) medium supplemented with 5% fetal bovine serum (FBS), 1 mg/mL hFGF-B, 0.5 mg/mL hEGF, 5 mg/mL insulin and gentamycin/amphotericin B. The cells plated at a density of 2.7×10^4 per one well of a 6-well dish in FBS. Cells were serum deprived for 24 h using serum-free medium, which consists of SmBM supplemented with gentamycin/amphotericin B. The culture technique was used in experiments involving stimulation with 10 ng/mL TNF- α , 100 ng/mL IL-13 or sterile phosphate buffered saline (PBS) as vehicle control for 30 min or 1 h. BMS-345541, IKK inhibitor, was coincubated with TNF- α or IL-13.

2.3. Nuclear protein extraction

Cells were washed in PBS and lysed in 15 mM KCl, 10 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (pH 7.6), 2 mM $MgCl_2$, 0.1 mM ethylenediaminetetraacetic acid (EDTA), 1 mM dithiothreitol (DTT), 0.1% (v/v) Nonidet P-40, 0.5 mM phen-

ylmethylsulfonyl fluoride (PMSF), 2.5 μ g/ml leupeptin, and 5 μ g/ml aprotinin for 10 min on ice. Nuclei were pelleted by centrifugation at 14,000g for 20 s at 4 °C. Proteins were extracted from nuclei by incubation at 4 °C with vigorous vortexing in buffer (420 mM NaCl, 20 mM HEPES (pH 7.9), 0.2 mM EDTA, 25% (v/v) glycerol, 1 mM DTT, 0.5 mM PMSF, 2.5 μ g/ml leupeptin, and 5 μ g/ml aprotinin). Nuclear debris was pelleted by centrifugation at 13,000g for 30 min at 4 °C, and the supernatant extract was collected and stored at -85 °C until use.

2.4. Western blot analyses

To determine the levels of p65 in nucleus of hBSMCs and the phosphorylated levels of STAT6 in hBSMCs, the samples (10 mg of total protein per lane) were subjected to 10 or 7.5% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), and the proteins were then electrophoretically transferred to a poly vinylidene fluoride (PVDF) membrane. After blocking with 3% skim milk or 3% gelatin, the PVDF membrane was incubated with polyclonal rabbit anti-NF- κ B p65 (1:1000 dilution; Biolegend, San Diego, CA, USA), polyclonal rabbit anti-STAT6 (1:1000 dilution; Santa Cruz Biotechnology, Inc., CA, USA), or polyclonal rabbit anti-phosphorylated STAT6 antibody (1:1000 dilution; Santa Cruz). Then the membrane was incubated with horseradish peroxidase-conjugated donkey anti-rabbit IgG (1:2500 dilution; Amersham Biosciences, Co., NJ, USA), detected by an enhanced chemiluminescent system (Amersham Biosciences) and analyzed by a densitometry system. Detection of house-keeping gene was also performed on the same membrane by using polyclonal rabbit anti-histone H1 (1:1000 dilution; Santa Cruz), or monoclonal mouse anti- β -actin (1:5000 dilution; Santa Cruz) to confirm the same amount of proteins loaded.

Phosphorylated STAT6

Total STAT6




Fig. 1. Phosphorylation of signal transducer and activator of transcription 6 (STAT6) induced by TNF- α (10 ng/mL, 1 h) or IL-13 (100 ng/mL, 1 h) in cultured human bronchial smooth muscle cells (hBSMCs). Total proteins of hBSMCs were assayed for phosphorylated STAT6 by immunoblotting. (Upper photo) Typical blots for phosphorylated and total STAT6. The phosphorylation levels of STAT6 are summarized in the lower panel. IL-13 induced phosphorylation of STAT6, but TNF- α did not. ($n = 4$, *** $p < 0.001$ vs. control by one-way ANOVA with *post-hoc* Bonferroni/Dunn).

2.5. Statistical analyses

All the data were expressed as the mean with S.E.M. Statistical significance of difference was determined by unpaired Student's *t*-test or two-way analysis of variance (ANOVA) with *post-hoc* Bonferroni/Dunn (StatView for Macintosh ver. 5.0, SAS Institute, Inc., NC). A value of $p < 0.05$ was considered as significant.

3. Result

To determine whether TNF- α and IL-13 are capable of activating STAT6 and NF- κ B in hBSMCs, the levels of STAT6 phosphorylation and nuclear translocation of p65 were assessed by immunoblottings.

As shown in Fig. 1, IL-13 caused distinct phosphorylation of STAT6 in the hBSMCs: the level of phosphorylated STAT6 in the IL-13 (100 ng/mL)-treated cells was significantly increased as compared with that in the vehicle-treated control cells ($p < 0.001$). On the other hand, TNF- α (10 ng/mL) had no effect on the level of STAT6 phosphorylation (Fig. 1).

Fig. 2 shows the effects of TNF- α and IL-13 on the level of nuclear p65 in hBSMCs. As widely reported in various types of cells, TNF- α (10 ng/mL) caused a significant increase in nuclear p65 level (Fig. 2; $p < 0.01$), i.e., translocation of p65 to nuclei, in hBSMCs. Interestingly, a significant increase in nuclear p65 was also observed in the cells treated with 100 ng/mL of IL-13 (Fig. 2; $p < 0.05$), suggesting that IL-13 is capable of activating not only STAT6 but also NF- κ B in hBSMCs.

Next, the effect of an I κ B kinase (IKK) inhibitor, BMS-345541, on the IL-13-induced translocation of p65 was investigated to determine the role of IKK in the IL-13-induced activation of NF- κ B. As shown in Fig. 3, the TNF- α -induced translocation of p65 was completely blocked by BMS-345541 (0.3 μ M; $p < 0.01$). This concentra-

tion of BMS-345541 also inhibited the IL-13-induced translocation of p65 completely, indicating that the p65 translocation induced by IL-13 is mediated by an activation of IKK in hBSMCs.

4. Discussion

IL-13 is a Th2 cytokine that has emerged as a critical regulator of inflammatory immune responses, with key roles in asthma and parasite immunity [15,16]. IL-13 can be detected in the bronchial tissue [17], nasal lavage fluid [18], and induced sputum [17] of asthmatics. Following segmental allergen challenge, bronchoalveolar lavage (BAL) fluid contains IL-13 mRNA [19] and IL-13 protein [20], indicating that the cytokine is generated in the lung in response to respiratory provocation. Transgene pulmonary overexpression of IL-13 in mice is associated with several key pathological features of airways inflammation and remodeling, also observed in patients with chronic severe asthma, including lymphocyte and eosinophil accumulation, mucus cell metaplasia, subepithelial fibrosis and AHR [1]. Several studies indicate that IL-13 may cause AHR independently of inflammation and it is plausible that the effect may result from a direct effect on airway smooth muscle (ASM). ASM cells express functional IL-13 receptors and IL-13 is also linked to an augmented ASM contractility in rabbit tracheal strips [21,22].

IL-13 first binds to the IL-13R α 1 chain on the surface of cells. IL-4R α is recruited upon IL-13 interaction with IL-13R α 1, to form the high affinity receptor complex [23]. Heterodimerization of cell surface IL-13R α 1 and IL-4R α receptor chains initiates IL-13 signaling via recruitment of janus kinase (JAK) 1, JAK2 and tyrosine kinase (Tyk) 2, resulting in the phosphorylation of STAT6, a critical step in IL-13 and IL-4 dependent signaling [24].

The present study demonstrated for the first time that not only TNF- α (10 ng/mL) but also IL-13 (100 ng/mL) induces translocation




Fig. 2. Translocation of p65 to nuclei induced by TNF- α (10 ng/mL, 30 min) or IL-13 (100 ng/mL, 30 min) in cultured human bronchial smooth muscle cells (hBSMCs). Nuclear proteins of hBSMCs were assayed for p65 by immunoblottings. (Upper photo) Typical blots for p65 and histone H1. The expression levels of p65 are summarized in the lower panel. TNF- α or IL-13 induced translocation of NF- κ B to nuclei. ($n = 4$, * $p < 0.05$, ** $p < 0.01$ vs control by one-way ANOVA with *post-hoc* Bonferroni/Dunn).




Fig. 3. Effects of BMS-345541, an I κ B kinase (IKK) inhibitor, on translocation of p65 to nuclei induced by TNF- α (10 ng/mL) or IL-13 (100 ng/mL) in cultured human bronchial smooth muscle cells (hBSMCs). Nuclear proteins of hBSMCs were assayed for p65 by immunoblottings. (Upper photo) Typical blots for p65 and histone H1. The expression levels of p65 are summarized in the lower panel. BMS-345541 inhibited TNF- α - or IL-13-induced translocation of p65 to nuclei. ($n = 4$, * $p < 0.05$, *** $p < 0.001$ vs. control, ++ $p < 0.01$ vs. TNF- α alone and # $p < 0.05$ vs. IL-13 alone by two-way ANOVA with *post-hoc* Bonferroni/Dunn).




Fig. 4. Putative mechanisms of activation of signal transducer and activator of transcription 6 (STAT6) and NF- κ B by IL-13 and TNF- α in hBSMCs. The current findings suggest that, in addition to the activation of STAT6 pathway, IL-13 also activate NF- κ B via an activation of I κ B kinase (IKK), although the signaling between IL-13 receptors and IKK is not yet clear now.

of NF- κ B to nuclei in hBSMCs. A complex composed of IKKs (IKK α , IKK β and IKK γ) is an essential component in NF- κ B-signaling mechanisms participating in I κ B α phosphorylation. Although IKK α and IKK β are catalytic subunits, IKK γ is a regulatory subunit. IKK β is essential for activation of the classical NF- κ B pathway through phosphorylation of serine residues 32 and 36 of I κ B α , resulting in the ubiquitination and subsequent degradation of I κ B α by the 26S proteasome. The degradation of I κ B α exposes a nuclear translocation sequence facilitating translocation of NF- κ B to the nucleus. In spite of treatment with TNF- α or IL-13, the expression of total cellular p65 failed to show any change. Therefore, it is our judgment that upregulation of p65 in nuclei occurred by translocation of NF- κ B. Svetlana et al. [25] had demonstrated an activation of NF- κ B in mice when IL-13 was overexpressed in lung epithelia. In addition, IL-13-induced eosinophilia, inflammation, fibrosis, airway obstruction and alveolar remodeling were inhibited by interference with NF- κ B activation [25]. In our works, IL-13 also activated NF- κ B, and IL-13-induced translocation of NF- κ B to nuclei was completely inhibited by BMS-345541 (0.3 μ M), IKK inhibitor in hBSMCs. Because the TNF- α induced activation of NF- κ B is dependent on IKK β , we suggested that BMS-345541 (0.3 μ M) was perfectly inhibited IKK activity. Therefore, our result indicated that IL-13 activates NF- κ B via IKK β .

In conclusion, we suggest for the first time that IL-13 activates NF- κ B via an activation of IKK in hBSMCs (Fig. 4). Therefore, because the inhibition of NF- κ B activity suppressed not only TNF- α induced signaling but also IL-13 induced signaling partly, NF- κ B inhibitor can be a therapeutic target for bronchial asthma.

References

- [1] Zhu Z, Homer RJ, Wang Z, Chen Q, Geba GP, Wang J, et al. Pulmonary expression of interleukin-13 causes inflammation, mucus hypersecretion, subepithelial fibrosis, physiologic abnormalities, and eotaxin production. *J Clin Invest* 1999;103:779–88.
- [2] Zhu Z, Ma B, Homer RJ, Zheng T, Elias JA. Use of the tetracycline-controlled transcriptional silencer (tTS) to eliminate transgene leak in inducible overexpression transgenic mice. *J Biol Chem* 2001;276:25222–9.
- [3] Elias JA, Lee TCG, Homer RJ, Chen Q, Ma B, Blackburn M, et al. Transgenic modeling of interleukin-13 in the lung. *Chest* 2003;123:339S–45S.
- [4] Herrick CA, Bottomly K. To respond or not to respond: T cells in allergic asthma. *Nat Rev Immunol* 2003;3:405–12.
- [5] Woodruff Prescott G. Gene expression in asthmatic airway smooth muscle. *Proc Am Thorac Soc* 2008;5:113–8.
- [6] Kuperman DA, Huang X, Koth LL, Chang GH, Dolganov GM, Zhu Z, et al. Direct effects of interleukin-13 on epithelial cells cause airway hyperreactivity and mucus overproduction in asthma. *Nat Med* 2002;8:885–9.
- [7] Lee JH, Kaminski N, Dolganov G, Grunig G, Koth L, Solomon C, et al. Interleukin-13 induces dramatically different transcriptional programs in three human airway cell types. *Am J Respir Cell Mol Biol* 2001;5:74–85.
- [8] Lee PJ, Zhang X, Shan P, Ma B, Lee CG, Homer RJ, et al. ERK1/2 mitogen-activated protein kinase selectively mediates IL-13-induced lung inflammation and remodeling in vivo. *J Clin Invest* 2006;116:163–73.
- [9] Christman JW, Sadikot RT, Blackwell TS. The role of nuclear factor κ B in pulmonary diseases. *Chest* 2000;117:1482–7.
- [10] Zhou LF, Zhu Y, Cui XF, Xie WP, Hu AH, Yin KS. Arsenic trioxide, a potent inhibitor of NF- κ B, abrogates allergen-induced airway hyperresponsiveness and inflammation. *Respir Res* 2006;7:146.
- [11] Tanaka T, Grusby MJ, Kaisho T. PDLIM2-mediated termination of transcription factor NF- κ B activation by intranuclear sequestration and degradation of the p65 subunit. *Nat Immunol* 2007;8:584–91.
- [12] Hayden MS, Ghosh S. Signaling to NF- κ B. *Genes Dev* 2004;18:2195–224.
- [13] Janssens S, Tschopp J. Signals from within: the DNA-damage-induced NF- κ B response. *Cell Death Differ* 2006;13:773–84.
- [14] Ea CK, Deng L, Xia ZP, Pineda G, Chen ZJ. Activation of IKK by TNF α requires site-specific ubiquitination of RIP1 and polyubiquitin binding by NEMO. *Mol Cell* 2006;22:245–57.
- [15] Wynn TA. IL-13 effector functions. *Annu Rev Immunol* 2003;21:425–56.
- [16] Wills-Karp M. Interleukin-13 in asthma pathogenesis. *Immunol Rev* 2004;202:175–90.
- [17] Berry MA, Parker D, Neale N, Woodman L, Morgan A, Monk P, et al. Sputum and bronchial submucosal IL-13 expression in asthma and eosinophilic bronchitis. *J Allergy Clin Immunol* 2004;114:1106–9.
- [18] Noah TL, Tudor GE, Ivins SS, Murphy PC, Peden DB, Henderson FW. Repeated measurement of nasal lavage fluid chemokines in school-age children with asthma. *Ann Allergy Asthma Immunol* 2006;96:304–10.
- [19] Prieto J, Lensmar C, Roquet A, van der Ploeg I, Gigliotti D, Eklund A, et al. Increased interleukin-13 mRNA expression in bronchoalveolar lavage cells of atopic patients with mild asthma after repeated low-dose allergen provocations. *Respir Med* 2000;94:806–14.
- [20] Batra V, Musani AI, Hastie AT, Khurana S, Carpenter KA, Zangrilli JG, et al. Bronchoalveolar lavage fluid concentrations of transforming growth factor (TGF)- β 1, TGF- β 2, interleukin (IL)-4 and IL-13 after segmental allergen challenge and their effects on alpha-smooth muscle actin and collagen III synthesis by primary human lung fibroblasts. *Clin Exp Allergy* 2004;34:437–44.
- [21] Grunstein MM, Hakonarson H, Leiter J, Chen M, Whelan R, Grunstein JS, et al. IL-13-dependent autocrine signaling mediates altered responsiveness of IgE-sensitized airway smooth muscle. *Am J Physiol Lung Cell Mol Physiol* 2002;282:L520–8.
- [22] Laporte JC, Moore PE, Baraldo S, Jouvin MH, Church TL, Schwartzman IN, et al. Direct effects of interleukin-13 on signaling pathways for physiological responses in cultured human airway smooth muscle cells. *Am J Respir Crit Care Med* 2001;164:141–8.
- [23] Andrews AL, Holloway JW, Puddicombe SM, Holgate ST, Davies DE. Kinetic analysis of the interleukin-13 receptor complex. *J Biol Chem* 2002;277:46073–8.
- [24] Palmer-Crocker RL, Hughes CC, Pober JS. IL-4 and IL-13 activate the JAK2 tyrosine kinase and Stat6 in cultured human vascular endothelial cells through a common pathway that does not involve the gamma c chain. *J Clin Invest* 1996;98:604–9.
- [25] Chapoval SP, Al-Garawi A, Lora JM, Strickland I, Ma B, Lee PJ, et al. Inhibition of NF- κ B activation reduces the tissue effects of transgenic IL-13. *J Immunol* 2007;179:7030–41.