

Uniform Convergence of Bernstein Polynomial Sequence and its Derivatives

Florent ONANA ASSOUGA

Under the Supervision of
Prof. Walter Van Assche
KU Leuven, Belgium



AIMS

African Institute for
Mathematical Sciences
CAMEROON



KU Leuven

Department of Mathematics
Celestijnenlaan 200 B bus 2400
BE-3001 Leuven (Heverlee)

January 26, 2024



Table of contents


- 1 Motivation-Objective
- 2 Weierstrass'first Theorem and Uniform Convergence of Bernstein polynomial
 - Definition and Examples
 - Statement and Uniform convergence
- 3 Bernstein Polynomial and its Derivatives
 - Another representation of Bernstein polynomial
 - Derivatives of Bernstein Polynomials
 - Uniform Convergence of Derivatives
- 4 Applications of Bernstein Polynomial Approximation
 - shape-preservation and variation-diminution
 - Resluts
- 5 Conclusion



Motivation-Objective

Introduction

- We are interested on sequences of polynomials named after their creator **Sergei Natanovich Bernstein**.



He proved :

- The **Weierstrass approximation theorem**
- And the **Bernstein's theorem (approximation theory)** .

Figure: March 1880-October 1968



Motivation

- One might wonder why Bernstein created “new” polynomials for a constructive proof of Weierstrass’s Theorem , instead of using polynomials such as Taylor polynomials or interpolating polynomials that were already known to mathematics.



Motivation

- One might wonder why Bernstein created “new” polynomials for a constructive proof of Weierstrass’s Theorem , instead of using polynomials such as Taylor polynomials or interpolating polynomials that were already known to mathematics.
- For even setting aside questions of convergence, Taylor polynomials are applicable only to functions that are infinitely differentiable, and not to all continuous functions.



Objectives

- Consider how Bernstein discovered his polynomials, and extend it to a class of function continuous to any compact $[a, b]$



Objectives

- Consider how Bernstein discovered his polynomials, and extend it to a class of function continuous to any compact $[a, b]$
- Study the uniform convergence of Bernstein polynomial and its derivatives (based on additional assumptions)



Objectives

- Consider how Bernstein discovered his polynomials, and extend it to a class of function continuous to any compact $[a, b]$
- Study the uniform convergence of Bernstein polynomial and its derivatives (based on additional assumptions)
- See that although the convergence of the Bernstein polynomials is slow, they have compensating “shape-preserving” properties.



Definition (Bernstein polynomial)

Given a function f on $[0, 1]$, we define the Bernstein polynomial as

$$B_n(f, x) = \sum_{k=0}^n \binom{n}{k} f\left(\frac{k}{n}\right) x^k (1-x)^{n-k} \quad (1)$$

for each positive integer n .



Definition (Bernstein polynomial)

Given a function f on $[0, 1]$, we define the Bernstein polynomial as

$$B_n(f, x) = \sum_{k=0}^n \binom{n}{k} f\left(\frac{k}{n}\right) x^k (1-x)^{n-k} \quad (1)$$

for each positive integer n .

Example

- $B_n(1, x) = 1$, $B_n(x, x) = x$, $B_n(x^2, x) = x^2 + \frac{x(1-x)}{n}$
- $B_n(e^{\alpha x}, x) = \left(xe^{\frac{\alpha}{n}} + (1-x)\right)^n$



Theorem (Weierstrass's first Theorem)

If $f \in C([0, 1])$, then for every $\epsilon > 0$, there exists a polynomial p such that $\|f - p\|_\infty < \epsilon$.



Theorem (Weierstrass's first Theorem)

If $f \in C([0, 1])$, then for every $\epsilon > 0$, there exists a polynomial p such that $\|f - p\|_\infty < \epsilon$.

Proof.

It uses the Bernstein polynomial sequence and the Lemma of [Korovkin](#). □



Theorem (Weierstrass's first Theorem)

If $f \in C([0, 1])$, then for every $\epsilon > 0$, there exists a polynomial p such that $\|f - p\|_\infty < \epsilon$.

Proof.

It uses the Bernstein polynomial sequence and the Lemma of Korovkin. □

Lemma (Korovkin Lemma)

Suppose $H_n, n = 1, 2, 3, \dots$ is a sequence of positive linear operators on $C[a, b]$. If for $k = 1, 2, 3$ one has $\lim_{n \rightarrow \infty} \|H_n(x^k, x) - x^k\|_\infty = 0$, then it follows that for all $f \in C[a, b]$

$$\lim_{n \rightarrow \infty} \|H_n(f, x) - f(x)\|_\infty = 0. \quad (2)$$



Corollary

- If $f \in \mathcal{C}[0, 1]$, then

$$B_n(f, \cdot) \xrightarrow[0,1]{CU} f. \quad (3)$$

- If $f \in \mathcal{C}[a, b]$, we define the bijection

$$\begin{cases} h : [a, b] \rightarrow [0, 1] \\ t \mapsto h(t) = \frac{t-a}{b-a} \end{cases}, \text{ thus } f \circ h^{-1} \in \mathcal{C}[0, 1] \text{ and}$$

$$B_n(f \circ h^{-1}, h(\cdot)) \xrightarrow[a,b]{CU} f. \quad (4)$$

Another representation of Bernstein polynomial

Definition (Forward difference operator)

Let $f \in \mathcal{C}[0, 1]$ $x \in [0, 1]$ and $h > 0$ such that $x + h \in [0, 1]$.

- $\Delta_h f(x) = f(x + h) - f(x)$
- $\Delta_h^0 f(x) = f(x)$
- $\Delta_h^r f(x) = \Delta_h^{r-1}(\Delta_h f(x)) = \Delta_h^{r-1} f(x + h) - \Delta_h^{r-1} f(x) \quad \text{if} \quad r \geq 1$



Another representation of Bernstein polynomial

Definition (Forward difference operator)

Let $f \in \mathcal{C}[0, 1]$ $x \in [0, 1]$ and $h > 0$ such that $x + h \in [0, 1]$.

- $\Delta_h f(x) = f(x + h) - f(x)$
- $\Delta_h^0 f(x) = f(x)$
- $\Delta_h^r f(x) = \Delta_h^{r-1}(\Delta_h f(x)) = \Delta_h^{r-1} f(x + h) - \Delta_h^{r-1} f(x) \quad \text{if} \quad r \geq 1$

Proposition

The Bernstein polynomial may be expressed in the form

$$B_n(f, x) = \sum_{r=0}^n \binom{n}{r} \Delta_{\frac{1}{n}}^r f(0) x^r. \quad (5)$$



Derivatives of Bernstein Polynomials

- The derivative of the Bernstein polynomial $B_{n+1}(f, x)$ may be expressed in the form

$$B'_{n+1}(f, x) = (n+1) \sum_{r=0}^n \Delta_h f \left(\frac{r}{n+1} \right) \binom{n}{r} x^r (1-x)^{n-r} \quad (6)$$

for $n \geq 0$ and $h = \frac{1}{n+1}$.



Derivatives of Bernstein Polynomials

- The derivative of the Bernstein polynomial $B_{n+1}(f, x)$ may be expressed in the form

$$B'_{n+1}(f, x) = (n+1) \sum_{r=0}^n \Delta_h f \left(\frac{r}{n+1} \right) \binom{n}{r} x^r (1-x)^{n-r} \quad (6)$$

for $n \geq 0$ and $h = \frac{1}{n+1}$.

- For any integer $k \geq 0$, the k th derivative of $B_{n+k}(f, x)$ may be expressed in terms of the k th differences of f as

$$B_{n+k}^{(k)}(f, x) = \frac{(n+k)!}{n!} \sum_{r=0}^n \Delta_h^k f \left(\frac{r}{n+k} \right) \binom{n}{r} x^r (1-x)^{n-r} \quad (7)$$

for $n \geq 0$ and $h = \frac{1}{n+k}$.



Theorem

If $f \in \mathcal{C}^k[0, 1]$, for some integer $k \geq 0$ then $B_n^{(k)}(f, x)$ converges uniformly to $f^{(k)}$ on $[0, 1]$.



Theorem

If $f \in \mathcal{C}^k[0, 1]$, for some integer $k \geq 0$ then $B_n^{(k)}(f, x)$ converges uniformly to $f^{(k)}$ on $[0, 1]$.

Proof.

See [2]



- We now state results concerning the Bernstein polynomials for a convex function f .
- First we suppose that the definition convexity and its connection with second-order divided differences are known.



Theorem

① If f is convex on $[0, 1]$,

$$B_n(f, x) \geq f(x), \quad 0 \leq x \leq 1 \quad \forall n \geq 1 \quad (8)$$



Theorem

① If f is convex on $[0, 1]$,

$$B_n(f, x) \geq f(x), \quad 0 \leq x \leq 1 \quad \forall n \geq 1 \quad (8)$$

② If f is convex on $[0, 1]$,

$$B_{n-1}(f, x) \geq B_n(f, x), \quad 0 \leq x \leq 1 \quad \forall n \geq 2 \quad (9)$$



Shape-preservation

Theorem

① If f is convex on $[0, 1]$,

$$B_n(f, x) \geq f(x), \quad 0 \leq x \leq 1 \quad \forall n \geq 1 \quad (8)$$

② If f is convex on $[0, 1]$,

$$B_{n-1}(f, x) \geq B_n(f, x), \quad 0 \leq x \leq 1 \quad \forall n \geq 2 \quad (9)$$

remark

If $f \in \mathcal{C}[0, 1]$, the inequality in (9) is strict for $0 < x < 1$.



Results

We take the example of: $B_n(e^{\alpha x}, x) = (xe^{\frac{\alpha}{n}} + (1-x))^n$ for $\alpha = 2$.




Figure: Comparison of f and $B_n(f, \cdot)$

Results

We take the example of: $B_n(e^{\alpha x}, x) = (xe^{\frac{\alpha}{n}} + (1-x))^n$ for $\alpha = 2$.





Figure: Comparison of f and $B_n(f, \cdot)$

Figure: $B_{n-1}(f, \cdot)$ Versus $B_n(f, \cdot)$



Conclusion

- Bernstein polynomials also form the mathematical basis for **Bézier curves** , which later became important in computer graphics.



Conclusion

- Bernstein polynomials also form the mathematical basis for [Bézier curves](#) , which later became important in computer graphics.
- In the 1950s, [H. Bohman](#) and [P. P. Korovkin](#) obtained an amazing generalization of Bernstein's Theorem. They found that as far as convergence is concerned, the crucial properties of the Bernstein operator B_n are that $B_n f \rightarrow f$ uniformly on $[0, 1]$ for $f = 1, x$, and x^2 , and that B_n is a monotone linear operator.





Conclusion

- Bernstein polynomials also form the mathematical basis for [Bézier curves](#) , which later became important in computer graphics.
- In the 1950s, [H. Bohman](#) and [P. P. Korovkin](#) obtained an amazing generalization of Bernstein's Theorem. They found that as far as convergence is concerned, the crucial properties of the Bernstein operator B_n are that $B_n f \rightarrow f$ uniformly on $[0, 1]$ for $f = 1, x$, and x^2 , and that B_n is a monotone linear operator.
- Several generalizations have been proposed for Bernstein polynomials: [a generalization based on the q-integers](#)



References

-  George A. Anastassiou, Razvan A. Mezei, *Numerical Analysis Using Sage*, Springer-2015
-  Michael S. Floater, *On the convergence of derivatives of Bernstein approximation* , Journal of Approximation Theory 134 (2005) 130 – 135



**THANK YOU FOR YOUR KIND
ATTENTION**

