Chapter 14 Notes

Section 1

Notes

Let K and K' be classes, K be transitive, $K \subseteq K'$, and $\phi(x_1, ..., x_n)$ a formula whose constants are all in K. Note that, even if we assume $\phi(a_1, ..., a_n)$ is true over K iff it is true over K' (for all constants $a_1, ..., a_n$ in K), we *cannot* prove that $(\exists x)(\phi(x, x_1, ..., x_n))$ is such that $(\exists x)(\phi(x, a_1, ..., a_n))$ is true over K iff it is true over K' (for all constants $a_1, ..., a_n$ in K); call any ϕ *good* if this property does hold for it. For, let $a_1, ..., a_n$ be constants in K. On the one hand, suppose that $(\exists x)(\phi(x, a_1, ..., a_n))$ is true over K. Then for some $a \in K$, $\phi(a, a_1, ..., a_n)$ is true over K, and thus by assumption, $\phi(a, a_1, ..., a_n)$ is true over K', hence $(\exists x)(\phi(x, a_1, ..., a_n))$ is true over K'. However, on the other hand, suppose that $(\exists x)(\phi(x, a_1, ..., a_n))$ is true over K'. Then for some $a \in K'$, $\phi(a, a_1, ..., a_n)$ is true over K'. However, unlike in Exercise 1.1 (a), we *cannot* conclude that $a \in K$, meaning it does not necessarily follow that $\phi(a, a_1, ..., a_n)$ is true over K (and thus it does not necessarily follow that $(\exists x)(\phi(x, a_1, ..., a_n))$ is true over K.

Using the same definitions as in the preceeding paragraph, note the following. Suppose $\phi(x_1, ..., x_n)$ is only such that if $\phi(a_1, ..., a_n)$ is true over K (for constants $a_1, ..., a_n$ in K), then $\phi(a_1, ..., a_n)$ is true over K'. It certainly does *not* follow that $\sim\phi$ is good (or even satisfies the weaker property if if $\sim\phi(a_1, ..., a_n)$ is true over K, then it is true over K'). For, if $\sim\phi(a_1, ..., a_n)$ is true over K, then $\phi(a_1, ..., a_n)$ is false over K. But because we do not have that ϕ is good, we cannot conclude that $\phi(a_1, ..., a_n)$ is false over K' (now, if $\sim\phi(a_1, ..., a_n)$ was true over K', then $\phi(a_1, ..., a_n)$ would be false over K', and then by assumption, $\phi(a_1, ..., a_n)$ would also need to be false over K, hence $\sim\phi(a_1, ..., a_n)$ would be true over K).

The above paragraph explains why Σ-formulas do not include negations (unless the subformlula the negation is the main operator of is itself a Σ_0 formula).

Proposition 1.1

Note that part (3) of this proposition follows in a similar way as part (3) of Exercise 1.1 (b) is proven below.

Exercise 1.1 (a)

Suppose K and K' are classes, such that K is transitive, $K \subseteq K'$, $a_1, ..., a_n$ are elements of K, and $\phi(x_1, ..., x_n)$ is a formula whose constants are all in K. Suppose also that $\phi(x_1, ..., x_n)$ is Σ_0. We are to show that $\phi(a_1, ..., a_n)$ is true over K iff it is true over K'. We prove this by induction on the degree of $\phi(x_1, ..., x_n)$.

Base Case: If $\phi(x_1, ..., x_n)$ is of the form $x \in y$, then $a_1 \in a_2$ is true over K iff it is true over K' (as a_1 and a_2 are both in K', as $K \subseteq K'$).

If the result holds for some ϕ, then obviously it also holds for $\sim\phi$.

If the result holds for ϕ and ψ, then obviously it also holds for $\phi \wedge \psi$.

Suppose the result holds for some $\phi(x, y, x_1, ..., x_n)$ (whose free variables are all in the set $\{x, y, x_1, ..., x_n\}$). We must show that the result also holds for

$$(\exists x \in y)(\phi(x, y, x_1, ..., x_n))$$

Take any constants $a, b_1, ..., b_n$ of K. We must show that $(\exists x \in a)(\phi(x, a, b_1, ..., b_n))$ is true over K iff it is true over K'. On the one hand, suppose that it is true over K. Then for some $b \in a$, $\phi(b, a, b_1, ..., b_n)$ is true over K. But the constants $b, a, b_1, ..., b_n$ are all in K, meaning by assumption, $\phi(b, a, b_1, ..., b_n)$ is true over K', and thus $(\exists x \in a)(\phi(x, a, b_1, ..., b_n))$ is true over K'. On the other hand, suppose that $(\exists x \in a)(\phi(x, a, b_1, ..., b_n))$ is true over K'. Then for some $b \in a$, $\phi(b, a, b_1, ..., b_n)$ is true over K'. But K is transitive, and $a \in K$, meaning $b \in K$. Thus the constants $b, a, b_1, ..., b_n$ are all in K, meaning, by assumption, $\phi(b, a, b_1, ..., b_n)$ is true over K, and thus $(\exists x \in a)(\phi(x, a, b_1, ..., b_n))$ is true over K.

Exercise 1.1 (b)

Suppose K and K' are classes, such that K is transitive, $K \subseteq K'$, $a_1, ..., a_n$ are elements of K, and $\phi(x_1, ..., x_n)$ is a formula whose constants are all in K. Suppose also that $\phi(x_1, ..., x_n)$ is Σ. We are to show that $\phi(a_1, ..., a_n)$ is true over K iff it is true over K'. Call a formula ϕ good if it satisfies this result. We prove this by induction on the degree of $\phi(x_1, ..., x_n)$. In fact, we prove:

(1) Every Σ_0 formula is good.

(2) If ϕ, ψ are good, then so are $\phi \wedge \psi$ and $\phi \vee \psi$.

(3) If ϕ is good, so is $(\exists x)\phi$.

(4) If ϕ is good, so are $(\forall x \in y)\phi$ and $(\exists x \in y)\phi$.

Proof. (1) This follows by Exercise 1.1 (a).

(2) Obvious.

(3) Suppose $\phi(x, x_1, ..., x_n)$ is good (where all free variables of ϕ are in the set $\{x, x_1, ..., x_n\}$). We must show that $(\exists x)\phi(x, x_1, ..., x_n)$ is good. So suppose $(\exists x)\phi(x, a_1, ..., a_n)$ was true over K. Then, for some $a \in K$, $\phi(a, a_1, ..., a_n)$ is true over K, hence by assumption, $\phi(a, a_1, ..., a_n)$ is true over K', meaning $(\exists x)\phi(x, a_1, ..., a_n)$ is true over K'.

(4) Suppose $\phi(x, y, z_1, ..., z_n)$ is good. We wish to show that $(\forall x \in y)\phi(x, y, z_1, ..., z_n)$ is also good. Well, let $a, b_1, ..., b_n$ be constants in K. Suppose $(\forall x \in a)\phi(x, a, b_1, ..., b_n)$ is true over K. Because K is transitive, it follows that $\phi(c, a, b_1, ..., b_n)$ is true over K for all elements $c \in a$. Then, by the assumption that ϕ is good, it follows that $(\forall x \in a)\phi(x, a, b_1, ..., b_n)$ is true over K'.

As for the formula $(\exists x \in y)\phi$, this is $(\exists x)(x \in y \wedge \phi)$, which is good by parts (1), (2), and (3) of this exercise. □

Section 2

Notes

It is remarked in the last paragraph of this section that once we show $y = \mathcal{F}(x)$ is Σ, it will follow by Theorem 2.8 that the relation $y = L_\alpha$ is also Σ, and absolute. It does follow that it is Σ, but it does not immediately follow by the theorem that it is absolute - only that it is *upwards* absolute.

Propositions 2.6 and 2.7

There are multiple problems with these two propositions. First, the propositions, as stated, are clearly incorrect, as the proofs given require f to be Σ, which is not assumed (because f here is being treated as a constant, not a variable). To fix this, for any set x, let us define $x''(y)$ for any set y to be the set of all $z \in \text{Ran}(x)$ such that $\langle w, z \rangle \in x$ for some $w \in y$. This more general definiton allows us to correct the proof.

In particular, we show that the relation $y = z''(x)$ (this time, z is a *variable*, meaning this is a legitimate equation, without any prior assumptions of being Σ) is Σ (in fact Σ_0).

Proof. $y = z''(x)$ is the conjunction of the following two Σ_0 conditions:

$$(1)\ (\forall v \in x)(\forall w \in z)(\langle v, w\rangle \in z \supset w \in y)$$
$$(2)\ (\forall v \in y)(\exists w \in x)(\langle w, v\rangle \in z)$$

□

If we were to restrict z to functions, then the proof would involve adding the condition that z is a function (that is, add in $\mathrm{Fun}(z)$, which is Σ_0 by results from Chapter 12).

Theorem 2.8

There is an issue with the proof of this theorem. In particular, $f(\beta + 1) = F(f \restriction \beta)$ is used in the supposed Σ formula, yet it was never established that $f(\beta + 1) = F(f \restriction \beta)$ is itself Σ. Here is why this is.

First, note that the relation $y = f \restriction x$ is Σ, as it is equivalent to the conjunction of the following two formulas:

$$(1)\ (\forall z \in y)(\exists w \in \mathrm{Dom}(f))(\exists v \in \mathrm{Ran}(f))(w \in x \wedge z = \langle w, v\rangle)$$
$$(2)\ (\forall z \in f)((\exists w \in x)(\exists v \in \mathrm{Ran}(f))(z = \langle w, v\rangle) \supset z \in y)$$

It is obvious that the above result means $f(\beta + 1) = F(f \restriction \beta)$ is Σ. To be specific, it is then represented by the following Σ formula:

$$(1)\ (\exists z \in \mathrm{Ran}(f))(\exists y)(\exists w)(\langle \beta + 1, z\rangle \in f \wedge w = f \restriction \beta \wedge y = F(w) \wedge y = z)$$

There is another error in the proof of this theorem. The occurences of $(\forall \beta \in \alpha + 1)$ should be replaced by $(\forall \beta \in \alpha)$ for the following reason: If we took the previous formula $((\forall \beta \in \alpha + 1))$, the formula in the proof would tell us that, for α, $f(\alpha + 1) = F(f \restriction \alpha)$, meaning $f(\alpha + 1)$ is defined. But that is not how we defined f, as we needed $\mathrm{Dom}(f) = \alpha + 1$, meaning $\alpha + 1$ would *not* be in the domain of f.

In fact, it seems that $F(f \restriction \beta)$ is just a typo, and really should be $F(f(b))$. This is due to the definition of F^*: $F^*(\alpha + 1)$ is not equal to $F(F^* \restriction \alpha)$, but is equal to $F(F^*(\alpha))$.