

## **VACCINE SCHEDULE (JANUARY 2017)**

At birth (in hospital) - Hepatitis B

2 months – Pentacel(DTaP, Hib,IPV), PCV 13(Prevnar), Rotateq, Hep B

4 months – Pentacel(DTaP, Hib, IPV), PCV 13, Rotated

6 months – Pentacel (DTap, Hib, IPV), PCV 13, Rotatedq (no IPV if doing Separate vaccines)

9 months – Hep B

12 months – MMR, Varicella, Hep A

15 months – Pentacel (DTaP, Hib, IPV), PCV 13

2 years – Hep A

4-5 years – Dtap, IPV, MMR/Varicella (Proquad)

10 years- Tdap (Adacel –if not previously given and >2 years from last tetanus

11 + years – MCV4 (Menactra), HPV(Gardasil)

16-21 years – Menactra booster if needed, Trumenba

Gardasil: Offer to all males and females >11 years of age. Adolescents >15 years of age. The 2<sup>nd</sup> and 3<sup>rd</sup> doses are given 2 and 6 months after the first dose. The minimum interval between 1<sup>st</sup> and 2<sup>nd</sup> doses is 1 month. The minimum interval between 2<sup>nd</sup> and 3<sup>rd</sup> doses is 3 months. It must be 6 months between the 1<sup>st</sup> and 3<sup>rd</sup> dose. Adolescents 11-14 years of age 1<sup>st</sup> dose and second dose given 6 months after 1<sup>st</sup> dose.

Trumenba: Offer to all males and females >16 years of age. The 2<sup>nd</sup> dose is given 6 months after the 1<sup>st</sup> dose.