

# Quantum Many-Body Simulations of Double Dot System

---

Alocias Mariadason

Institute of Physics

# Table of contents

1. Introduction
2. Methods
3. Implementation
4. Summary and Conclusion

# Introduction

---

# Quantum-Dot

- Small semiconductor nanostructures
- 2-10 nanometers with 10—50 particles

# Quantum-Dot Model

- Schrödinger equation
  - $\mathcal{H}|\psi\rangle = E|\psi\rangle$

# Quantum-Dot Model

- Schrödinger equation

- $\mathcal{H}|\psi\rangle = E|\psi\rangle$

- Hamiltonian

- $$\mathcal{H} = -\frac{1}{2} \sum_i \nabla_i^2 + \sum_{i < j} f(r_i, r_j) - \frac{1}{2} \sum_k \frac{\nabla_k^2}{M_k} + \sum_{k < l} g(R_k, R_l) + V(R, r)$$

# Quantum-Dot Model

- Schrödinger equation
  - $\mathcal{H}|\psi\rangle = E|\psi\rangle$
- Hamiltonian
  - $\mathcal{H} = -\frac{1}{2} \sum_i \nabla_i^2 + \sum_{i<j} f(r_j, r_j) - \frac{1}{2} \sum_k \frac{\nabla_k^2}{M_k} + \sum_{k<l} g(R_k, R_l) + V(R, r)$
- Born-Oppenheimer Approximation
  - Ignore Nuclei
  - $\sum_k \frac{\nabla_k^2}{M_k}$  gone
  - $\sum_{k<l} g(R_k, R_l)$  constant

# Quantum-Dot Model

- Schrödinger equation

- $\mathcal{H}|\psi\rangle = E|\psi\rangle$

- Hamiltonian

- $\mathcal{H} = -\frac{1}{2} \sum_i \nabla_i^2 + \sum_{i<j} f(r_j, r_j) - \frac{1}{2} \sum_k \frac{\nabla_k^2}{M_k} + \sum_{k<l} g(R_k, R_l) + V(R, r)$

- Born-Oppenheimer Approximation

- Ignore Nuclei

- $\sum_k \frac{\nabla_k^2}{M_k}$  gone

- $\sum_{k<l} g(R_k, R_l)$  constant

- $\mathcal{H} = -\frac{1}{2} \sum_i \nabla_i^2 + \sum_{i<j} f(r_j, r_j) + V(R, r)$



# Quantum-Dot Model

- Interaction



- $f(r_i, r_j) = \frac{1}{|r_i - r_j|}$

# Quantum-Dot Model

- Interaction: Coulomb repulsion
  - $f(r_i, r_j) = \frac{1}{|r_i - r_j|}$
- Confinement: Harmonic Oscillator<sup>1</sup>, Double-Well<sup>2</sup>

$$V(r) = \frac{1}{2} \omega m r^2$$

$$V(R, r) = \frac{1}{2} m \omega^2 (r^2 - \delta R |x| + R^2)$$



<sup>1</sup>S. Kvaal. "Harmonic Oscillator Eigenfunction Expansions, Quantum dots, and Effective Interactions". In: *Phys. Rev. B* 80 (4 2009), p. 045321.

<sup>2</sup>M. J. A. Schuetz et al. "Nuclear Spin Dynamics in Double Quantum Dots: Multistability, Dynamical Polarization, Criticality, and Entanglement". In: *Phys. Rev. B* 89 (19 2014), p. 195310.

**#NucleiHaveFeelingsTo**

# Methods

---

# Methods

---

- Hartree-Fock(HF)
- Variational Monte-Carlo(VMC)

# Implementation

---

## Summary and Conclusion

---

**Questions?**



Questions

Questions?