

Quantum Many-Body Simulations of Double Dot System

Alfred Alocias Mariadason

June 29, 2018

Contents

1. Introduction
2. Methods
3. Wavefunction
4. Implementation
5. Results
6. Summary and Conclusion

Introduction

Quantum-Dot Model

- Schrödinger equation: $\mathcal{H}|\psi\rangle = E|\psi\rangle$, $\mathcal{H} = -\sum_i \frac{\nabla_i^2}{2} + f(\mathbf{r}) + V(R, \mathbf{r})$

Quantum-Dot Model

- Schrödinger equation: $\mathcal{H}|\psi\rangle = E|\psi\rangle$, $\mathcal{H} = -\sum_i \frac{\nabla_i^2}{2} + f(\mathbf{r}) + V(R, \mathbf{r})$
- Interaction: $f(\mathbf{r}) = \sum_{i < j} \frac{1}{|r_i - r_j|}$

Quantum-Dot Model


- Schrödinger equation: $\mathcal{H}|\psi\rangle = E|\psi\rangle$, $\mathcal{H} = -\sum_i \frac{\nabla_i^2}{2} + f(\mathbf{r}) + V(\mathbf{R}, \mathbf{r})$
- Interaction: $f(\mathbf{r}) = \sum_{i < j} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}$
- Confinement: Harmonic Oscillator¹, Double-Well²

¹S. Kvaal. "Harmonic Oscillator Eigenfunction Expansions, Quantum dots, and Effective Interactions". In: *Phys. Rev. B* (2009).

²M. J. A. Schuetz et al. "Nuclear Spin Dynamics in Double Quantum Dots: Multistability, Dynamical Polarization, Criticality, and Entanglement". In: *Phys. Rev. B* (2014).

Quantum-Dot Model

- Schrödinger equation: $\mathcal{H}|\psi\rangle = E|\psi\rangle$, $\mathcal{H} = -\sum_i \frac{\nabla_i^2}{2} + f(\mathbf{r}) + V(\mathbf{R}, \mathbf{r})$
- Interaction: $f(\mathbf{r}) = \sum_{i < j} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}$
- Confinement: Harmonic Oscillator¹, Double-Well²
 $V(\mathbf{r}) = \frac{1}{2}m\omega^2 r^2$ $V(\mathbf{r}) = \frac{1}{2}m\omega^2(r^2 - \delta R|x| + R^2)$



¹S. Kvaal. "Harmonic Oscillator Eigenfunction Expansions, Quantum dots, and Effective Interactions". In: *Phys. Rev. B* (2009).

²M. J. A. Schuetz et al. "Nuclear Spin Dynamics in Double Quantum Dots: Multistability, Dynamical Polarization, Criticality, and Entanglement". In: *Phys. Rev. B* (2014).

Methods

**Hartree-Fock
Variational Monte-Carlo**

Methods: Variational Principle

$$E_0 \leq \frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle}$$

Slater Determinant and Energy Functional

Methods: Slater Determinant and Energy Functional

- Pauli Principle

Methods: Slater Determinant and Energy Functional

- Pauli Principle
- Slater Determinant

Methods: Slater Determinant and Energy Functional

- Pauli Principle
- Slater Determinant

- $\Psi_T^{AS} = \frac{1}{\sqrt{N!}} \sum_P (-1)^P \mathcal{P}_P \prod_i \psi_i$

Methods: Slater Determinant and Energy Functional

- Pauli Principle
- Slater Determinant

- $\Psi_T^{AS} = \frac{1}{\sqrt{N!}} \sum_P (-1)^P \mathcal{P}_P \prod_i \psi_i$

- $\Psi_T^S = \sqrt{\frac{\prod_{i=1}^N n_i!}{N!}} \sum_P \mathcal{P}_P \prod_i \psi_i$

Methods: Slater Determinant and Energy Functional

- Pauli Principle
- Slater Determinant

$$\bullet \Psi_T^{AS} = \frac{1}{\sqrt{N!}} \sum_P (-1)^P \mathcal{P}_P \prod_i \psi_i$$

$$\bullet \Psi_T^S = \sqrt{\frac{\prod_{i=1}^N n_i!}{N!}} \sum_P \mathcal{P}_P \prod_i \psi_i$$

$$\bullet E[\Psi] = \frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \sum_p \langle p | \mathcal{H}_0 | p \rangle + \frac{1}{2} \sum_{p,q} [\langle pq | f_{12} | pq \rangle \pm \langle pq | f_{12} | qp \rangle]$$



Hartree-Fock

Methods: Hartree-Fock

- Single Slater determinant wavefunction

Methods: Hartree-Fock

- Single Slater determinant wavefunction
- Constrained minimization

Methods: Hartree-Fock

- Single Slater determinant wavefunction
- Constrained minimization
 - Spin orthogonality: $\langle \psi_i | \psi_j \rangle = \delta_{ij}$

Methods: Hartree-Fock

- Single Slater determinant wavefunction
- Constrained minimization
 - Spin orthogonality: $\langle \psi_i | \psi_j \rangle = \delta_{ij}$
 - Fock-operator: $\mathcal{F} \equiv \mathcal{H}_0 + \mathcal{J} - \mathcal{K}$
 - $\mathcal{J} \equiv \langle \psi_k^* | f_{12} | \psi_k \rangle = \int \psi_k^*(\mathbf{r}) f_{12} \psi_k(\mathbf{r}) d\mathbf{r}$
 - $\mathcal{K} \equiv \langle \psi_k^* | f_{12} | \psi \rangle = \int \psi_k^*(\mathbf{r}) f_{12} \psi(\mathbf{r}) d\mathbf{r}$

Methods: Hartree-Fock

- Single Slater determinant wavefunction
- Constrained minimization
 - Spin orthogonality: $\langle \psi_i | \psi_j \rangle = \delta_{ij}$
 - Fock-operator: $\mathcal{F} \equiv \mathcal{H}_0 + \mathcal{J} - \mathcal{K}$
 - $\mathcal{J} \equiv \langle \psi_k^* | f_{12} | \psi_k \rangle = \int \psi_k^*(\mathbf{r}) f_{12} \psi_k(\mathbf{r}) d\mathbf{r}$
 - $\mathcal{K} \equiv \langle \psi_k^* | f_{12} | \psi \rangle = \int \psi_k^*(\mathbf{r}) f_{12} \psi(\mathbf{r}) d\mathbf{r}$
 - $\mathcal{F} |\psi\rangle = \epsilon |\psi\rangle, \epsilon = (\epsilon_0, \dots, \epsilon_N)$

$$\mathcal{F} \left| \psi \right\rangle = \pmb{\varepsilon} \left| \psi \right\rangle, \quad \pmb{\varepsilon} = (\varepsilon_0, \dots, \varepsilon_N)$$

Methods: Hartree-Fock

- Pair spins as: $\{\psi_{2l-1}, \psi_{2l}\} = \{\phi_l(\mathbf{r})\alpha(s), \phi_l(\mathbf{r})\beta(s)\}$

Methods: Hartree-Fock

- Pair spins as: $\{\psi_{2l-1}, \psi_{2l}\} = \{\phi_l(\mathbf{r})\alpha(s), \phi_l(\mathbf{r})\beta(s)\}$
- Expand: $\phi_l(\mathbf{r}) = \sum_{p=1}^L C_{pi} \chi_p(\mathbf{r})$

Methods: Hartree-Fock

- Pair spins as: $\{\psi_{2I-1}, \psi_{2I}\} = \{\phi_I(\mathbf{r})\alpha(s), \phi_I(\mathbf{r})\beta(s)\}$
- Expand: $\phi_i(\mathbf{r}) = \sum_{p=1}^L C_{pi} \chi_p(\mathbf{r})$
- Roothan-Hall: $\mathbf{F}\mathbf{C}_i = \boldsymbol{\varepsilon} S \mathbf{C}_i$
 - $F_{pq} = h_{pq} + \sum_{pq} \rho_{pq} (2\langle pq | f_{12} | rs \rangle - \langle pq | f_{12} | sr \rangle)$
 - $h_{pq} \equiv \langle p | h | q \rangle$
 - $\rho_{pq} \equiv \sum_{i=1}^{\frac{N}{2}} C_{pi} C_{qi}^*$
 - $S_{pq} \equiv \langle p | q \rangle$

Methods: Hartree-Fock

- Pair spins as: $\{\psi_{2I-1}, \psi_{2I}\} = \{\phi_I(\mathbf{r})\alpha(s), \phi_I(\mathbf{r})\beta(s)\}$
- Expand: $\phi_i(\mathbf{r}) = \sum_{p=1}^L C_{pi} \chi_p(\mathbf{r})$
- Roothan-Hall: $\mathbf{F}\mathbf{C}_i = \boldsymbol{\varepsilon} S \mathbf{C}_i$
 - $F_{pq} = h_{pq} + \sum_{pq} \rho_{pq} (2\langle pq|f_{12}|rs\rangle - \langle pq|f_{12}|sr\rangle)$
 - $h_{pq} \equiv \langle p|h|q\rangle$
 - $\rho_{pq} \equiv \sum_{i=1}^{\frac{N}{2}} C_{pi} C_{qi}^*$
 - $S_{pq} \equiv \langle p|q\rangle$
- Poople-Nesbet: $\mathbf{F}^+ \mathbf{C}^+ = \boldsymbol{\varepsilon}^+ S \mathbf{C}^+, \mathbf{F}^- \mathbf{C}^- = \boldsymbol{\varepsilon}^- S \mathbf{C}^-$

Variational Monte-Carlo

Methods: Variational Monte-Carlo

- Variational Principle

Methods: Variational Monte-Carlo

- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$

Methods: Variational Monte-Carlo

- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$

Methods: Variational Monte-Carlo

- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$
 - $P(R) \equiv \frac{|\Psi|^2}{\langle \Psi | \Psi \rangle}$

Methods: Variational Monte-Carlo

- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$
 - $P(R) \equiv \frac{|\Psi|^2}{\langle \Psi | \Psi \rangle}$
- Metropolis-Hastings Algorithm

Methods: Variational Monte-Carlo

- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$
 - $P(R) \equiv \frac{|\Psi|^2}{\langle \Psi | \Psi \rangle}$
- Metropolis-Hastings Algorithm
 - $r^{(b)} = r^{(a)} + \Delta t \xi$

Methods: Variational Monte-Carlo


- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$
 - $P(R) \equiv \frac{|\Psi|^2}{\langle \Psi | \Psi \rangle}$
- Metropolis-Hastings Algorithm
 - $r^{(b)} = r^{(a)} + \Delta t \xi$
 - $A_{a \rightarrow b} = \min\left(\frac{P_{a \rightarrow b}}{P_{b \rightarrow a}}, \frac{T_{a \rightarrow b}}{T_{b \rightarrow a}}, 1\right)$

Methods: Variational Monte-Carlo


- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$
 - $P(R) \equiv \frac{|\Psi|^2}{\langle \Psi | \Psi \rangle}$
- Metropolis-Hastings Algorithm
 - $r^{(b)} = r^{(a)} + \Delta t \xi$
 - $A_{a \rightarrow b} = \min\left(\frac{P_{a \rightarrow b}}{P_{b \rightarrow a}}, \frac{T_{a \rightarrow b}}{T_{b \rightarrow a}}\right), 1$
 - Importance Sampling
 - Focker-Planck equation and Langevin equation
 - $r^{(b)} = r^{(a)} + D \Delta t F^{(a)} + \sqrt{\Delta t} \xi$
 - Quantum force: $F = \frac{2}{\Psi} \nabla \Psi$
 - $\frac{T(b, a, \Delta t)}{T(a, b, \Delta t)} = \text{Greensfunction ratio}$

Minimization


Single-Well



Rosenbrock



Minimization: Gradient Descent



Minimization: Gradient Descent

x_0	γ	Iterations	x_m	$f(x_m)$
(5, 5)	0.9	20	(-0.072, -0.072)	0.010
(5, 5)	0.9	50	(-8.920×10^{-5} , -8.920×10^{-5})	1.591×10^{-8}
(5, 5)	0.9	100	(-1.273×10^{-9} , -1.273×10^{-9})	3.242×10^{-18}
(5, 5)	0.5	20	(0.0, 0.0)	0.0
(5, 5)	0.5	50	(0.0, 0.0)	0.0
(5, 5)	0.5	100	(0.0, 0.0)	0.0
(5, 5)	0.1	20	(0.072, 0.072)	0.010
(5, 5)	0.1	50	(8.920×10^{-5} , 8.920×10^{-5})	1.591×10^{-8}
(5, 5)	0.1	100	(1.273×10^{-9} , 1.273×10^{-9})	3.242×10^{-18}


x_0	γ	Iterations	x_m	$f(x_m)$
(0, 0.5)	0.001	100	(0.181, 0.030)	0.034
(0, 0.5)	0.001	500	(0.512, 0.258)	0.327
(0, 0.5)	0.001	1000	(0.675, 0.454)	0.106
(0, 0.5)	0.001	100000	(1.000, 1.000)	0.0
(0, 0.5)	0.0001	100	(0.027, 0.068)	1.399
(0, 0.5)	0.0001	500	(0.105, 0.009)	0.801
(0, 0.5)	0.0001	1000	(0.184, 0.031)	0.666
(0, 0.5)	0.0001	100000	(0.994, 0.989)	3.131×10^{-5}

Minimization: Gradient Descent

x_0	γ	Iterations	x_m	$f(x_m)$
(5, 5)	0.9	20	(-0.072, -0.072)	0.010
(5, 5)	0.9	50	(-8.920×10^{-5} , -8.920×10^{-5})	1.591×10^{-8}
(5, 5)	0.9	100	(-1.273×10^{-9} , -1.273×10^{-9})	3.242×10^{-18}
(5, 5)	0.5	20	(0.0, 0.0)	0.0
(5, 5)	0.5	50	(0.0, 0.0)	0.0
(5, 5)	0.5	100	(0.0, 0.0)	0.0
(5, 5)	0.1	20	(0.072, 0.072)	0.010
(5, 5)	0.1	50	(8.920×10^{-5} , 8.920×10^{-5})	1.591×10^{-8}
(5, 5)	0.1	100	(1.273×10^{-9} , 1.273×10^{-9})	3.242×10^{-18}

x_0	γ	Iterations	x_m	$f(x_m)$
(0, 0.5)	0.001	100	(0.181, 0.030)	0.034
(0, 0.5)	0.001	500	(0.512, 0.258)	0.327
(0, 0.5)	0.001	1000	(0.675, 0.454)	0.106
(0, 0.5)	0.001	100000	(1.000, 1.000)	0.0
(0, 0.5)	0.0001	100	(0.027, 0.068)	1.399
(0, 0.5)	0.0001	500	(0.105, 0.009)	0.801
(0, 0.5)	0.0001	1000	(0.184, 0.031)	0.666
(0, 0.5)	0.0001	100000	(0.994, 0.989)	3.131×10^{-5}

Minimization: Quasi-Newton BFGS



Minimization: Quasi-Newton BFGS

x_0	Iterations	x_m	$f(x_m)$
(1,1)	1	(-0.071,-0.071)	1.000
(-1,2)	1	(0.447,-0.894)	1.000
(1,1)	2	(0.000,0.000)	0.000
(-1,2)	2	(0.000,0.000)	0.000
(10,10)	1	(-0.071,-0.071)	1.000
(10,10)	2	(0.000,0.000)	0.000
(100,100)	1	(-0.071,-0.071)	1.000
(100,100)	2	(0.000,0.000)	0.000

x_0	Iterations	x_m	$f(x_m)$
(-0.5,2.0)	1	(-0.706,0.708)	7.280
(-0.5,2.0)	2	(-0.780,0.649)	3.342
(-0.5,2.0)	10	(0.238,0.051)	0.584
(-0.5,2.0)	30	(1.000,1,000)	0.000
(5.5,-10.0)	1	(-0.996,0.091)	85.214
(5.5,-10.0)	2	(-0.908,1.087)	10.549
(5.5,-10.0)	10	(0.027,0.012)	0.9613
(5.5,-10.0)	30	(1.000,1,000)	0.000

Minimization: Quasi-Newton BFGS

x_0	Iterations	x_m	$f(x_m)$
(1,1)	1	(-0.071,-0.071)	1.000
(-1,2)	1	(0.447,-0.894)	1.000
(1,1)	2	(0.000,0.000)	0.000
(-1,2)	2	(0.000,0.000)	0.000
(10,10)	1	(-0.071,-0.071)	1.000
(10,10)	2	(0.000,0.000)	0.000
(100,100)	1	(-0.071,-0.071)	1.000
(100,100)	2	(0.000,0.000)	0.000

x_0	Iterations	x_m	$f(x_m)$
(-0.5,2.0)	1	(-0.706,0.708)	7.280
(-0.5,2.0)	2	(-0.780,0.649)	3.342
(-0.5,2.0)	10	(0.238,0.051)	0.584
(-0.5,2.0)	30	(1.000,1,000)	0.000
(5.5,-10.0)	1	(-0.996,0.091)	85.214
(5.5,-10.0)	2	(-0.908,1.087)	10.549
(5.5,-10.0)	10	(0.027,0.012)	0.9613
(5.5,-10.0)	30	(1.000,1,000)	0.000

Minimization: Simulated Annealing

Wavefunction

Wavefunction

$$\phi_i(\mathbf{r}) = \sum_{p=1}^L C_{pi} \chi_p(\mathbf{r})$$

Wavefunction

$$\phi_i(\mathbf{r}) = \sum_{p=1}^L C_{pi} \chi_p(\mathbf{r})$$

Wavefunction: Integral Elements

$$\langle \phi_i(\mathbf{r}) | \phi_j(\mathbf{r}) \rangle$$

$$\langle \phi_i(\mathbf{r}) | x_d^k | \phi_j(\mathbf{r}) \rangle$$

$$\langle \phi_i(\mathbf{r}) | \nabla^2 | \phi_j(\mathbf{r}) \rangle$$

$$\langle \phi_i(\mathbf{r}_1) \phi_j(\mathbf{r}_2) | f_{12} | \phi_k(\mathbf{r}_1) \phi_l(\mathbf{r}_2) \rangle$$

Wavefunction: Single-Well

- Hermite Function: $\psi_n(\mathbf{r}) \equiv \prod_d N_d H_{n_d}(\sqrt{\omega}x_d) \exp\left(-\frac{\omega}{2}x_d^2\right)$

Wavefunction: Single-Well

- Hermite Function: $\psi_n(\mathbf{r}) \equiv \prod_d N_d H_{n_d}(\sqrt{\omega}x_d) \exp\left(-\frac{\omega}{2}x_d^2\right)$
- Solution in polar³

³E. Anisimovas and A. Matulis. “Energy spectra of few-electron quantum dots”. In: *Journal of Physics: Condensed Matter* (1998), p. 601.

Wavefunction: Single-Well

- Hermite Function: $\psi_n(\mathbf{r}) \equiv \prod_d N_d H_{n_d}(\sqrt{\omega}x_d) \exp\left(-\frac{\omega}{2}x_d^2\right)$
- Solution in polar³
- $\psi_n(\mathbf{r}) = \prod_d N_d \sum_{l=1}^{n_d} C_{n_d l}^{\text{Hermite}} g_l\left(\frac{\omega}{2}, \mathbf{r}, \mathbf{0}\right), \quad g_l\left(\frac{\omega}{2}, \mathbf{r}, \mathbf{0}\right) = x_d^{(l)} \exp\left(-\frac{\omega^2}{2}x_d^2\right)$

³E. Anisimovas and A. Matulis. “Energy spectra of few-electron quantum dots”. In: *Journal of Physics: Condensed Matter* (1998), p. 601.

Wavefunction: Single-Well

- Hermite Function: $\psi_n(\mathbf{r}) \equiv \prod_d N_d H_{n_d}(\sqrt{\omega}x_d) \exp\left(-\frac{\omega}{2}x_d^2\right)$
- Solution in polar³
- $\psi_n(\mathbf{r}) = \prod_d N_d \sum_{l=1}^{n_d} C_{n_d l}^{\text{Hermite}} g_l\left(\frac{\omega}{2}, \mathbf{r}, \mathbf{0}\right), \quad g_l\left(\frac{\omega}{2}, \mathbf{r}, \mathbf{0}\right) = x_d^{(l)} \exp\left(-\frac{\omega^2}{2}x_d^2\right)$
- Solution in Cartesian⁴

$$\langle g_i(\mathbf{r}) | g_j(\mathbf{r}) \rangle$$

$$\langle g_i(\mathbf{r}) | x_d' | g_j(\mathbf{r}) \rangle$$

$$\langle g_i(\mathbf{r}) | \nabla^2 | g_j(\mathbf{r}) \rangle$$

$$\langle g_i(\mathbf{r}_1) g_j(\mathbf{r}_2) | f_{12} | g_k(\mathbf{r}_1) g_l(\mathbf{r}_2) \rangle$$

³E. Anisimovas and A. Matulis. “Energy spectra of few-electron quantum dots”. In: *Journal of Physics: Condensed Matter* (1998), p. 601.

⁴J. Olsen T. Helgaker P. Jørgensen. *Molecular Electronic-Structure Theory*. Wiley, 2014. isbn: 978-0-47-196755-2. doi: 10.1002/9781119019572.

Wavefunction: Single-Well Integral Elements

$$\langle \psi_i^{\text{HO}} | \psi_j^{\text{HO}} \rangle = N_i \delta_{ij}$$

$$\langle \psi_i^{\text{HO}} | h^{\text{HO}} | \psi_j^{\text{HO}} \rangle = N_i \epsilon_i^{\text{HO}} \delta_{ij}$$

$$\langle \psi_i^{\text{HO}} \psi_j^{\text{HO}} | \frac{1}{r_{12}} | \psi_k^{\text{HO}} \psi_l^{\text{HO}} \rangle = \frac{aN_{ijkl}}{\sqrt{2\omega}} \sum_{tuvw}^{ijkl} H_{tuvw}^{ijkl} \sum_{pq}^{t+v, u+w} E_p^{tv} E_q^{uw} (-1)^q \xi_{p+q}(\frac{\omega}{2}, 0)$$

$$E_t^{i_d+1} = \frac{1}{2\omega} E_{t-1}^i \quad \xi_{t_d+1}^n = t_d \xi_{t_d-1}^{n+1}$$

$$E_0^0 = K_{AB} \quad \xi_0^n = (-b)^n \zeta_n(0)$$

$$\zeta_n^{\text{2D}}(x) = \int_{-1}^1 \frac{u^{2n}}{\sqrt{1-u^2}} e^{-u^2} du \quad \zeta_n^{\text{3D}}(x) = \int_{-1}^1 u^{2n} e^{-u^2} du$$

$$b = \begin{cases} \frac{\omega}{2}, & \text{2D} \\ \omega, & \text{3D} \end{cases}$$

Wavefunction: Double-Well

- Perturbation of harmonic oscillator: $U^{\text{DW}}(r) = V^{\text{HO}}(r) + V_n^{\text{DW}}(r)$

Wavefunction: Double-Well

- Perturbation of harmonic oscillator: $U^{\text{DW}}(r) = V^{\text{HO}}(r) + V_n^{\text{DW}}(r)$
- Expand in HO-functions: $|\psi_p^{\text{DW}}\rangle = \sum_l C_{lp}^{\text{DW}} |\psi_l^{\text{HO}}\rangle$

Wavefunction: Double-Well

- Perturbation of harmonic oscillator: $U^{\text{DW}}(r) = V^{\text{HO}}(r) + V_n^{\text{DW}}(r)$
- Expand in HO-functions: $\left| \psi_p^{\text{DW}} \right\rangle = \sum_l C_{lp}^{\text{DW}} \left| \psi_l^{\text{HO}} \right\rangle$
- Eigenvalue equation: $H^{\text{DW}} C^{\text{DW}} = \epsilon^{\text{DW}} C^{\text{DW}}$
 - $H_{ij}^{\text{DW}} = \epsilon_i^{\text{HO}} \delta_{ij} + \langle \psi_i^{\text{HO}} | V_n^{\text{DW}} | \psi_j^{\text{HO}} \rangle$

Wavefunction: Double-Well

- Perturbation of harmonic oscillator: $U^{\text{DW}}(r) = V^{\text{HO}}(r) + V_n^{\text{DW}}(r)$
- Expand in HO-functions: $\left| \psi_p^{\text{DW}} \right\rangle = \sum_l C_{lp}^{\text{DW}} \left| \psi_l^{\text{HO}} \right\rangle$
- Eigenvalue equation: $H^{\text{DW}} C^{\text{DW}} = \epsilon^{\text{DW}} C^{\text{DW}}$
 - $H_{ij}^{\text{DW}} = \epsilon_i^{\text{HO}} \delta_{ij} + \langle \psi_i^{\text{HO}} \left| V_n^{\text{DW}} \right| \psi_j^{\text{HO}} \rangle$
- Integral-Elements

$$\langle \psi_p^{\text{DW}} \left| \psi_q^{\text{DW}} \right\rangle = \delta_{pq}$$

$$\langle \psi_p^{\text{DW}} \left| h^{\text{DW}} \right| \psi_q^{\text{DW}} \rangle = \epsilon_p^{\text{DW}} \delta_{pq}$$

$$\left\langle \psi_p^{\text{DW}} \psi_q^{\text{DW}} \left| \frac{1}{r_{12}} \right| \psi_r^{\text{DW}} \psi_s^{\text{DW}} \right\rangle = \sum_{tuvw}^{ijkl} C_{tp}^{\text{DW}} C_{uq}^{\text{DW}} C_{vr}^{\text{DW}} C_{ws}^{\text{DW}} \left\langle \psi_t^{\text{HO}} \psi_u^{\text{HO}} \left| \frac{1}{r_{12}} \right| \psi_v^{\text{HO}} \psi_w^{\text{HO}} \right\rangle$$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi = \det(\Phi) J$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi = \det(\Phi) J$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{a\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{a\omega}x_d) e^{-\frac{a\omega}{2}x_d^2}$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi = \det(\Phi) J$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{\alpha\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{\alpha\omega}x_d) e^{-\frac{\alpha\omega}{2}x_d^2}$
- Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\omega}r_i)$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi = \det(\Phi) J$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{\alpha\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{\alpha\omega}x_d) e^{-\frac{\alpha\omega}{2}x_d^2}$
- Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\omega}r_i)$
- Modified Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\alpha\omega}r_i)$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi = \det(\Phi) J$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{\alpha\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{\alpha\omega}x_d) e^{-\frac{\alpha\omega}{2}x_d^2}$
- Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\omega}r_i)$
- Modified Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\alpha\omega}r_i)$
- Padé-Jastrow: $J_{\text{Padé}} = \prod_{i < j} \exp\left(\frac{a_{ij}r_{ij}}{1 + \beta r_{ij}}\right)$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi = \det(\Phi) J$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{\alpha\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{\alpha\omega}x_d) e^{-\frac{\alpha\omega}{2}x_d^2}$
- Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\omega}r_i)$
- Modified Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\alpha\omega}r_i)$
- Padé-Jastrow: $J_{\text{Padé}} = \prod_{i < j} \exp\left(\frac{a_{ij} r_{ij}}{1 + \beta r_{ij}}\right)$
- NQS: $J_{\text{NQS}} = e^{-\sum_{i=1}^N \frac{(r_i - \mathbf{a}_i)^2}{2\sigma^2}} \prod_j^M \left(1 + \exp\left(\mathbf{b}_j + \sum_{i=1}^N \sum_{d=1}^D \frac{x_i^{(d)} \mathbf{w}_{i+d,j}}{\sigma^2}\right) \right)$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi = \det(\Phi) J$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{\alpha\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{\alpha\omega}x_d) e^{-\frac{\alpha\omega}{2}x_d^2}$
- Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\omega}r_i)$
- Modified Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\alpha\omega}r_i)$
- Padé-Jastrow: $J_{\text{Padé}} = \prod_{i < j} \exp\left(\frac{a_{ij} r_{ij}}{1 + \beta r_{ij}}\right)$
- NQS: $J_{\text{NQS}} = e^{-\sum_{i=1}^N \frac{(r_i - a_i)^2}{2\sigma^2}} \prod_j^M \left(1 + \exp\left(b_j + \sum_{i=1}^N \sum_{d=1}^D \frac{x_i^{(d)} w_{i+d,j}}{\sigma^2}\right)\right)$
- Padé-NQS: $J = J_{\text{Padé}} J_{\text{NQS}}$

Implementation

Implementation

- C++ and Eigen

Implementation

- C++ and Eigen
 - Performance

Implementation

- C++ and Eigen
 - Performance
 - Generalization


Implementation

- C++ and Eigen
 - Performance
 - Generalization
- Python


Implementation

- C++ and Eigen
 - Performance
 - Generalization
- Python
 - Generate C++ code


Implementation: Cartesian




$(2, 0)$




$(1, 1)$




$(0, 2)$



$(1, 0)$




$(0, 1)$




$(0, 0)$

Implementation: Hartree-Fock




Implementation: Hartree-Fock




- Parallelization

Implementation: Hartree-Fock




- Parallelization
 - Two-body element is computationally expensive
 - $S_i = \sum_{j=0}^{P_i} \prod_d (n_{j_d} + 1)$

Implementation: Hartree-Fock




- Parallelization
 - Two-body element is computationally expensive
 - $S_i = \sum_{j=0}^{P_i} \prod_d (n_{j_d} + 1)$
 - Hartree-Fock algorithm only run on one process

Implementation: Hartree-Fock




- Parallelization
 - Two-body element is computationally expensive
 - $S_i = \sum_{j=0}^{P_i} \prod_d (n_{j_d} + 1)$
 - Hartree-Fock algorithm only run on one process
 - Tabulation of Two-Body matrix

Implementation: Variational Monte-Carlo




Implementation: Variational Monte-Carlo



- **Hermite** generated with Python and SymPy

Implementation: Variational Monte-Carlo







- **Hermite** generated with Python and SymPy
- Wavefunction class generated with Python


Results

Benchmark





Results: Benchmark



Results: Benchmark



Results: Benchmark



Results: Benchmark

ω [a.u]	N			
	2	6	12	20
0.1	0.4407(4)	3.5650(4)	12.3164(4)	30.0480(4)
0.28	1.0020(4)	7.6198(4)	25.5948(3)	61.8090(3)
0.5	1.6650(4)	11.8017(4)	39.3166(3)	93.9240(2)
1.0	3.0000(5)	20.2863(3)	68.1465(3)	156.2778(2)

ω [a.u]	N	
	2	8
0.1	0.50006(5)	5.80479(4)
0.28	1.20156(5)	12.48178(4)
0.5	2.00027(5)	19.33356(4)
1.0	3.72985(5)	33.30958(4)

$$\psi = \psi^{\text{HO}}(\sqrt{\alpha\omega}) J_{\text{Pad\'e}}$$

Results: Benchmark

ω [a.u]	N			
	2	6	12	20
0.1	0.46552(5){15}	3.70137(4){36}	12.64342(4){91}	-
0.28	1.04939(4){6}	7.89627(4){36}	26.21301(4){66}	62.93503(5){120}
0.5	1.70130(4){6}	12.02776(4){21}	39.76442(3){45}	95.21976(3){91}
1.0	3.05625(4){6}	20.45876(3){36}	66.37115(3){45}	157.41119(3){78}

ω [a.u]	N			
	2	6	12	20
0.10	0.44473(5){15}	3.63897(4){36}	12.46408(4){91}	-
0.28	1.04978(4){6}	7.72929(4){36}	25.96595(4){66}	62.65652(3){120}
0.50	1.66418(4){6}	11.97781(4){21}	39.57182(3){45}	94.76303(3){91}
1.00	3.00624(4){6}	20.38811(3){36}	66.28996(3){45}	157.46167(3){78}

$$\psi_p = \sum_l C_{lp} \psi_l^{\text{HO}}(\sqrt{\omega} r) J_{\text{Pad\'e}}, \quad \psi_p = \sum_l C_{lp} \psi_l^{\text{HO}}(\sqrt{\alpha \omega} r) J_{\text{Pad\'e}}$$

Results: Benchmark


ω	N	
	2	8
0.1	0.51122(5){70}	5.87372(4){120}
0.28	1.21844(5){70}	12.36177(4){168}
0.5	2.02030(4){20}	19.15006(4){112}
1.0	3.72918(5){20}	33.58046(4){168}

ω	N	
	2	8
0.1	0.50751(5){70}	5.84082(4){240}
0.28	1.20320(5){20}	12.37435(4){168}
0.5	2.01439(4){20}	19.09917(4){112}
1.0	3.72959(5){70}	33.04162(4){168}


$$\psi_p = \sum_l C_{lp} \psi_l^{\text{HO}}(\sqrt{\omega} r) J_{\text{Pad\'e}}, \quad \psi_p = \sum_l C_{lp} \psi_l^{\text{HO}}(\sqrt{\alpha\omega} r) J_{\text{Pad\'e}}$$

Results: Double-Well Hartree-Fock

2D



3D



Results: Double-Well Variational Monte-Carlo

ω	N			
	2	4	6	8
1.0	2.42238(4){10}	7.95247(4){42}	16.61419(4){44}	27.54453(3){50}

$$\psi_p = \sum_l C_{lp}^{\text{HF}} \sum_k C_{kl}^{\text{DW}} \psi_k^{\text{HO}} (\sqrt{\omega} r) J_{\text{Pad\'e}}$$

ω	N			
	2	4	6	8
1.0	2.36618(4){10}	7.90232(4){42}	16.55609(4){44}	27.58524(4){50}

$$\psi_p = \sum_l C_{lp}^{\text{HF}} \sum_k C_{kl}^{\text{DW}} \psi_k^{\text{HO}} (\sqrt{\alpha \omega} r) J_{\text{Pad\'e}}$$

Results: Double-Well Variational Monte-Carlo

ω	N		
	2	4	8
1.0	3.25118(4){11}	9.17489(4){17}	28.49671(4){26}

$$\psi_p = \sum_l C_{lp}^{\text{HF}} \sum_k C_{kl}^{\text{DW}} \psi_k^{\text{HO}} (\sqrt{\omega} r) J_{\text{Pad\'e}}$$





ω	N		
	2	4	8
1.0	3.22226(4){11}	9.17013(4){17}	28.62826(4){26}

$$\psi_p = \sum_l C_{lp}^{\text{HF}} \sum_k C_{kl}^{\text{DW}} \psi_k^{\text{HO}} (\sqrt{\alpha\omega} r) J_{\text{Pad\'e}}$$





Results: NQS-Jastrow

$$J_{\text{NQS}} = e^{-\sum_{i=1}^N \frac{(r_i - a_i)^2}{2\sigma^2}} \prod_j^M \left(1 + e^{b_j + \sum_{i=1}^N \sum_{d=1}^D \frac{x_i^{(d)} w_{i+d,j}}{\sigma^2}} \right)$$

Results: NQS-Jastrow Harmonic Oscillator



Results: NQS-Jastrow Double-Well





Summary and Conclusion

Summary

- Schrödinger equation: $\mathcal{H}|\psi\rangle = E|\psi\rangle$, $\mathcal{H} = -\sum_i \frac{\nabla_i^2}{2} + f(\mathbf{r}) + V(\mathbf{R}, \mathbf{r})$
- Interaction: $f(\mathbf{r}) = \sum_{i < j} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}$
- Confinement: Harmonic Oscillator, Double-Well

$$V(\mathbf{r}) = \frac{1}{2} m \omega^2 r^2$$

$$V(\mathbf{r}) = \frac{1}{2} m \omega^2 (r^2 - \delta R |x| + R^2)$$



**Hartree-Fock
Variational Monte-Carlo**

Summary

$$\begin{aligned}\left\langle \psi_i^{\text{HO}} \middle| \psi_j^{\text{HO}} \right\rangle &= N_i \delta_{ij} \\ \left\langle \psi_i^{\text{HO}} \middle| h^{\text{HO}} \right\rangle &= N_i \varepsilon_i^{\text{HO}} \delta_{ij}\end{aligned}$$

$$\left\langle \psi_i^{\text{HO}} \psi_j^{\text{HO}} \middle| \frac{1}{r_{12}} \right\rangle = \frac{aN_{ijkl}}{\sqrt{2\omega}} \sum_{tuvw}^{ijkl} H_{tuvw}^{ijkl} \sum_{pq}^{t+v, u+w} E_p^{tv} E_q^{uw} (-1)^q \xi_{p+q}(\frac{\omega}{2}, \mathbf{0})$$

$$\begin{aligned}E_t^{i_d+1} &= \frac{1}{2\omega} E_{t-1}^i & \xi_{t_d+1}^n &= t_d \xi_{t_d-1}^{n+1} \\ E_0^0 &= K_{AB} & \xi_0^n &= (-b)^n \zeta_n(0)\end{aligned}$$

$$\zeta_n^{\text{2D}}(x) = \int_{-1}^1 \frac{u^{2n}}{\sqrt{1-u^2}} e^{-u^2 x} du \quad \zeta_n^{\text{3D}}(x) = \int_{-1}^1 u^{2n} e^{-u^2 x} du$$

$$b = \begin{cases} \frac{\omega}{2}, & \text{2D} \\ \omega, & \text{3D} \end{cases}$$

Summary

- Perturbation of harmonic oscillator: $U^{\text{DW}}(r) = V^{\text{HO}}(r) + V_n^{\text{DW}}(r)$
- Expand in HO-functions: $\left| \psi_p^{\text{DW}} \right\rangle = \sum_l C_{lp}^{\text{DW}} \left| \psi_l^{\text{HO}} \right\rangle$
- Eigenvalue equation: $H^{\text{DW}} C^{\text{DW}} = \epsilon^{\text{DW}} C^{\text{DW}}$
 - $H_{ij}^{\text{DW}} = \epsilon_i^{\text{HO}} \delta_{ij} + \langle \psi_i^{\text{HO}} \left| V_n^{\text{DW}} \right| \psi_j^{\text{HO}} \rangle$
- Integral-Elements

$$\langle \psi_p^{\text{DW}} \left| \psi_q^{\text{DW}} \right\rangle = \delta_{pq}$$

$$\langle \psi_p^{\text{DW}} \left| h^{\text{DW}} \right| \psi_q^{\text{DW}} \rangle = \epsilon_p^{\text{DW}} \delta_{pq}$$

$$\left\langle \psi_p^{\text{DW}} \psi_q^{\text{DW}} \left| \frac{1}{r_{12}} \right| \psi_r^{\text{DW}} \psi_s^{\text{DW}} \right\rangle = \sum_{tuvw}^{ijkl} C_{tp}^{\text{DW}} C_{uq}^{\text{DW}} C_{vr}^{\text{DW}} C_{ws}^{\text{DW}} \left\langle \psi_t^{\text{HO}} \psi_u^{\text{HO}} \left| \frac{1}{r_{12}} \right| \psi_v^{\text{HO}} \psi_w^{\text{HO}} \right\rangle$$

Conclusion

- Single-Well and Double-well

Conclusion

- Single-Well and Double-well
- General C++ code implemented

Conclusion

- Single-Well and Double-well
- General C++ code implemented
- Hartree-Fock limit reached

Conclusion

- Single-Well and Double-well
- General C++ code implemented
- Hartree-Fock limit reached
- Hartree-Fock energies improved with Variational Monte-Carlo

Conclusion

- Single-Well and Double-well
- General C++ code implemented
- Hartree-Fock limit reached
- Hartree-Fock energies improved with Variational Monte-Carlo
- Stable minimization with NQS-Jastrow

Further Work

- Binding with Python

Further Work

- Binding with Python
- Rewrite with sparse matrices in Hartree-Fock

Further Work

- Binding with Python
- Rewrite with sparse matrices in Hartree-Fock
- Extend to non-isotropic gaussians

End

Questions?

Methods: Hartree-Fock

- Assumptions
 - The Born-Oppenheimer approximation holds.
 - All relativistic effects are negligible.
 - The wavefunction can be described by a single *Slater determinant*.
 - The Mean Field Approximation holds.

Methods: Hartree-Fock


- Assumptions
 - The Born-Oppenheimer approximation holds.
 - All relativistic effects are negligible.
 - The wavefunction can be described by a single *Slater determinant*.
 - The Mean Field Approximation holds.

Methods: Hartree-Fock


- Assumptions
 - The Born-Oppenheimer approximation holds.
 - All relativistic effects are negligible.
 - The wavefunction can be described by a single *Slater determinant*.
 - The Mean Field Approximation holds.

Methods: Hartree-Fock

- Assumptions
 - The Born-Oppenheimer approximation holds.
 - All relativistic effects are negligible.
 - The wavefunction can be described by a single *Slater determinant*.
 - The Mean Field Approximation holds.



Methods: Hartree-Fock



Methods: Variational Monte-Carlo

