

Quantum Many-Body Simulations of Double Dot System

Alfred Alocias Mariadason

June 27, 2018



Contents

1. Introduction
2. Methods
3. Wavefunction
4. Implementation
5. Results
6. Summary and Conclusion

Introduction

1. Small semiconductor nanostructures

Quantum-Dot Model

- Schrödinger equation

Quantum-Dot Model

- Schrödinger equation
 - $\mathcal{H}|\psi\rangle = E|\psi\rangle$

Quantum-Dot Model

- Schrödinger equation
 - $\mathcal{H}|\psi\rangle = E|\psi\rangle$
- Hamiltonian

Quantum-Dot Model

- Schrödinger equation
 - $\mathcal{H}|\psi\rangle = E|\psi\rangle$
- Hamiltonian
 - $$\mathcal{H} = -\frac{1}{2} \sum_i \nabla_i^2 + \sum_{i < j} f(\mathbf{r}_i, \mathbf{r}_j) - \frac{1}{2} \sum_k \frac{\nabla_k^2}{M_k} + \sum_{k < l} g(\mathbf{R}_k, \mathbf{R}_l) + V(\mathbf{R}, \mathbf{r})$$

Quantum-Dot Model

- Schrödinger equation
 - $\mathcal{H}|\psi\rangle = E|\psi\rangle$
- Hamiltonian
 - $$\mathcal{H} = -\frac{1}{2} \sum_i \nabla_i^2 + \sum_{i < j} f(\mathbf{r}_i, \mathbf{r}_j) - \frac{1}{2} \sum_k \frac{\nabla_k^2}{M_k} + \sum_{k < l} g(\mathbf{R}_k, \mathbf{R}_l) + V(\mathbf{R}, \mathbf{r})$$
- Born-Oppenheimer Approximation

Quantum-Dot Model

- Schrödinger equation
 - $\mathcal{H}|\psi\rangle = E|\psi\rangle$
- Hamiltonian
 - $$\mathcal{H} = -\frac{1}{2} \sum_i \nabla_i^2 + \sum_{i < j} f(\mathbf{r}_i, \mathbf{r}_j) - \frac{1}{2} \sum_k \frac{\nabla_k^2}{M_k} + \sum_{k < l} g(\mathbf{R}_k, \mathbf{R}_l) + V(\mathbf{R}, \mathbf{r})$$
- Born-Oppenheimer Approximation
 - Ignore Nuclei

Quantum-Dot Model

- Schrödinger equation
 - $\mathcal{H}|\psi\rangle = E|\psi\rangle$
- Hamiltonian
 - $$\mathcal{H} = -\frac{1}{2} \sum_i \nabla_i^2 + \sum_{i < j} f(\mathbf{r}_i, \mathbf{r}_j) - \frac{1}{2} \sum_k \frac{\nabla_k^2}{M_k} + \sum_{k < l} g(\mathbf{R}_k, \mathbf{R}_l) + V(\mathbf{R}, \mathbf{r})$$
- Born-Oppenheimer Approximation
 - Ignore Nuclei

Quantum-Dot Model

- Schrödinger equation
 - $\mathcal{H}|\psi\rangle = E|\psi\rangle$
- Hamiltonian
 - $$\mathcal{H} = -\frac{1}{2} \sum_i \nabla_i^2 + \sum_{i < j} f(\mathbf{r}_i, \mathbf{r}_j) - \frac{1}{2} \sum_k \frac{\nabla_k^2}{M_k} + \sum_{k < l} g(R_k, R_l) + V(\mathbf{R}, \mathbf{r})$$
- Born-Oppenheimer Approximation
 - Ignore Nuclei

Quantum-Dot Model

- Interaction:

Quantum-Dot Model

- Interaction:

- $f(\mathbf{r}_i, \mathbf{r}_j) = \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}$, Coulomb Repulsion

Quantum-Dot Model


- Interaction:
 - $f(r_i, r_j) = \frac{1}{|r_i - r_j|}$, Coulomb Repulsion
- Confinement: Harmonic Oscillator¹, Double-Well²

¹S. Kvaal. "Harmonic Oscillator Eigenfunction Expansions, Quantum dots, and Effective Interactions". In: *Phys. Rev. B* (2009).

²M. J. A. Schuetz et al. "Nuclear Spin Dynamics in Double Quantum Dots: Multistability, Dynamical Polarization, Criticality, and Entanglement". In: *Phys. Rev. B* (2014).

Quantum-Dot Model

- Interaction:
 - $f(r_i, r_j) = \frac{1}{|r_i - r_j|}$, Coulomb Repulsion
- Confinement: Harmonic Oscillator¹, Double-Well²
 $V(\mathbf{r}) = \frac{1}{2}m\omega^2 r^2$ $V(\mathbf{r}) = \frac{1}{2}m\omega^2(r^2 - \delta R|x| + R^2)$



¹S. Kvaal. "Harmonic Oscillator Eigenfunction Expansions, Quantum dots, and Effective Interactions". In: *Phys. Rev. B* (2009).

²M. J. A. Schuetz et al. "Nuclear Spin Dynamics in Double Quantum Dots: Multistability, Dynamical Polarization, Criticality, and Entanglement". In: *Phys. Rev. B* (2014).

Methods

Methods

Hartree-Fock
Variational Monte-Carlo

Methods: Variational Principle

$$E_0 \leq \frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle}$$

Slater Determinant and Energy Functional

Methods: Slater Determinant and Energy Functional

- Pauli Principle

Methods: Slater Determinant and Energy Functional

- Pauli Principle
- Slater Determinant

Methods: Slater Determinant and Energy Functional

- Pauli Principle
- Slater Determinant

- $\Psi_T^{AS} = \frac{1}{\sqrt{N!}} \sum_P (-1)^P \mathcal{P}_P \prod_i \psi_i$

Methods: Slater Determinant and Energy Functional

- Pauli Principle
- Slater Determinant

- $\Psi_T^{AS} = \frac{1}{\sqrt{N!}} \sum_P (-1)^P \mathcal{P}_P \prod_i \psi_i$

- $\Psi_T^S = \sqrt{\frac{\prod_{i=1}^N n_i!}{N!}} \sum_P \mathcal{P}_P \prod_i \psi_i$

Methods: Slater Determinant and Energy Functional

- Pauli Principle
- Slater Determinant

- $\Psi_T^{AS} = \frac{1}{\sqrt{N!}} \sum_P (-1)^P \mathcal{P}_P \prod_i \psi_i$

- $\Psi_T^S = \sqrt{\frac{\prod_{i=1}^N n_i!}{N!}} \sum_P \mathcal{P}_P \prod_i \psi_i$

- $E[\Psi] = \frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \sum_p \langle p | \mathcal{H}_0 | p \rangle + \frac{1}{2} \sum_{p,q} [\langle pq | f_{12} | pq \rangle \pm \langle pq | f_{12} | qp \rangle]$

Hartree-Fock

Methods: Hartree-Fock

- Assumptions

Methods: Hartree-Fock

- Assumptions
 - The Born-Oppenheimer approximation holds.


Methods: Hartree-Fock

- Assumptions
 - The Born-Oppenheimer approximation holds.
 - All relativistic effects are negligible.

Methods: Hartree-Fock

- Assumptions


- The Born-Oppenheimer approximation holds.
- All relativistic effects are negligible.
- The wavefunction can be described by a single *Slater determinant*.



Methods: Hartree-Fock

- Assumptions

- The Born-Oppenheimer approximation holds.
- All relativistic effects are negligible.
- The wavefunction can be described by a single *Slater determinant*.
- The Mean Field Approximation holds.



Methods: Hartree-Fock

- Constrained minimization

Methods: Hartree-Fock

- Constrained minimization
 - Spin orthogonality: $\langle \psi_i | \psi_j \rangle = \delta_{ij}$

Methods: Hartree-Fock

- Constrained minimization
 - Spin orthogonality: $\langle \psi_i | \psi_j \rangle = \delta_{ij}$
 - Fock-operator: $\mathcal{F} \equiv \mathcal{H}_0 + \mathcal{J} - \mathcal{K}$

Methods: Hartree-Fock

- Constrained minimization
 - Spin orthogonality: $\langle \psi_i | \psi_j \rangle = \delta_{ij}$
 - Fock-operator: $\mathcal{F} \equiv \mathcal{H}_0 + \mathcal{J} - \mathcal{K}$
 - $\mathcal{J} \equiv \langle \psi_k^* | f_{12} | \psi_k \rangle = \int \psi_k^*(\mathbf{r}) f_{12} \psi_k(\mathbf{r}) d\mathbf{r}$
 - $\mathcal{K} \equiv \langle \psi_k^* | f_{12} | \psi \rangle = \int \psi_k^*(\mathbf{r}) f_{12} \psi(\mathbf{r}) d\mathbf{r}$

Methods: Hartree-Fock

- Constrained minimization
 - Spin orthogonality: $\langle \psi_i | \psi_j \rangle = \delta_{ij}$
 - Fock-operator: $\mathcal{F} \equiv \mathcal{H}_0 + \mathcal{J} - \mathcal{K}$
 - $\mathcal{J} \equiv \langle \psi_k^* | f_{12} | \psi_k \rangle = \int \psi_k^*(\mathbf{r}) f_{12} \psi_k(\mathbf{r}) d\mathbf{r}$
 - $\mathcal{K} \equiv \langle \psi_k^* | f_{12} | \psi \rangle = \int \psi_k^*(\mathbf{r}) f_{12} \psi(\mathbf{r}) d\mathbf{r}$
 - $\mathcal{F} |\psi\rangle = \epsilon |\psi\rangle, \epsilon = (\epsilon_0, \dots, \epsilon_N)$

$$\mathcal{F} \left| \psi \right\rangle = \pmb{\varepsilon} \left| \psi \right\rangle, \quad \pmb{\varepsilon} = (\varepsilon_0, \dots, \varepsilon_N)$$

Methods: Hartree-Fock

- Integrate out spin

Methods: Hartree-Fock

- Integrate out spin
- Pair spins as: $\{\psi_{2l-1}, \psi_{2l}\} = \{\phi_l(\mathbf{r})\alpha(s), \phi_l(\mathbf{r})\beta(s)\}$

Methods: Hartree-Fock

- Integrate out spin
- Pair spins as: $\{\psi_{2I-1}, \psi_{2I}\} = \{\phi_I(\mathbf{r})\alpha(s), \phi_I(\mathbf{r})\beta(s)\}$
- Expand: $\phi_i(\mathbf{r}) = \sum_{p=1}^L C_{pi} \chi_p(\mathbf{r})$

Methods: Hartree-Fock

- Integrate out spin
- Pair spins as: $\{\psi_{2I-1}, \psi_{2I}\} = \{\phi_I(\mathbf{r})\alpha(s), \phi_I(\mathbf{r})\beta(s)\}$
- Expand: $\phi_i(\mathbf{r}) = \sum_{p=1}^L C_{pi} \chi_p(\mathbf{r})$
- Roothan-Hall: $\mathbf{FC}_i = \epsilon S \mathbf{C}_i$
 - $F_{pq} = h_{pq} + \sum_{pq} \rho_{pq} (2D_{prqs} \pm D_{prsq})$
 - $h_{pq} \equiv \langle p | h | q \rangle$
 - $\rho_{pq} \equiv \sum_{i=1}^{\frac{N}{2}} C_{pi} C_{qi}^*$
 - $D_{pqrs} \equiv \langle pq | f_{12} | rs \rangle$
 - $S_{pq} \equiv \langle p | q \rangle$

Methods: Hartree-Fock

- Integrate out spin
- Pair spins as: $\{\psi_{2I-1}, \psi_{2I}\} = \{\phi_I(\mathbf{r})\alpha(s), \phi_I(\mathbf{r})\beta(s)\}$
- Expand: $\phi_i(\mathbf{r}) = \sum_{p=1}^L C_{pi} \chi_p(\mathbf{r})$
- Roothan-Hall: $\mathbf{F}\mathbf{C}_i = \boldsymbol{\epsilon} \mathbf{S} \mathbf{C}_i$
 - $F_{pq} = h_{pq} + \sum_{pq} \rho_{pq} (2D_{prqs} \pm D_{prsq})$
 - $h_{pq} \equiv \langle p | h | q \rangle$
 - $\rho_{pq} \equiv \sum_{i=1}^{\frac{N}{2}} C_{pi} C_{qi}^*$
 - $D_{pqrs} \equiv \langle pq | f_{12} | rs \rangle$
 - $S_{pq} \equiv \langle p | q \rangle$
- Poople-Nesbet: $\mathbf{F}^+ \mathbf{C}^+ = \boldsymbol{\epsilon} \mathbf{S} \mathbf{C}^+, \mathbf{F}^- \mathbf{C}^- = \boldsymbol{\epsilon}^- \mathbf{S} \mathbf{C}^-$
 - $F_{pq}^\pm = h_{pq} + \sum_{k_\pm} \sum_{rs} C_{rk_\pm}^{\pm\dagger} C_{sk_\pm}^{\pm\dagger} [D_{prqs} - D_{prsq}] + \sum_{k_\mp} \sum_{rs} C_{rk_\mp}^{\mp\dagger} C_{sk_\mp}^{\mp\dagger} D_{prqs}$

Variational Monte-Carlo

Methods: Variational Monte-Carlo

- Variational Principle

Methods: Variational Monte-Carlo

- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$

Methods: Variational Monte-Carlo

- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$

Methods: Variational Monte-Carlo

- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$
 - $P(R) \equiv \frac{|\Psi|^2}{\langle \Psi | \Psi \rangle}$

Methods: Variational Monte-Carlo

- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$
 - $P(R) \equiv \frac{|\Psi|^2}{\langle \Psi | \Psi \rangle}$
- Metropolis-Hastings Algorithm

Methods: Variational Monte-Carlo

- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$
 - $P(R) \equiv \frac{|\Psi|^2}{\langle \Psi | \Psi \rangle}$
- Metropolis-Hastings Algorithm
 - $r^{(b)} = r^{(a)} + \Delta t \xi$

Methods: Variational Monte-Carlo


- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$
 - $P(R) \equiv \frac{|\Psi|^2}{\langle \Psi | \Psi \rangle}$
- Metropolis-Hastings Algorithm
 - $r^{(b)} = r^{(a)} + \Delta t \xi$
 - $A_{a \rightarrow b} = \min\left(\frac{P_{a \rightarrow b}}{P_{b \rightarrow a}}, \frac{T_{a \rightarrow b}}{T_{b \rightarrow a}}, 1\right)$

Methods: Variational Monte-Carlo


- Variational Principle
- Rewrite expectation value: $\frac{\langle \Psi | \mathcal{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \int \frac{\Psi^* \mathcal{H} \Psi}{\int \Psi^* \Psi dr} dr = \int \frac{|\Psi|^2}{\int \Psi^* \Psi dr} E_L dr$
 - $E_L(R; \alpha) \equiv \frac{1}{\Psi} \mathcal{H} \Psi$
 - $P(R) \equiv \frac{|\Psi|^2}{\langle \Psi | \Psi \rangle}$
- Metropolis-Hastings Algorithm
 - $r^{(b)} = r^{(a)} + \Delta t \xi$
 - $A_{a \rightarrow b} = \min\left(\frac{P_{a \rightarrow b}}{P_{b \rightarrow a}}, \frac{T_{a \rightarrow b}}{T_{b \rightarrow a}}, 1\right)$
 - Importance Sampling
 - $r^{\text{new}} = r^{\text{old}} + D \Delta t F^{\text{old}} + \sqrt{\Delta t} \xi$
 - $F = \frac{2}{\Psi} \nabla \Psi$
 - $\frac{T(b, a, \Delta t)}{T(a, b, \Delta t)} = \sum_i \exp\left(-\frac{(r_i^{(b)} - r_i^{(a)} - D \Delta t F_i^{(a)})^2}{4 D \Delta t} + \frac{(r_i^{(a)} - r_i^{(b)} - D \Delta t F_i^{(b)})^2}{4 D \Delta t}\right)$

Minimization


Single-Well



Rosenbrock



Minimization: Gradient Descent



Minimization: Gradient Descent

x_0	γ	Iterations	x_m	$f(x_m)$
(5, 5)	0.9	20	(-0.072, -0.072)	0.010
(5, 5)	0.9	50	(-8.920×10^{-5} , -8.920×10^{-5})	1.591×10^{-8}
(5, 5)	0.9	100	(-1.273×10^{-9} , -1.273×10^{-9})	3.242×10^{-18}
(5, 5)	0.5	20	(0.0, 0.0)	0.0
(5, 5)	0.5	50	(0.0, 0.0)	0.0
(5, 5)	0.5	100	(0.0, 0.0)	0.0
(5, 5)	0.1	20	(0.072, 0.072)	0.010
(5, 5)	0.1	50	(8.920×10^{-5} , 8.920×10^{-5})	1.591×10^{-8}
(5, 5)	0.1	100	(1.273×10^{-9} , 1.273×10^{-9})	3.242×10^{-18}


x_0	γ	Iterations	x_m	$f(x_m)$
(0, 0.5)	0.001	100	(0.181, 0.030)	0.034
(0, 0.5)	0.001	500	(0.512, 0.258)	0.327
(0, 0.5)	0.001	1000	(0.675, 0.454)	0.106
(0, 0.5)	0.001	100000	(1.000, 1.000)	0.0
(0, 0.5)	0.0001	100	(0.027, 0.068)	1.399
(0, 0.5)	0.0001	500	(0.105, 0.009)	0.801
(0, 0.5)	0.0001	1000	(0.184, 0.031)	0.666
(0, 0.5)	0.0001	100000	(0.994, 0.989)	3.131×10^{-5}

Minimization: Gradient Descent

x_0	γ	Iterations	x_m	$f(x_m)$
(5, 5)	0.9	20	(-0.072, -0.072)	0.010
(5, 5)	0.9	50	(-8.920×10^{-5} , -8.920×10^{-5})	1.591×10^{-8}
(5, 5)	0.9	100	(-1.273×10^{-9} , -1.273×10^{-9})	3.242×10^{-18}
(5, 5)	0.5	20	(0.0, 0.0)	0.0
(5, 5)	0.5	50	(0.0, 0.0)	0.0
(5, 5)	0.5	100	(0.0, 0.0)	0.0
(5, 5)	0.1	20	(0.072, 0.072)	0.010
(5, 5)	0.1	50	(8.920×10^{-5} , 8.920×10^{-5})	1.591×10^{-8}
(5, 5)	0.1	100	(1.273×10^{-9} , 1.273×10^{-9})	3.242×10^{-18}

x_0	γ	Iterations	x_m	$f(x_m)$
(0, 0.5)	0.001	100	(0.181, 0.030)	0.034
(0, 0.5)	0.001	500	(0.512, 0.258)	0.327
(0, 0.5)	0.001	1000	(0.675, 0.454)	0.106
(0, 0.5)	0.001	100000	(1.000, 1.000)	0.0
(0, 0.5)	0.0001	100	(0.027, 0.068)	1.399
(0, 0.5)	0.0001	500	(0.105, 0.009)	0.801
(0, 0.5)	0.0001	1000	(0.184, 0.031)	0.666
(0, 0.5)	0.0001	100000	(0.994, 0.989)	3.131×10^{-5}

Minimization: Quasi-Newton BFGS



Minimization: Quasi-Newton BFGS

x_0	Iterations	x_m	$f(x_m)$
(1,1)	1	(-0.071,-0.071)	1.000
(-1,2)	1	(0.447,-0.894)	1.000
(1,1)	2	(0.000,0.000)	0.000
(-1,2)	2	(0.000,0.000)	0.000
(10,10)	1	(-0.071,-0.071)	1.000
(10,10)	2	(0.000,0.000)	0.000
(100,100)	1	(-0.071,-0.071)	1.000
(100,100)	2	(0.000,0.000)	0.000

x_0	Iterations	x_m	$f(x_m)$
(-0.5,2.0)	1	(-0.706,0.708)	7.280
(-0.5,2.0)	2	(-0.780,0.649)	3.342
(-0.5,2.0)	10	(0.238,0.051)	0.584
(-0.5,2.0)	30	(1.000,1,000)	0.000
(5.5,-10.0)	1	(-0.996,0.091)	85.214
(5.5,-10.0)	2	(-0.908,1.087)	10.549
(5.5,-10.0)	10	(0.027,0.012)	0.9613
(5.5,-10.0)	30	(1.000,1,000)	0.000

Minimization: Quasi-Newton BFGS


x_0	Iterations	x_m	$f(x_m)$
(1,1)	1	(-0.071,-0.071)	1.000
(-1,2)	1	(0.447,-0.894)	1.000
(1,1)	2	(0.000,0.000)	0.000
(-1,2)	2	(0.000,0.000)	0.000
(10,10)	1	(-0.071,-0.071)	1.000
(10,10)	2	(0.000,0.000)	0.000
(100,100)	1	(-0.071,-0.071)	1.000
(100,100)	2	(0.000,0.000)	0.000

x_0	Iterations	x_m	$f(x_m)$
(-0.5,2.0)	1	(-0.706,0.708)	7.280
(-0.5,2.0)	2	(-0.780,0.649)	3.342
(-0.5,2.0)	10	(0.238,0.051)	0.584
(-0.5,2.0)	30	(1.000,1,000)	0.000
(5.5,-10.0)	1	(-0.996,0.091)	85.214
(5.5,-10.0)	2	(-0.908,1.087)	10.549
(5.5,-10.0)	10	(0.027,0.012)	0.9613
(5.5,-10.0)	30	(1.000,1,000)	0.000


Minimization: Simulated Annealing

Minimization: Simulated Annealing

Ackley



Rastrigin



Wavefunction

Wavefunction

$$\phi_i(\mathbf{r}) = \sum_{p=1}^L C_{pi} \chi_p(\mathbf{r})$$

Wavefunction

$$\phi_i(\mathbf{r}) = \sum_{p=1}^L C_{pi} \chi_p(\mathbf{r})$$

Wavefunction: Integral Elements

$$\langle \phi_i(\mathbf{r}) | \phi_j(\mathbf{r}) \rangle$$

$$\langle \phi_i(\mathbf{r}) | x_d^k | \phi_j(\mathbf{r}) \rangle$$

$$\langle \phi_i(\mathbf{r}) | \nabla^2 | \phi_j(\mathbf{r}) \rangle$$

$$\langle \phi_i(\mathbf{r}_1) \phi_j(\mathbf{r}_2) | f_{12} | \phi_k(\mathbf{r}_1) \phi_l(\mathbf{r}_2) \rangle$$

Wavefunction: Single-Well

- Hermite Function: $\psi_n(\mathbf{r}) \equiv \prod_d N_d H_{n_d}(\sqrt{\omega}x_d) \exp\left(-\frac{\omega}{2}x_d^2\right)$

Wavefunction: Single-Well

- Hermite Function: $\psi_n(\mathbf{r}) \equiv \prod_d N_d H_{n_d}(\sqrt{\omega}x_d) \exp\left(-\frac{\omega}{2}x_d^2\right)$
- Solution in polar³

³E. Anisimovas and A. Matulis. “Energy spectra of few-electron quantum dots”. In: *Journal of Physics: Condensed Matter* (1998), p. 601.

Wavefunction: Single-Well

- Hermite Function: $\psi_n(\mathbf{r}) \equiv \prod_d N_d H_{n_d}(\sqrt{\omega}x_d) \exp\left(-\frac{\omega}{2}x_d^2\right)$
- Solution in polar³
- $\psi_n(\mathbf{r}) = \prod_d N_d \sum_{l=1}^{n_d} C_{n_d l}^{\text{Hermite}} g_l\left(\frac{\omega}{2}, \mathbf{r}, \mathbf{0}\right), \quad g_l\left(\frac{\omega}{2}, \mathbf{r}, \mathbf{0}\right) = x_d^{(l)} \exp\left(-\frac{\omega^2}{2}x_d^2\right)$

³E. Anisimovas and A. Matulis. “Energy spectra of few-electron quantum dots”. In: *Journal of Physics: Condensed Matter* (1998), p. 601.

Wavefunction: Single-Well

- Hermite Function: $\psi_n(\mathbf{r}) \equiv \prod_d N_d H_{n_d}(\sqrt{\omega}x_d) \exp\left(-\frac{\omega}{2}x_d^2\right)$
- Solution in polar³
- $\psi_n(\mathbf{r}) = \prod_d N_d \sum_{l=1}^{n_d} C_{n_d l}^{\text{Hermite}} g_l\left(\frac{\omega}{2}, \mathbf{r}, \mathbf{0}\right), \quad g_l\left(\frac{\omega}{2}, \mathbf{r}, \mathbf{0}\right) = x_d^{(l)} \exp\left(-\frac{\omega^2}{2}x_d^2\right)$
- Solution in Cartesian⁴

$$\langle g_i(\mathbf{r}) | g_j(\mathbf{r}) \rangle$$

$$\langle g_i(\mathbf{r}) | x_d^k | g_j(\mathbf{r}) \rangle$$

$$\langle g_i(\mathbf{r}) | \nabla^2 | g_j(\mathbf{r}) \rangle$$

$$\langle g_i(\mathbf{r}_1) g_j(\mathbf{r}_2) | f_{12} | g_k(\mathbf{r}_1) g_l(\mathbf{r}_2) \rangle$$

³E. Anisimovas and A. Matulis. “Energy spectra of few-electron quantum dots”. In: *Journal of Physics: Condensed Matter* (1998), p. 601.

⁴J. Olsen T. Helgaker P. Jørgensen. *Molecular Electronic-Structure Theory*. Wiley, 2014. isbn: 978-0-47-196755-2. doi: 10.1002/9781119019572.

Wavefunction: Single-Well Integral Elements

$$\begin{aligned}\langle \psi_i^{\text{HO}} | \psi_j^{\text{HO}} \rangle &= N_i \delta_{ij} \\ \langle \psi_i^{\text{HO}} | h^{\text{HO}} | \psi_j^{\text{HO}} \rangle &= N_i \epsilon_i^{\text{HO}} \delta_{ij}\end{aligned}$$

$$\langle \psi_i^{\text{HO}} \psi_j^{\text{HO}} | \frac{1}{r_{12}} | \psi_k^{\text{HO}} \psi_l^{\text{HO}} \rangle = \frac{aN_{ijkl}}{\sqrt{2\omega}} \sum_{tuvw}^{ijkl} H_{tuvw}^{ijkl} \sum_{pq}^{t+v,u+w} E_p^{tv} E_q^{uw} (-1)^q \xi_{p+q} \left(\frac{\omega}{2}, \mathbf{0} \right)$$

$$E_t^{i+1,j} = \frac{1}{2\omega} E_{t-1}^{ij}$$

$$E_t^{i,j+1} = \frac{1}{2\omega} E_{t-1}^{ij}$$

$$E_0^{00} = K_{AB}$$

$$\xi_{t+1,u}^n = t \xi_{t-1,u}^{n+1}$$

$$\xi_{t+1,u,v}^n = t \xi_{t-1,u,v}^{n+1}$$

$$\xi_{t,u+1}^n = u \xi_{t,u-1}^{n+1}$$

$$\xi_{t,u+1,v}^n = u \xi_{t,u-1,v}^{n+1}$$

$$\xi_{00}^n = \left(\frac{-\omega}{2} \right)^n \zeta_n(0)$$

$$\xi_{t,u,v+1}^n = u \xi_{t,u,v-1}^{n+1}$$

$$\xi_{000}^n = (-\omega)^n \zeta_n(0)$$

$$\zeta_n(x) = \int_{-1}^1 \frac{u^{2n}}{\sqrt{1-u^2}} e^{-u^2} du$$

$$\zeta_n(x) = \int_{-1}^1 u^{2n} e^{-u^2} du$$

Wavefunction: Double-Well

- Perturbation of harmonic oscillator: $U^{\text{DW}}(r) = V^{\text{HO}}(r) + V_n^{\text{DW}}(r)$

Wavefunction: Double-Well

- Perturbation of harmonic oscillator: $U^{\text{DW}}(r) = V^{\text{HO}}(r) + V_n^{\text{DW}}(r)$
- Expand in HO-functions: $\left| \psi_p^{\text{DW}} \right\rangle = \sum_l C_{lp}^{\text{DW}} \left| \psi_l^{\text{HO}} \right\rangle$

Wavefunction: Double-Well

- Perturbation of harmonic oscillator: $U^{\text{DW}}(r) = V^{\text{HO}}(r) + V_n^{\text{DW}}(r)$
- Expand in HO-functions: $\left| \psi_p^{\text{DW}} \right\rangle = \sum_l C_{lp}^{\text{DW}} \left| \psi_l^{\text{HO}} \right\rangle$
- Eigenvalue equation: $H^{\text{DW}} C^{\text{DW}} = \epsilon^{\text{DW}} C^{\text{DW}}$
 - $H_{ij}^{\text{DW}} = \epsilon_i^{\text{HO}} \delta_{ij} + \langle \psi_i^{\text{HO}} | V_n^{\text{DW}} | \psi_j^{\text{HO}} \rangle$

Wavefunction: Double-Well

- Perturbation of harmonic oscillator: $U^{\text{DW}}(r) = V^{\text{HO}}(r) + V_n^{\text{DW}}(r)$
- Expand in HO-functions: $\left| \psi_p^{\text{DW}} \right\rangle = \sum_l C_{lp}^{\text{DW}} \left| \psi_l^{\text{HO}} \right\rangle$
- Eigenvalue equation: $H^{\text{DW}} C^{\text{DW}} = \epsilon^{\text{DW}} C^{\text{DW}}$
 - $H_{ij}^{\text{DW}} = \epsilon_i^{\text{HO}} \delta_{ij} + \langle \psi_i^{\text{HO}} \left| V_n^{\text{DW}} \right| \psi_j^{\text{HO}} \rangle$
- Integral-Elements

$$\langle \psi_p^{\text{DW}} \left| \psi_q^{\text{DW}} \right\rangle = \delta_{pq}$$

$$\langle \psi_p^{\text{DW}} \left| h^{\text{DW}} \right| \psi_q^{\text{DW}} \rangle = \epsilon_p^{\text{DW}} \delta_{pq}$$

$$\left\langle \psi_p^{\text{DW}} \psi_q^{\text{DW}} \left| \frac{1}{r_{12}} \right| \psi_r^{\text{DW}} \psi_s^{\text{DW}} \right\rangle = \sum_{tuvw}^{ijkl} C_{tp}^{\text{DW}} C_{uq}^{\text{DW}} C_{vr}^{\text{DW}} C_{ws}^{\text{DW}} \left\langle \psi_t^{\text{HO}} \psi_u^{\text{HO}} \left| \frac{1}{r_{12}} \right| \psi_v^{\text{HO}} \psi_w^{\text{HO}} \right\rangle$$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi_T = \det(\Phi(\mathbf{R}; \boldsymbol{\alpha}))\xi(s)$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi_T = \det(\Phi(\mathbf{R}; \boldsymbol{\alpha}))\xi(s)$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{\alpha\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{\alpha\omega}x_d) e^{-\frac{\alpha\omega}{2}x_d^2}$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi_T = \det(\Phi(\mathbf{R}; \boldsymbol{\alpha}))\xi(s)$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{\alpha\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{\alpha\omega}x_d) e^{-\frac{\alpha\omega}{2}x_d^2}$
- Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\omega}r_i)$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi_T = \det(\Phi(\mathbf{R}; \boldsymbol{\alpha}))\xi(s)$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{\alpha\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{\alpha\omega}x_d)e^{-\frac{\alpha\omega}{2}x_d^2}$
- Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\omega}r_i)$
- Modified Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\alpha\omega}r_i)$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi_T = \det(\Phi(\mathbf{R}; \boldsymbol{\alpha}))\xi(s)$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{\alpha\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{\alpha\omega}x_d)e^{-\frac{\alpha\omega}{2}x_d^2}$
- Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\omega}r_i)$
- Modified Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\alpha\omega}r_i)$
- Padé-Jastrow: $J_{\text{Padé}} = \prod_{i < j} e^{\frac{a_{ij} r_{ij}}{1 + \beta r_{ij}}}$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi_T = \det(\Phi(\mathbf{R}; \boldsymbol{\alpha})) \xi(s)$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{\alpha\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{\alpha\omega}x_d) e^{-\frac{\alpha\omega}{2}x_d^2}$
- Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\omega}r_i)$
- Modified Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\alpha\omega}r_i)$
- Padé-Jastrow: $J_{\text{Padé}} = \prod_{i < j} e^{\frac{a_{ij} r_{ij}}{1 + \beta r_{ij}}}$
- NQS: $J_{\text{NQS}} = e^{-\sum_{i=1}^N \frac{(r_i - a_i)^2}{2\sigma^2}} \prod_j^M \left(1 + e^{b_j + \sum_{i=1}^N \sum_{d=1}^D \frac{x_i^{(d)} w_{i+d,j}}{\sigma^2}} \right)$

Wavefunction: Slater-Jastrow

- Slater determinant: $\psi_T = \det(\Phi(\mathbf{R}; \boldsymbol{\alpha})) \xi(s)$
- Modified Hermite: $\Phi_{ij} = \psi_{n_j}^{\text{HO}}(\sqrt{\alpha\omega}r_i) = \prod_d N_d H_{n_d}(\sqrt{\alpha\omega}x_d) e^{-\frac{\alpha\omega}{2}x_d^2}$
- Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\omega}r_i)$
- Modified Hartree-Fock: $\Phi_{ij} = \sum_l C_{jl} \psi_{n_l}^{\text{HO}}(\sqrt{\alpha\omega}r_i)$
- Padé-Jastrow: $J_{\text{Padé}} = \prod_{i < j} e^{\frac{a_{ij} r_{ij}}{1 + \beta r_{ij}}}$
- NQS: $J_{\text{NQS}} = e^{-\sum_{i=1}^N \frac{(r_i - a_i)^2}{2\sigma^2}} \prod_j^M \left(1 + e^{b_j + \sum_{i=1}^N \sum_{d=1}^D \frac{x_j^{(d)} w_{i+d,j}}{\sigma^2}} \right)$
- Padé-NQS: $J = J_{\text{Padé}} J_{\text{NQS}}$

Implementation

Implementation

- C++ and Eigen

Implementation

- C++ and Eigen
 - Performance

Implementation

- C++ and Eigen
 - Performance
 - Generalization


Implementation

- C++ and Eigen
 - Performance
 - Generalization
- Python


Implementation

- C++ and Eigen
 - Performance
 - Generalization
- Python
 - Generate C++ code


Implementation: Cartesian




(2,0)




(1,1)




(0,2)



(1,0)




(0,1)




(0,0)

Implementation: Hartree-Fock




Implementation: Hartree-Fock




- Parallelization

Implementation: Hartree-Fock




- Parallelization
 - Two-body element is computationally expensive
 - $S_i = \sum_{j=0}^{P_i} \prod_d (n_{j_d} + 1)$

Implementation: Hartree-Fock




- Parallelization
 - Two-body element is computationally expensive
 - $S_i = \sum_{j=0}^{P_i} \prod_d (n_{jd} + 1)$
- Hartree-Fock algorithm only run on one process

Implementation: Hartree-Fock




- Parallelization
 - Two-body element is computationally expensive
 - $S_i = \sum_{j=0}^{P_i} \prod_d (n_{jd} + 1)$
- Hartree-Fock algorithm only run on one process
- Tabulation of Two-Body matrix

Implementation: Variational Monte-Carlo



Implementation: Variational Monte-Carlo




- **Hermite** generated with Python and SymPy

Implementation: Variational Monte-Carlo


- `set`: Called during initialization (before each sampling)
- `reSetAll`: Sets all matrices to zero (used in testing)
- `initializeMatrices`: Allocate memory
- `update`: Update positions and wavefunction
- `reset`: Revert to previous positions and wavefunction
- `resetGradient`: Revert to previous gradient
- `acceptState`: Update previous positions and wavefunction to current
- `acceptGradient`: Update previous gradient to current one

Implementation: Variational Monte-Carlo



- **Hermite** generated with Python and SymPy

Implementation: Variational Monte-Carlo







- Hermite generated with Python and SymPy
- Wavefunction class can be created with Python


Results

Benchmark





Results: Benchmark



Results: Benchmark



Results: Benchmark



Results: Benchmark

ω [a.u]	N			
	2	6	12	20
0.1	0.4407(4)	3.5650(4)	12.3164(4)	30.0480(4)
0.28	1.0020(4)	7.6198(4)	25.5948(3)	61.8090(3)
0.5	1.6650(4)	11.8017(4)	39.3166(3)	93.9240(2)
1.0	3.0000(5)	20.2863(3)	68.1465(3)	156.2778(2)

ω [a.u]	N	
	2	8
0.1	0.50006(5)	5.80479(4)
0.28	1.20156(5)	12.48178(4)
0.5	2.00027(5)	19.33356(4)
1.0	3.72985(5)	33.30958(4)

$$\psi = \psi^{\text{HO}}(\sqrt{\alpha\omega}) J_{\text{Pad\'e}}$$

Results: Benchmark

ω [a.u]	N			
	2	6	12	20
0.1	0.46552(5){15}	3.70137(4){36}	12.64342(4){91}	-
0.28	1.04939(4){6}	7.89627(4){36}	26.21301(4){66}	62.93503(5){120}
0.5	1.70130(4){6}	12.02776(4){21}	39.76442(3){45}	95.21976(3){91}
1.0	3.05625(4){6}	20.45876(3){36}	66.37115(3){45}	157.41119(3){78}

ω [a.u]	N			
	2	6	12	20
0.10	0.44473(5){15}	3.63897(4){36}	12.46408(4){91}	-
0.28	1.04978(4){6}	7.72929(4){36}	25.96595(4){66}	62.65652(3){120}
0.50	1.66418(4){6}	11.97781(4){21}	39.57182(3){45}	94.76303(3){91}
1.00	3.00624(4){6}	20.38811(3){36}	66.28996(3){45}	157.46167(3){78}

$$\psi_p = \sum_l C_{lp} \psi_l^{\text{HO}}(\sqrt{\omega} r) J_{\text{Pad\'e}}, \quad \psi_p = \sum_l C_{lp} \psi_l^{\text{HO}}(\sqrt{\alpha\omega} r) J_{\text{Pad\'e}}$$


Results: Benchmark

ω	N	
	2	8
0.1	0.51122(5){70}	5.87372(4){120}
0.28	1.21844(5){70}	12.36177(4){168}
0.5	2.02030(4){20}	19.15006(4){112}
1.0	3.72918(5){20}	33.58046(4){168}

ω	N	
	2	8
0.1	0.50751(5){70}	5.84082(4){240}
0.28	1.20320(5){20}	12.37435(4){168}
0.5	2.01439(4){20}	19.09917(4){112}
1.0	3.72959(5){70}	33.04162(4){168}

$$\psi_p = \sum_l C_{lp} \psi_l^{\text{HO}}(\sqrt{\omega} r) J_{\text{Pad\'e}}, \quad \psi_p = \sum_l C_{lp} \psi_l^{\text{HO}}(\sqrt{\alpha\omega} r) J_{\text{Pad\'e}}$$

Results: Double-Well Hartree-Fock



Results: Double-Well Variational Monte-Carlo

ω	N			
	2	4	6	8
1.0	2.42238(4){10}	7.95247(4){42}	16.61419(4){44}	27.54453(3){50}

$$\psi_p = \sum_l C_{lp}^{\text{HF}} \sum_k C_{kl}^{\text{DW}} \psi_k^{\text{HO}} (\sqrt{\omega} r) J_{\text{Pad\'e}}$$

ω	N			
	2	4	6	8
1.0	2.36618(4){10}	7.90232(4){42}	16.55609(4){44}	27.58524(4){50}

$$\psi_p = \sum_l C_{lp}^{\text{HF}} \sum_k C_{kl}^{\text{DW}} \psi_k^{\text{HO}} (\sqrt{\alpha \omega} r) J_{\text{Pad\'e}}$$

Results: Double-Well Variational Monte-Carlo

ω	N		
	2	4	8
1.0	3.25118(4){11}	9.17489(4){17}	28.49671(4){26}

$$\psi_p = \sum_l C_{lp}^{\text{HF}} \sum_k C_{kl}^{\text{DW}} \psi_k^{\text{HO}} (\sqrt{\omega} r) J_{\text{Pad\'e}}$$





ω	N		
	2	4	8
1.0	3.22226(4){11}	9.17013(4){17}	28.62826(4){26}

$$\psi_p = \sum_l C_{lp}^{\text{HF}} \sum_k C_{kl}^{\text{DW}} \psi_k^{\text{HO}} (\sqrt{\alpha \omega} r) J_{\text{Pad\'e}}$$



Results: NQS-Jastrow

$$J_{\text{NQS}} = e^{-\sum_{i=1}^N \frac{(r_i - a_i)^2}{2\sigma^2}} \prod_j^M \left(1 + e^{b_j + \sum_{i=1}^N \sum_{d=1}^D \frac{x_i^{(d)} w_{i+d,j}}{\sigma^2}} \right)$$

Results: NQS-Jastrow Harmonic Oscillator



Results: NQS-Jastrow Double-Well



Summary and Conclusion

Summary



- Interaction:

- $f(\mathbf{r}_i, \mathbf{r}_j) = \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}$, Coulomb Repulsion

- Confinement: Harmonic Oscillator, Double-Well

$$V(\mathbf{r}) = \frac{1}{2} m \omega^2 r^2$$

$$V(\mathbf{r}) = \frac{1}{2} m \omega^2 (r^2 - \delta R|x| + R^2)$$



**Hartree-Fock
Variational Monte-Carlo**

Summary

$$\begin{aligned}\left\langle \psi_i^{\text{HO}} \middle| \psi_j^{\text{HO}} \right\rangle &= N_i \delta_{ij} \\ \left\langle \psi_i^{\text{HO}} \middle| h^{\text{HO}} \right\rangle &= N_i \epsilon_i^{\text{HO}} \delta_{ij} \\ \left\langle \psi_i^{\text{HO}} \psi_j^{\text{HO}} \middle| \frac{1}{r_{12}} \right\rangle \left\langle \psi_k^{\text{HO}} \psi_l^{\text{HO}} \right\rangle &= \frac{aN_{ijkl}}{\sqrt{2\omega}} \sum_{tuvw}^{ijkl} H_{tuvw}^{ijkl} \sum_{pq}^{t+v,u+w} E_p^{tv} E_q^{uw} (-1)^q \xi_{p+q} \left(\frac{\omega}{2}, \mathbf{0} \right)\end{aligned}$$

$$E_t^{i+1,j} = \frac{1}{2(\alpha + \beta)} E_{t-1}^{ij} - \frac{\beta}{\alpha + \beta} (A_x - B_x) E_t^{ij} + (t+1) E_{t+1}^{ij}$$

$$E_t^{i,j+1} = \frac{1}{2(\alpha + \beta)} E_{t-1}^{ij} - \frac{\alpha}{\alpha + \beta} (A_y - B_y) E_t^{ij} + (t+1) E_{t+1}^{ij}$$

$$E_0^{00} = K_{AB}$$

$$\xi_{t+1,u}^n = t \xi_{t-1,u}^{n+1} + X_{AB} \xi_{t,u}^{n+1}$$

$$\xi_{t,u+1}^n = u \xi_{t,u-1}^{n+1} + Y_{AB} \xi_{t,u}^{n+1}$$

$$\xi_{00}^n = \left(\frac{-2\alpha\beta}{\alpha + \beta} \right)^n \zeta_n \left(\frac{\alpha\beta}{\alpha + \beta} R_{AB}^2 \right)$$

$$\zeta_n(x) = \int_{-1}^1 \frac{u^{2n}}{\sqrt{1-u^2}} e^{-u^2 x} du$$

$$\xi_{t+1,u,v}^n = t \xi_{t-1,u,v}^{n+1} + X_{AB} \xi_{t,u,v}^{n+1}$$

$$\xi_{t,u+1,v}^n = u \xi_{t,u-1,v}^{n+1} + Y_{AB} \xi_{t,u,v}^{n+1}$$

$$\xi_{t,u,v+1}^n = u \xi_{t,u,v-1}^{n+1} + Y_{AB} \xi_{t,u,v}^{n+1}$$

$$\xi_{000}^n = (-2\alpha\beta)^n \zeta_n \left(\frac{\alpha\beta}{\alpha + \beta} R_{AB}^2 \right)$$

$$\zeta_n(x) = \int_{-1}^1 u^{2n} e^{-u^2 x} du$$

Summary

- Perturbation of harmonic oscillator: $U^{\text{DW}}(r) = V^{\text{HO}}(r) + V_n^{\text{DW}}(r)$
- Expand in HO-functions: $\left| \psi_p^{\text{DW}} \right\rangle = \sum_l C_{lp}^{\text{DW}} \left| \psi_l^{\text{HO}} \right\rangle$
- Eigenvalue equation: $H^{\text{DW}} C^{\text{DW}} = \epsilon^{\text{DW}} C^{\text{DW}}$
 - $H_{ij}^{\text{DW}} = \epsilon_i^{\text{HO}} \delta_{ij} + \langle \psi_i^{\text{HO}} \left| V_n^{\text{DW}} \right| \psi_j^{\text{HO}} \rangle$
- Integral-Elements

$$\langle \psi_p^{\text{DW}} \left| \psi_q^{\text{DW}} \right\rangle = \delta_{pq}$$

$$\langle \psi_p^{\text{DW}} \left| h^{\text{DW}} \right| \psi_q^{\text{DW}} \rangle = \epsilon_p^{\text{DW}} \delta_{pq}$$

$$\left\langle \psi_p^{\text{DW}} \psi_q^{\text{DW}} \left| \frac{1}{r_{12}} \right| \psi_r^{\text{DW}} \psi_s^{\text{DW}} \right\rangle = \sum_{tuvw}^{ijkl} C_{tp}^{\text{DW}} C_{uq}^{\text{DW}} C_{vr}^{\text{DW}} C_{ws}^{\text{DW}} \left\langle \psi_t^{\text{HO}} \psi_u^{\text{HO}} \left| \frac{1}{r_{12}} \right| \psi_v^{\text{HO}} \psi_w^{\text{HO}} \right\rangle$$

Conclusion

- Single-Well and Double-well

Conclusion

- Single-Well and Double-well
- General C++ code implemented

Conclusion

- Single-Well and Double-well
- General C++ code implemented
- Hartree-Fock limit reached

Conclusion

- Single-Well and Double-well
- General C++ code implemented
- Hartree-Fock limit reached
- Hartree-Fock energies improved with Variational Monte-Carlo

Conclusion

- Single-Well and Double-well
- General C++ code implemented
- Hartree-Fock limit reached
- Hartree-Fock energies improved with Variational Monte-Carlo
- Stable minimization with NQS-Jastrow

Further Work

- Binding with Python

Further Work

- Binding with Python
- Optimize and clean-up code

Further Work

- Binding with Python
- Optimize and clean-up code
- Rewrite with sparse matrices in Hartree-Fock

Further Work


- Binding with Python
- Optimize and clean-up code
- Rewrite with sparse matrices in Hartree-Fock
- Extend to non-isotropic gaussians

Questions?

Questions

Questions?

Methods: Hartree-Fock



Methods: Variational Monte-Carlo

