

Work in progress

by

Alfred Alocias Mariadason

THESIS

for the degree of

MASTER OF SCIENCE



Faculty of Mathematics and Natural Sciences
University of Oslo

May 2018

TODO: write creative commons licence

Abstract

Work in progress

Acknowledgements

Work in progress

Contents

1		9
1	Introduction	9
1.1	bleh	9
2		11
1	Computational Quantum Mechanics	11
1.1	Metropolis-Hastings Algorithm	11

Symbols List

Work in progress make for $w_{i \rightarrow j} \equiv w(i|j)$

Source Code

Work in progress

Chapter 1

1 Introduction

1.1 bleh

Chapter 2

In this chapter we will address » LIST METHODS « regarding computational quantum mechanics and further deepen into Hartree-Fock methods and Variational Monte Carlo method. Optimization of calculation is also given while structure of program is given in » REF TO PROGRAM STRUCTURE CHAPTER «. General statistical theory used is given in » REF TO STATISTICS CHAPTER «

1 Computational Quantum Mechanics

1.1 Metropolis-Hastings Algorithm

The Metropolis algorithm bases itself on moves (also called transitions) as given in a Markov process. » REF THIS HERE «. This process is given by

$$w_i(t + \varepsilon) = \sum_j w_{i \rightarrow j} w_j(t) \quad (2.1)$$

where $w(j \rightarrow i)$ is just a transition from state j to state i . In order for the transition chain to reach a desired convergence while reversibility is kept, the well known condition for detailed balance must be fulfilled » REF HERE DETAILED BALANCE «. If detailed balance is true, then the following relations is true

$$w_i T_{i \rightarrow j} A_{i \rightarrow j} = w_j T_{j \rightarrow i} A_{j \rightarrow i} \Rightarrow \frac{w_i}{w_j} = \frac{T_{j \rightarrow i} A_{j \rightarrow i}}{T_{i \rightarrow j} A_{i \rightarrow j}} \quad (2.2)$$

We have here introduced two scenarios, the transition from configuration i to configuration j and the reverse process j to i . Solving the acceptance A for the two cases where the ratio in 2.2 is either 1 (in which case the proposed state j is accepted and transitions is made) and when the ratio is less then 1. The Metropolis algorithm would in this case not automatically reject the latter case, but rather reject it with a proposed uniform probability. Introducing now a probability distribution function(PDF) P the acceptance A can be expressed as

$$A_{i \rightarrow j} = \min(P_{i \rightarrow j} T_{i \rightarrow j}, 1) \quad (2.3)$$

The so-called selection probability T is defined specifically for each problem. For our case the PDF in question is the absolute square of the wave function and the selection T is a Green's function given in » REF GREENS «.