

Asymptotic Behavior of the Number of Interval Sums

Ohto Katila ohto.katila@gmail.com

May 9, 2026

Abstract

In this paper we extend the paper of O’Sullivan et al. on interval sums of prime powers to a class of regularly varying functions with index $\alpha > 1$, under mild additional assumptions. Let $n_f(x)$ be the length of the largest interval with sum less than x and $S_f(x)$ denote the number of interval sums less than x . Our main theorem states that $S_f(x) \sim n_f(x)^2 C(\alpha)$, where $C(\alpha)$ is constant dependent only on α . This paper applies tools from the theory of regularly varying functions and relates the number of blocks of length i to $\phi(x) = (1 + x^{\alpha+1})^{\frac{1}{\alpha+1}} - x$.

1 Introduction

Interval sum is the sum of consecutive terms in a sequence. Determining the asymptotic behavior of the count of interval sums less than x has been investigated for primes and prime powers [3, 4, 5]. Moser [3] determined the asymptotic behavior of interval sums of primes. Tongsomporn et al. [5] derived bounds for the number of interval sums of prime squares, though the bounds were not asymptotic to each other. O’Sullivan et al. [4] found asymptotic upper and lower bounds for the number of interval sums of prime powers.

To the best of our knowledge, no asymptotic formula is known for more general sequences or even prime powers. An asymptotic formula is found for the number of interval sums for a class of sequences $(f(i))_{i=1}^{\infty}$, where f is measurable, locally bounded away from 0 and ∞ , eventually increasing, and regularly varying with index $\alpha > 1$. This generalizes and extends O’Sullivan et al.’s work and provides exact asymptotics for the number of interval sums of prime powers.

Our main theorem provides an asymptotic for the interval sum counting function

$$S_f(x) = \#\left\{(i, j) \in \mathbb{N}^2 \mid i \leq j, f(i) + f(i+1) + \cdots + f(j) \leq x\right\}. \quad (1)$$

That is

$$S_f(x) \sim n_f(x)^2 C(\alpha) \quad x \rightarrow \infty \quad (2)$$

where $n := n_f(x)$ is the length of the largest interval, with following asymptotic relation to x , $x \sim \frac{nf(n)}{\alpha+1}$, $x \rightarrow \infty$ and

$$C(\alpha) = \frac{1}{2(\alpha+1)} \frac{\Gamma(\frac{1}{\alpha+1})\Gamma(\frac{\alpha-1}{\alpha+1})}{\Gamma(\frac{\alpha}{\alpha+1})}. \quad (3)$$

The proof relies on classical results from the theory of regular variation, in particular the uniform convergence theorem and Potter's bound [1]. The function $S_f(x)$ can be split into a sum of $s_{i,f}$ that counts the number of intervals of length i . The main idea in the proof is that $s_{i,f}$ can be related to the inverse of

$$\phi(x) = (1 + x^{\alpha+1})^{\frac{1}{\alpha+1}} - x \quad (4)$$

which allows us to calculate the exact coefficient in the asymptotic.

This paper is organized as follows. Section 2 defines our notation and some global assumptions. Section 3 covers preliminaries and one lemma. Section 4 proves a lemma that relates ϕ and $s_{i,f}$, a lemma that calculates the integral of the inverse of ϕ , one technical lemma bounding the sum of $s_{i,f}$ for small i and finally proves the main theorem.

2 Notation

- The function $f: [0, \infty) \rightarrow [0, \infty)$ is assumed to satisfy following criterion: measurable, eventually increasing, locally bounded away from 0 and ∞ , and regularly varying with index α (see definition 1 or [1]).
- $S_f(x) = \# \{(i, j) \in \mathbb{N}^2 \mid i \leq j, f(i) + f(i+1) + \dots + f(j) \leq x\}$ – the number of interval sums below x .
- $s_{k,f}(x) = \# \{(i, j) \in \mathbb{N}^2 \mid j - i = k - 1, f(i) + f(i+1) + \dots + f(j) \leq x\}$ – the number of interval sums of length k below x .
- $f(x) \sim g(x)$, $x \rightarrow a$ – meaning $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = 1$. The part " $x \rightarrow a$ " is often omitted when it is clear from context.
- $f(x) \lesssim g(x)$ – meaning $\limsup \frac{f(x)}{g(x)} \leq 1$.
- $f(x) \gtrsim g(x)$ – meaning $g(x) \lesssim f(x)$.
- sums of the form $\sum_{i=x}^y F(i)$ are denoted with $\sum_{x \leq i < y} F(i)$.
- $f(x) = O(g(x))$ – there exists $C > 0$ such that $|f(x)| \leq Cg(x)$ for all sufficiently large x .
- $f(x) = o(g(x))$ – shorthand for

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 0$$

- Define $n := n_f(x)$, where

$$n_f(x) = \max \left\{ n \in \mathbb{N} \mid \sum_{i=1}^n f(i) \leq x \right\} \quad (5)$$

that is, length of the largest interval.

3 Preliminaries

Definition 1. Function $f: [0, \infty) \rightarrow [0, \infty)$ is **regularly varying** with index α if for all $\lambda > 0$

$$\lim_{x \rightarrow \infty} \frac{f(\lambda x)}{f(x)} = \lambda^\alpha. \quad (6)$$

Definition 2. Function $L: [0, \infty) \rightarrow [0, \infty)$ is **slowly varying** if for all $\lambda > 0$

$$\lim_{x \rightarrow \infty} \frac{L(\lambda x)}{L(x)} = 1. \quad (7)$$

Since f is measurable and regularly varying with index α , we have $f(x) = x^\alpha L(x)$, for slowly varying L (see Theorem 1.4.1 in [1]).

Lemma 1. *We have*

$$x \sim \frac{n^{\alpha+1} L(n)}{\alpha + 1} = \frac{n f(n)}{\alpha + 1} \quad (8)$$

for $\alpha > -1$. Note $n := n_f(x)$ (see notation 5).

Proof. From Karamata's theorem (see proposition 1.5.8 in [1] or Proposition B.1 in the appendix)

$$\int_1^n f(i) \sim \frac{n f(n)}{\alpha + 1}. \quad (9)$$

From eventual monotonicity and regular variation with index $\alpha > -1$ of f it follows that

$$\sum_{i=1}^n f(i) \sim \int_1^n f(i) \quad (10)$$

From definition of $n = n_f(x)$ the result follows. $\square$

4 Proof of the Main Theorem

Lemma 2. *Let $\varepsilon > 0$ and $1 - \varepsilon \geq \lambda \geq \varepsilon$. Let $l = \phi^{-1}(\lambda)$ and $k = l + \lambda$, then uniformly in λ*

$$x \sim \sum_{i=n\phi^{-1}(\lambda)}^{n(\phi^{-1}(\lambda)+\lambda)} f(i) \quad (11)$$

where l is unique and $\phi(x) = (1 + x^{\alpha+1})^{\frac{1}{\alpha+1}} - x$.

Proof. We are going to consider sums of the form $\sum_{i=ln}^{kn} f(i)$. Let $l, k > 0$ be such that $k-l = \lambda$ and $k^{\alpha+1} - l^{\alpha+1} = 1$. Such k and l exists, by monotonicity of $(l+\lambda)^{\alpha+1} - l^{\alpha+1}$. Now Using mean-value theorem we get that for some $\zeta \in (l, k)$

$$1 = (l + \lambda)^{\alpha+1} - l^{\alpha+1} = (\alpha + 1)\zeta^{\alpha}\lambda > (\alpha + 1)l^{\alpha}\varepsilon. \quad (12)$$

Thus

$$l < \left(\frac{1}{(\alpha + 1)\varepsilon} \right)^{\frac{1}{\alpha}}. \quad (13)$$

To show that k, l are bounded below note that, if $l < \varepsilon$ then

$$1 = (l + \lambda)^{\alpha+1} - l^{\alpha+1} < (\varepsilon + \lambda)^{\alpha+1} - \varepsilon^{\alpha+1} = 1 - \varepsilon^{\alpha+1} < 1 \quad (14)$$

which is a contradiction.

Now k and l are in a compact subset of $(0, \infty)$, thus we can use the uniform convergence theorem (see Theorem 1.2.1 in [1] or Theorem B.1 in the Appendix). Using uniform convergence theorem and lemma 1

$$x \sim \frac{1}{\alpha + 1} L(n) n^{\alpha+1} = \quad (15)$$

$$\frac{1}{\alpha + 1} L(n) n^{\alpha+1} (k^{\alpha+1} - l^{\alpha+1}) = \quad (16)$$

$$L(n) \int_{nl}^{nk} t^{\alpha} dt \sim \quad (17)$$

$$L(n) \sum_{i=nl}^{nk} i^{\alpha} = \quad (18)$$

$$\sum_{i=nl}^{nk} i^{\alpha} L(n) \sim \sum_{i=nl}^{nk} f(i). \quad (19)$$

It follows that uniformly

$$x \sim \sum_{i=nl}^{nk} f(i). \quad (20)$$

Using $k^{\alpha+1} - l^{\alpha+1} = 1$ we get

$$k^{\alpha+1} - l^{\alpha+1} = 1 \quad (21)$$

$$k - l = (1 + l^{\alpha+1})^{\frac{1}{\alpha+1}} - l \quad (22)$$

$$\lambda = (1 + l^{\alpha+1})^{\frac{1}{\alpha+1}} - l \quad (23)$$

$$l = \phi^{-1}(\lambda) \quad (24)$$

In lemma 4 it is proven that the inverse of ϕ is well defined. The value of l is unique, because otherwise $k^{\alpha+1} - l^{\alpha+1} \neq 1$, thus same reasoning as 15 to 19 gives a contradiction. $\square$

Lemma 3. Let $\varepsilon > 0$ and $1 - \varepsilon \geq \lambda \geq \varepsilon$, then $s_{\lambda n, f}(x)$ has the following asymptotic uniformly

$$s_{\lambda n, f}(x) \sim n\phi^{-1}(\lambda) \quad (25)$$

where $\phi(x) = (1 + x^{\alpha+1})^{\frac{1}{\alpha+1}} - x$.

Proof. Since $s_{\lambda n, f}(x)$ counts the number of intervals of length λn , we get from lemma 2

$$s_{\lambda n, f}(x) \sim n\phi^{-1}(\lambda). \quad (26)$$

We will prove uniformity of the asymptotic by contradiction. Assume that it is not uniform then there exists $\varepsilon_0 > 0$ such that for all X_0 there exists $x > X_0$ and $\lambda_x \in [\varepsilon, 1 - \varepsilon]$ such that

$$\left| \frac{s_{\lambda_x n, f}(x)}{n\phi^{-1}(\lambda_x)} - 1 \right| > \varepsilon_0 \quad (27)$$

Now take sequence (x_m, λ_m) that satisfies 27. There are two cases

Case 1: $s_{\lambda_n, f}(x) > n\phi^{-1}(\lambda)(1 + \varepsilon_0)$

It would follow that

$$x \sim \sum_{i=s_{\lambda_n, f}(x)}^{s_{\lambda_n, f}(x)+n\lambda} f(i) > \sum_{i=n\phi^{-1}(\lambda)(1+\varepsilon_0)}^{n(\phi^{-1}(\lambda)(1+\varepsilon_0)+\lambda)} f(i) \quad (28)$$

We also have by same argument as steps from 19 to 16 that

$$\sum_{i=n\phi^{-1}(\lambda)(1+\varepsilon_0)}^{n(\phi^{-1}(\lambda)(1+\varepsilon_0)+\lambda)} f(i) \sim x((\phi^{-1}(\lambda)(1+\varepsilon_0)+\lambda)^{\alpha+1} - ((\phi^{-1}(\lambda)(1+\varepsilon_0))^{\alpha+1}) \quad (29)$$

We know that

$$\left(\phi^{-1}(\lambda)(1+\varepsilon_0)+\lambda \right)^{\alpha+1} - \left(\phi^{-1}(\lambda)(1+\varepsilon_0) \right)^{\alpha+1} = 1 + F(\lambda) \quad (30)$$

for some continuous F and that $F(\lambda) > 0$, hence it has minimum $c > 0$ in the compact λ -set. Thus independent λ we get that

$$x \gtrsim x(1+c) \quad (31)$$

which is a contradiction.

Case 2: $s_{\lambda_n, f}(x) < n\phi^{-1}(\lambda)(1 - \varepsilon_0)$

This is same argument as case 1. $\square$

Lemma 4. The function $\phi : (0, \infty) \rightarrow (0, 1)$, $\phi(x) = (1 + x^{\alpha+1})^{\frac{1}{\alpha+1}} - x$ is a strictly decreasing bijection, has an inverse, and

$$\int_0^1 \phi^{-1}(t) dt = \frac{1}{2(\alpha+1)} \frac{\Gamma(\frac{1}{\alpha+1})\Gamma(\frac{\alpha-1}{\alpha+1})}{\Gamma(\frac{\alpha}{\alpha+1})} \quad (32)$$

for $\alpha > 1$.

Proof. It is straightforward to verify that ϕ has an inverse and its integral converges. Doing change of variables and using analytic continuation of the beta function, gives the evaluation. For full details see the appendix A. $\square$

Lemma 5. *Let $\varepsilon > 0$, $\alpha > 1$, and $0 \leq l \leq 1 - \varepsilon$ then*

$$\sum_{i=nl}^{n(l+\varepsilon)} s_{i,f}(x) = O(n^2) \varepsilon^{\frac{1}{2}(1-\frac{1}{\alpha})}, \quad (33)$$

where the $O(n^2)$ term is independent of ε .

Proof. Since

$$x \geq f(i) + \dots + f(i+k-1) \geq kf(i) \quad (34)$$

$$f^{-1}\left(\frac{x}{k}\right) \geq i, \quad (35)$$

it follows that

$$s_{k,f}(x) \leq f^{-1}\left(\frac{x}{k}\right). \quad (36)$$

The inverse of a regularly varying function is regularly varying with index $\frac{1}{\alpha}$ (see Theorem 1.5.12 of [1] or Theorem B.3 in the Appendix). Let $f^{-1}(x) = x^{\frac{1}{\alpha}} \tilde{L}(x)$. We get from monotonicity of f that

$$\sum_{i=nl}^{n(l+\varepsilon)} f^{-1}\left(\frac{x}{i}\right) \sim \int_{nl}^{n(l+\varepsilon)} f^{-1}\left(\frac{x}{t}\right) dt = x^{\frac{1}{\alpha}} \int_{nl}^{n(l+\varepsilon)} t^{-\frac{1}{\alpha}} \tilde{L}\left(\frac{x}{t}\right) dt. \quad (37)$$

Since $t \leq n$, we get by Potter's bound (ii)

$$\tilde{L}\left(\frac{x}{t}\right) \leq M \tilde{L}\left(\frac{x}{n}\right) \left(\frac{n}{t}\right)^{\delta}, \quad (38)$$

where M is the constant from Potter's bound and $\delta = \frac{1}{2}(1 - \frac{1}{\alpha})$. Now we can bound the integral

$$x^{\frac{1}{\alpha}} \int_{nl}^{n(l+\varepsilon)} t^{-\frac{1}{\alpha}} \tilde{L}\left(\frac{x}{t}\right) dt \leq x^{\frac{1}{\alpha}} M \tilde{L}\left(\frac{x}{n}\right) n^{\delta} \int_{nl}^{n(l+\varepsilon)} t^{-\frac{1}{\alpha}-\delta} dt. \quad (39)$$

By definition

$$f^{-1}(f(n)) \sim n \quad (40)$$

$$n \tilde{L}(n^{\alpha} L(n)) L(n)^{1/\alpha} \sim n \quad (41)$$

$$\tilde{L}(n^{\alpha} L(n)) L(n)^{1/\alpha} \sim 1. \quad (42)$$

Using the relation between x and n obtained in Lemma 1

$$\tilde{L}\left(\frac{x}{n}\right) \sim \tilde{L}(n^\alpha L(n)). \quad (43)$$

Thus, from equations 43 and 42

$$x^{\frac{1}{\alpha}} M \tilde{L}\left(\frac{x}{n}\right) n^\delta \int_{nl}^{n(l+\varepsilon)} t^{-\frac{1}{\alpha}-\delta} \quad (44)$$

$$\sim M n^2 L(n)^{\frac{1}{\alpha}} \tilde{L}(n^\alpha L(n)) \frac{2\alpha}{\alpha-1} ((l+\varepsilon)^{\frac{1}{2}(1-\frac{1}{\alpha})} - l^{\frac{1}{2}(1-\frac{1}{\alpha})}) \quad (45)$$

$$\sim M n^2 \frac{2\alpha}{\alpha-1} ((l+\varepsilon)^{\frac{1}{2}(1-\frac{1}{\alpha})} - l^{\frac{1}{2}(1-\frac{1}{\alpha})}) \quad (46)$$

$$= O(n^2) ((l+\varepsilon)^{\frac{1}{2}(1-\frac{1}{\alpha})} - l^{\frac{1}{2}(1-\frac{1}{\alpha})}) \quad (47)$$

$$= O(n^2) \varepsilon^{\frac{1}{2}(1-\frac{1}{\alpha})}. \quad (48)$$

□

Theorem 1. *Under the assumptions and notation in 2, the function $S_f(x)$ has the asymptotic $n_f(x)^2 C(\alpha)$, $x \rightarrow \infty$ for $\alpha > 1$, where*

$$C(\alpha) = \frac{1}{2(\alpha+1)} \frac{\Gamma(\frac{1}{\alpha+1}) \Gamma(\frac{\alpha-1}{\alpha+1})}{\Gamma(\frac{\alpha}{\alpha+1})}. \quad (49)$$

Proof. Let $\varepsilon > 0$. From the definition of S and $s_{i,f}$ it follows that

$$S_f(x) = \sum_{i=1}^n s_{i,f}(x). \quad (50)$$

Splitting the sum

$$S_f(x) = \sum_{i=1}^{n\varepsilon} s_{i,f}(x) + \sum_{i=n\varepsilon}^{n(1-\varepsilon)} s_{i,f}(x) + \sum_{i=n(1-\varepsilon)}^n s_{i,f}(x). \quad (51)$$

Using Lemma 3 and continuity of ϕ^{-1} , we get

$$\sum_{i=n\varepsilon}^{n(1-\varepsilon)} s_{i,f}(x) \sim \sum_{i=n\varepsilon}^{n(1-\varepsilon)} n \phi^{-1}\left(\frac{i}{n}\right) \quad (52)$$

$$= n \sum_{i=n\varepsilon}^{n(1-\varepsilon)} \phi^{-1}\left(\frac{i}{n}\right) \quad (53)$$

$$\sim n \int_{n\varepsilon}^{n(1-\varepsilon)} \phi^{-1}\left(\frac{t}{n}\right) dt \quad (54)$$

$$= n^2 \int_{\varepsilon}^{1-\varepsilon} \phi^{-1}(t) dt. \quad (55)$$

Since the integral converges

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\varepsilon}^{1-\varepsilon} \phi^{-1}(t) dt = \int_0^1 \phi^{-1}(t) dt \quad (56)$$

From Lemma 5, it follows that the first and third sum in 51 are negligible in the asymptotic, hence taking $\varepsilon \rightarrow 0$ finishes the proof. $\square$

Corollary 1. *The number of interval sums for $f(x) = x^\alpha, \alpha > 1$ is asymptotic to*

$$x^{\frac{2}{1+\alpha}} (\alpha + 1)^{\frac{2}{1+\alpha}} C(\alpha) \quad (57)$$

Proof. From theorem 1 and inverting lemma 1, it follows that

$$S_f(x) \sim n^2 C(\alpha) \sim ((\alpha + 1)^{\frac{1}{1+\alpha}} x^{\frac{1}{1+\alpha}})^2 C(\alpha) \sim x^{\frac{2}{1+\alpha}} (\alpha + 1)^{\frac{2}{1+\alpha}} C(\alpha) \quad (58)$$

$\square$

Corollary 2. *Number of interval sums for prime powers $p^\alpha, \alpha > 1$ is asymptotic to*

$$(1 + \alpha)^2 C(\alpha) \left(\frac{x}{\log(x)^\alpha} \right)^{\frac{2}{\alpha+1}} \quad (59)$$

Proof. From the prime number theorem, it follows that $f(x) \sim (x \log(x))^\alpha$. Now using lemma 1 it follows that $x \sim \frac{1}{\alpha+1} n^{\alpha+1} \log^\alpha(n)$. It is straightforward to verify that $n = (1 + \alpha) \left(\frac{x}{\log^\alpha(x)} \right)^{\frac{1}{1+\alpha}}$ is the asymptotic inverse. By theorem 1 the corollary follows. $\square$

A Appendix: The Integral of the Function Phi

Here we give detailed evaluation of the integral of ϕ .

Proof. Taking the derivative of ϕ

$$\frac{d\phi}{dx} = (1 + x^{\alpha+1})^{-\frac{\alpha}{\alpha+1}} x^\alpha - 1 < 0 \quad (60)$$

As $\phi(0) = 1$ and $\lim_{x \rightarrow \infty} \phi(x) = 0$, ϕ is a bijection, hence it has an inverse. Since ϕ is a strictly decreasing bijection

$$\int_0^1 \phi^{-1}(x) dx = \int_0^\infty \phi(x) dx. \quad (61)$$

The integral converges as

$$\phi(x) = (1 + x^{\alpha+1})^{\frac{1}{\alpha+1}} - x < \frac{1}{1 + \alpha} x^{-\alpha}. \quad (62)$$

Doing a change of variables $t = \frac{x^{\alpha+1}}{1+x^{\alpha+1}}$, $x = (\frac{t}{1-t})^{\frac{1}{\alpha+1}}$, we get

$$\int_0^\infty \phi(x)dx = \int_0^1 (1-t)^{-\frac{1}{\alpha+1}} (1-t^{\frac{1}{\alpha+1}}) \frac{1}{\alpha+1} \left(\frac{t}{1-t}\right)^{-\frac{\alpha}{\alpha+1}} (1-t)^{-2} dt \quad (63)$$

$$= \frac{1}{\alpha+1} \int_0^1 (1-t)^{-\frac{2}{\alpha+1}-1} (t^{-\frac{\alpha}{\alpha+1}} - t^{\frac{1-\alpha}{\alpha+1}}) dt. \quad (64)$$

Let $g_r(t) = (1-t)^{r-\frac{2}{\alpha+1}-1} (t^{-\frac{\alpha}{\alpha+1}} - t^{\frac{1-\alpha}{\alpha+1}})$ and $g(t) = (1-t)^{-\frac{2}{\alpha+1}-1} (t^{-\frac{\alpha}{\alpha+1}} - t^{\frac{1-\alpha}{\alpha+1}})$. Since for each $r > 0$, $g_r(t) \rightarrow g(t)$, $r \rightarrow 0$ pointwise and $|g_r(t)| < |g(t)|$, we can use the dominated convergence theorem. Thus

$$\int_0^1 (1-t)^{-\frac{2}{\alpha+1}-1} (t^{-\frac{\alpha}{\alpha+1}} - t^{\frac{1-\alpha}{\alpha+1}}) dt = \lim_{r \rightarrow 0^+} \int_0^1 (1-t)^{r-\frac{2}{\alpha+1}-1} (t^{-\frac{\alpha}{\alpha+1}} - t^{\frac{1-\alpha}{\alpha+1}}) dt. \quad (65)$$

Let B be the beta function (see NIST DLMF § 5.12.1 [2])

$$B(z_1, z_2) = \int_0^1 t^{z_1-1} (1-t)^{z_2-1} dt = \frac{\Gamma(z_1)\Gamma(z_2)}{\Gamma(z_1+z_2)}. \quad (66)$$

The beta function can be analytically extended using Pochhammer's integral (see NIST DLMF § 5.12.12 [2]).

$$\int_0^\infty \phi(x)dx = \frac{1}{\alpha+1} \int_0^1 (1-t)^{-\frac{2}{\alpha+1}-1} (t^{-\frac{\alpha}{\alpha+1}} - t^{\frac{1-\alpha}{\alpha+1}}) dt \quad (67)$$

$$= \frac{1}{\alpha+1} \lim_{r \rightarrow 0^+} \int_0^1 (1-t)^{r-\frac{2}{\alpha+1}-1} (t^{-\frac{\alpha}{\alpha+1}} - t^{\frac{1-\alpha}{\alpha+1}}) dt \quad (68)$$

$$= \frac{1}{\alpha+1} \lim_{r \rightarrow 0^+} \left(B\left(\frac{1}{\alpha+1}, r - \frac{2}{\alpha+1}\right) - B\left(\frac{2}{\alpha+1}, r - \frac{2}{\alpha+1}\right) \right). \quad (69)$$

Using the gamma expression for beta functions, we get

$$\int_0^\infty \phi(x)dx = \frac{1}{\alpha+1} \lim_{r \rightarrow 0^+} \left(\frac{\Gamma(\frac{1}{\alpha+1})\Gamma(r - \frac{2}{\alpha+1})}{\Gamma(\frac{1}{\alpha+1} + r - \frac{2}{\alpha+1})} - \frac{\Gamma(\frac{2}{\alpha+1})\Gamma(r - \frac{2}{\alpha+1})}{\Gamma(\frac{2}{\alpha+1} + r - \frac{2}{\alpha+1})} \right) \quad (70)$$

$$= \frac{1}{\alpha+1} \lim_{r \rightarrow 0^+} \left(\frac{\Gamma(\frac{1}{\alpha+1})\Gamma(-\frac{2}{\alpha+1})}{\Gamma(-\frac{1}{\alpha+1})} - \frac{\Gamma(\frac{2}{\alpha+1})\Gamma(-\frac{2}{\alpha+1})}{\Gamma(r)} \right). \quad (71)$$

Since $\Gamma(t) = \frac{\Gamma(1+t)}{t}$, we have $\Gamma(t) \sim \frac{1}{t}$, $t \rightarrow 0$. Thus

$$\lim_{r \rightarrow 0} \frac{\Gamma(\frac{2}{\alpha+1})\Gamma(-\frac{2}{\alpha+1})}{\Gamma(r)} = 0 \quad (72)$$

so the expression simplifies to

$$\int_0^\infty \phi(x) dx = \frac{1}{\alpha+1} \frac{\Gamma(\frac{1}{\alpha+1})\Gamma(-\frac{2}{\alpha+1})}{\Gamma(-\frac{1}{\alpha+1})} \quad (73)$$

$$= \frac{1}{2(\alpha+1)} \frac{\Gamma(\frac{1}{\alpha+1})\Gamma(\frac{\alpha-1}{\alpha+1})}{\Gamma(\frac{\alpha}{\alpha+1})} \quad (74)$$

where the last equality follows from $z\Gamma(z) = \Gamma(z+1)$. $\square$

B Appendix: Selected Results on Regular Variation

For convenience, we reproduce several standard theorems from [1] we assume through this section that f and l are measurable.

Theorem B.1 (Uniform Convergence Theorem). *If l is slowly varying then*

$$\frac{l(\lambda x)}{l(x)} \rightarrow 1 \quad (x \rightarrow \infty)$$

uniformly in each compact λ -set in $(0, \infty)$.

Proof. See Theorem 1.2.1 in [1]. $\square$

Theorem B.2 (Potters Theorem). *(i) If l is slowly varying function then for any chosen constants $A > 1$, $\delta > 0$ there exists $X = X(A, \delta)$ such that*

$$\frac{l(y)}{l(x)} \leq A \max \left\{ \left(\frac{y}{x} \right)^\delta, \left(\frac{x}{y} \right)^\delta \right\} \quad (x \geq X, y \geq X)$$

(ii) If further, l is bounded away from 0 and ∞ in every compact subset of $[0, \infty)$, then for every $\delta > 0$ there exists $A' = A'(\delta) > 1$ such that

$$\frac{l(y)}{l(x)} \leq A' \max \left\{ \left(\frac{y}{x} \right)^\delta, \left(\frac{x}{y} \right)^\delta \right\} \quad (x \geq 0, y \geq 0)$$

(iii) If f is regularly varying of index ρ then for any chosen $A > 1$, $\delta > 0$ there exists $X = X(A, \delta)$ such that

$$\frac{f(y)}{f(x)} \leq A \max \left\{ \left(\frac{y}{x} \right)^{\rho+\delta}, \left(\frac{x}{y} \right)^{\rho+\delta} \right\} \quad (x \geq X, y \geq X)$$

Proof. See Theorem 1.5.6 in [1]. $\square$

Proposition B.1 (Karamata’s Theorem). *If l is slowly varying, X is so large that $l(x)$ is locally bounded in $[X, \infty)$, and $\alpha > -1$, then*

$$\int_X^x t^\alpha l(t) dt \sim \frac{x^{\alpha+1} l(x)}{\alpha + 1} \quad (x \rightarrow \infty)$$

Proof. See Proposition 1.5.8 in [1]. □

Definition B.1 (Generalized inverse). Generalized inverse of f is defined by

$$f^{\leftarrow}(x) = \inf \{x \mid f(x) = y\} \quad (75)$$

Theorem B.3. *If $f \in RV_\alpha$ with $\alpha > 0$, there exists $g \in RV_{\frac{1}{\alpha}}$ with*

$$f(g(x)) \sim g(f(x)) \sim x \quad (x \rightarrow \infty).$$

Here g (an asymptotic inverse of f) is determined uniquely to within asymptotic equivalence, and one version of g is $f^{\leftarrow}$.

Proof. See theorem 1.5.12 in [1]. □

References

- [1] N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular Variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1987.
- [2] *NIST Digital Library of Mathematical Functions*. <https://dlmf.nist.gov/>, Release 1.2.4 of 2025-03-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [3] L. Moser. On the sum of consecutive primes. *Canadian Mathematical Bulletin*, 6(2):159–161, 1963.
- [4] C. O’Sullivan, J. P. Sorenson, and A. Stahl. Algorithms and bounds on the sums of powers of consecutive primes. *Integers*, 24, 2024.
- [5] A. Tongsomporn, S. Wananiyakul, and J. Steuding. Sums of consecutive prime squares. *Integers*, 22, 2022.