



Session 4

# Lung absorption

MoBi<sup>®</sup> and PK-Sim<sup>®</sup> interaction

April 23, 2021

Erik Sjögren, PhD

Senior MIDD Consultant, Pharmetheus AB Associate Professor, Uppsala University



### Acknowledgement & Information

Johanna Eriksson, PhD
Hans Lennernäs, Prof. Uppsala University
Helena Thörn, PhD
... contributors at AstraZenca



European Journal of Pharmaceutics and Biopharmaceutics



Volume 156, November 2020, Pages 191-202

Pulmonary drug absorption and systemic exposure in human: Predictions using physiologically based biopharmaceutics modeling

Johanna Eriksson  $^{\rm a},$  Helena Thörn  $^{\rm b},$  Hans Lennernäs  $^{\rm a},$  Erik Sjögren  $^{\rm a} \stackrel{\textstyle \sim}{\sim} \boxtimes$ 

#### Show more 🗸

https://doi.org/10.1016/j.ejpb.2020.09.004

Get rights and content

Drug Absorption Parameters Obtained Using the Isolated Perfused Rat Lung Model Are Predictive of Rat In Vivo Lung Absorption.

Eriksson J, Sjögren E, Lennernäs H, Thörn H.

AAPS J. 2020 May 11;22(3):71. doi: 10.1208/s12248-020-00456-x.

PMID: 32394314 Free PMC article.

Pulmonary Dissolution of Poorly Soluble Compounds Studied in an ex Vivo Rat Lung Model.

Eriksson J, Thörn H, Sjögren E, Holmstén L, Rubin K, Lennernäs H.

Mol Pharm. 2019 Jul 1;16(7):3053-3064. doi: 10.1021/acs.molpharmaceut.9b00289. Epub 2019 Jun 11.

PMID: 31136181

Pulmonary absorption - estimation of effective pulmonary permeability and tissue retention of ten drugs using an ex vivo rat model and computational analysis.

Eriksson J, Sjögren E, Thörn H, Rubin K, Bäckman P, Lennernäs H.

Eur J Pharm Biopharm. 2018 Mar;124:1-12. doi: 10.1016/j.ejpb.2017.11.013. Epub 2017 Nov 27.

PMID: 29191716



#### Session 4

### Lung absorption: MoBi® to PK-Sim®




- Aim
- Background
  - Setting of investigation
  - Experimental model
  - Analysis of experimental data
- OSP Model development
- Application (Showcase MoBi–PK-Sim interaction)

### Aim

Show how experimental data and the approach of experimental data analysis can be used to further inform the development of a physiologically based translational model for pulmonary drug delivery.

Application of MoBi and PK-Sim to integrate experimental biopharmaceutics aspects to a PBPK framework.

Application of PK-Sim to evaluate population variability for a model developed in MoBi.



Increased leverage of generated pre-clinical ex-situ information for clinical predictions.



### Background – Setting of showcase

#### Lung administration

- Rapid onset of action
- High concentration at site of target
- Lower risk for systemic adverse effects
- Non-invasive method
- Little first pass effects

#### Drug delivery complexities

- Narrow and "sensitive" formulation space
- Device and patient dependencies
- Systemic concentrations for assessments

#### Context of presented work

- Formulation evaluation and optimization
- Assessments based on in vitro and pre-clinical data





### Background – Experimental model

#### Isolated Perfused Lung (IPL) experimental setup

- 1. Lungs and heart removed from rat
- 2. Pulmonary circulation perfused with a buffer
- 3. Lungs ventilated with negative pressure
- 4. Drug administered by tidal breathing

- Intact lung physiology
- No influence of systemic distribution
- Formulations as an aerosol
- Calculate dose






6




### Background – Analysis of experimental data



- 1




### Background – Analysis of experimental data





### Background – Analysis of experimental data

#### **Curve-fitting by estimation of:**






### OSP - Model development


Model developed in MoBi using PK-Sim PBPK model structure as back bone

Addition of IPL analysis structure

Human physiology

Addition of extra processes (particle distribution, MCC, trachea etc)







### OSP - Model development

- Particle size distribution
  - Input: Median and GSD → output: particle size distribution in 8 bins assuming log-normal distribution
  - Manual input also possible
- Particle dissolution:  $\frac{dm}{dt} = \frac{D \times SAi(t) \times (C_S C)}{h}$
- Deposition pattern fraction of dose in bronchial, alveolar and extrathoracic compartment (stomach)
- Mucociliary clearance of undissolved drug from bronchial compartment to stomach:
  - First order process  $(k_{mcc} = 0.58 h^{-1})$



### OSP - Model evaluation

The performance of the translational approach (input and model) evaluated towards clinical data after administration of solutions, suspensions and particulates.



European Journal of Pharmaceutics and **Biopharmaceutics** 



Get rights and content

Volume 156, November 2020, Pages 191-202

Pulmonary drug absorption and systemic exposure in human: Predictions using physiologically based biopharmaceutics modeling

Johanna Eriksson <sup>a</sup>, Helena Thörn <sup>b</sup>, Hans Lennernäs <sup>a</sup>, Erik Sjögren <sup>a</sup> △ ⊠ Show more V + Add to Mendeley <a Share >> Cite</a> https://doi.org/10.1016/j.ejpb.2020.09.004



### Showcase MoBi – PK-Sim interaction


#### Showcases

- MoBi ↔ PK-Sim
- Applications
  - Population dependencies
    - Permeability change (*smokers*)
    - Deposition change (asthmatics)
  - Drug dependencies
    - Solubility
    - · Particle size
  - Local lung vs systemic concentrations for the populations




### MoBi ↔ PK-Sim

#### PK-Sim → MoBi



Right click simulation Send to MoBi...

#### MoBi → PK-Sim





Right click simulation
Send simulation to PK-Sim for Population Simulation...



### MoBi ↔ PK-Sim

#### MoBi → PK-Sim








## Application


ParticleSize: 1- 5 µm

Solubility =  $100 \mu g/ml$ 


Normal


Cp-Range 5% to 95%
Clung-Range 5% to 95%
Cu cell-Range 5% to 95%
Cu ce






Solubility =  $1 \mu g/ml$ 







© Pharmetheus Confidential




### **Application**

ParticleSize: 2 -10 µm No change in deposition

Solubility =  $100 \mu g/ml$ 

Solubility =  $1 \mu g/ml$ 



Normal

0.1

0.01

0.001

0.0001

0.3



Cp-Range 5% to 95%

Clung-Range 5% to 95%

- Cp-Geometric Mean


Clung-Geometric Mean


1.5


Time [h]

- Cu.cell-Geometric Mean


Cu.cell-Range 5% to 95%












#### Permeability



© Pharmetheus Confidential



### Summary:

- Clinical predictions performed via direct physiologically based translation of pre-clinical ex-situ information.
- Integration of data analysis model structure dose reduce risks of model misconceptions and use of parameter input.
- System based virtual population simulations to assess dependencies using population database in OSP.

