

# **Accelerating Drug Discovery: systematic application of PBPK/PD modelling supporting candidate selection**

30-Sep-2025

Gregori Gerebtzoff



# Agenda




1. Introduction



2. Leveraging PK-Sim  
snapshots



3. Data and application



4. Outlook




# Introduction



# Motivation


- At Novartis, early-stage PBPK modeling on low molecular weight compounds is rarely done due to lack of experimental data, lack of a streamlined process, and challenge to get and process all required data
- Our overall goal is to enable a better understanding of distribution on a project basis, and to couple multi-scale modeling to pharmacodynamic models
- We developed an integrated, high-throughput PBPK platform connected to our corporate database and in-house machine learning models to execute and visualize simulations at scale, leveraging the OSP suite
- The workflow can be applied at all stages of drug development, including to yet-to-be-synthesized virtual compounds, and supports portfolio-wide screening for complex mechanisms underlying disposition

# Strengthening the Discovery PBPK strategy by developing a HT platform



# High-throughput generation of PK-Sim simulations

- We developed a set of Python packages to collect data, run simulations, process and visualize results, and run non-compartmental analysis on simulations to compare with observed data
- To run simulations, we leverage PK-Sim through **OSPSuite-R** functions, developed in collaboration with ESQ Labs. The **pksimHT** Python package is wrapping these R functions





# Leveraging PKSim's Snapshots



# OSP Open Systems Pharmacology - Snapshot

“A project snapshot contains the minimal amount of information required to recreate the project from scratch. This includes the information on primary substance specific input parameters (e.g. molecular properties like molecular weight, lipophilicity, etc.) and required inputs (e.g. demographic characteristics) for defining the system parameters.

Project snapshots are human-readable text files in JSON format.”

Source: <https://docs.open-systems-pharmacology.org/working-with-pk-sim/pk-sim-documentation/importing-exporting-project-data-models>

```
Internal {
  Version: 74,
  Individuals: [...],
  Populations: [
    {
      Name: "Healthy adults",
      Seed: 128991875,
      Settings: {
        NumberOfIndividuals: 100,
        ProportionOfFemales: 50,
        Age: {
          Min: 20.0,
          Max: 40.0,
          Unit: "year(s)"
        },
        Individual: {...}
      }
    }
  ],
  Compounds: [
    {
      Name: "SuperDrug",
      PlasmaProteinBindingPartner: "Albumin",
      Lipophilicity: [...],
      FractionUnbound: [...],
      Solubility: [...],
      CalculationMethods: [
        "Cellular partition coefficient method - PK-Sim Standard",
        "Cellular permeability - PK-Sim Standard"
      ],
      Parameters: [...]
    }
  ],
  Protocols: [
    {
      Name: "Bolus",
      ApplicationType: "IntravenousBolus",
      DosingInterval: "Single",
      Parameters: [...]
    }
  ],
  Simulations: [
    {
      Name: "S1",
      Model: "4Comp",
      Solver: { },
      OutputSchema: [...],
      Population: "Healthy adults",
      Compounds: [
        {
          Name: "SuperDrug",
          CalculationMethods: [
            "Cellular partition coefficient method - PK-Sim Standard",
            "Cellular permeability - PK-Sim Standard"
          ],
          Protocol: {
            Name: "Bolus"
          }
        }
      ],
      HasResults: false
    }
  ],
}
```



# What do our snapshots contain?

**Building Blocks**

- Individuals
  - Mouse
  - Rat
- Populations
- Compounds
  - propranolol
- Formulations
  - Default particle dissolution
  - Default Weibull
  - Solution
- Administration Protocols
  - IV 1 mg/kg
  - IV 1 mg/kg QD
  - IV 1.8 mg/kg QD
  - PO 1 mg/kg
  - PO 10 mg/kg
  - PO 3 mg/kg QD
- Events
- Observers
- Observed Data
  - propranolol
    - Mouse
      - IV
        - propranolol hydrochloride-2 - MAP-USCA-14-0261 - Mouse (Male) - Blood IV 1 mg/kg Solution: 100% Saline (Arm Group 1, n=3)
        - propranolol hydrochloride-2 - MAP-USCA-14-0262 - Mouse (Male) - Blood IV 1 mg/kg Solution: 100% Saline (Arm Group 1, n=2)
        - propranolol hydrochloride-2 - MAP-USCA-14-0263 - Mouse (Male) - Blood IV 1 mg/kg Solution: 100% Saline (Arm Group 1, n=2)

**propranolol hydrochloride-2 - MAP-CHBS-13-G001 - Rat (Male) - Blood IV 1 mg/kg Single (Other) Solution: NaCl [0.9%] (Arm Group 0, n=4)**

Molecular weight: 259.35 g/mol  
LLOQ: 0.31 nM

**Meta Data**

Name	Value
Molecule	propranolol
Species	Rat
Gender	Male
Study Id	MAP-CHBS-13-G001
Route	IV
Dose	1 mg/kg
Dose-normalized	False
Organ	Venous Blood
Compartment	Whole Blood
Dosage Form	Solution
Administration Frequency	Single
Sampling	Other
Study Type	In Vivo PK one-in-one
Arm Group	0

**Observed Data**

Time [h]	Concentration [nM]	Std Dev [nM]
0.08	777.00	82.40
0.25	309.00	37.30
0.50	119.00	20.20
1.00	44.20	7.97
2.00	12.50	0.11
3.00	4.35	0.56
4.00	2.41	0.29
6.00	1.03	0.35
8.00	0.54	0.35

**Simulations**









- Simulations
  - propranolol
    - Mouse
      - IV
        - propranolol - Mouse - IV 1 mg/kg
          - 4Comp
          - Mouse
          - propranolol
          - IV 1 mg/kg
          - propranolol hydrochloride-2 - MAP-USCA-14-0261
          - propranolol hydrochloride-2 - MAP-USCA-14-0262
          - propranolol hydrochloride-2 - MAP-USCA-14-0263
          - propranolol hydrochloride-2 - MAP-USCA-14-0294

**Concentration [nM] vs Time [h]**


Logarithmic Scale

# Getting *in vivo*, *in vitro* and *in silico* data from our corporate database

- *In vivo* concentration-time PK data are collected from 8 individual assays (various species in current and legacy systems)
- Physchem (pKa, solubility, lipophilicity) and ADME (plasma protein binding, intrinsic clearance, blood:plasma ratio,  $f_{u,mic}$ ) *in vitro* and *in silico* data are collected from 48 individual assays
- For each physchem and ADME property, a decision tree is implemented to return the “best” possible value
- The script can generate two flavors of the snapshot:
  - A compact version which only returns the “best” values
  - A complete version which return all available values, but the “best” value will be tagged accordingly

Experiment		Lipophilicity	Value Origin	Default	
Lipophilicity determination in 1-octanol/water	...	3.21 Log Units	Database-In Vitro-Potentiometric titration, average of 3 experiments	<input checked="" type="checkbox"/>	 
Direct logP/D in 1-octanol/buffer	...	2.80 Log Units	Database-In Vitro-Direct, from one experiment	<input type="checkbox"/>	 
High-Throughput Octanol/Water Partition Coefficient	...	3.40 Log Units	Database-In Vitro-High-throughput, from one experiment	<input type="checkbox"/>	 
NIBR logP and logD model (multi-task)	...	3.06 Log Units	Database-Other-In silico	<input type="checkbox"/>	 

## PPB decision tree



# Generate PKSim Snapshot files

## Arguments

### Preferred Assays

You can select the preferred assays for the PKSim snapshot. If you leave it blank (None), all assays will be included, and best selected.

### Cellular Partition Coefficient Method

CPC Method

Rodgers and Rowland

### Species, for observed values

Species

Mouse × Rat × Dog × Monkey ×

### Yes/No Options

- ☒ **Add Human?** Adds human ADME data to the snapshot. Note that you still have to add human simulations under 'Default simulations' if desired.
- ☐ **Normalize Dose?** Pulls dose-normalized experimental C-t data and sets all simulations to 1mg/kg (other than simulations specified in Add simulations below)
- ☒ **Use Legacy data?** Pulls data from three 'Legacy Assessment' assays and the Aragen Ct assay
- ☐ **Use Insilico data only?** Uses in silico options for all compound properties and doesn't pull observed data
- ☐ **Run NCA?** Calculates NCA clearance and other PK parameters from observed data (when available) and uses the observed NCA clearance to calculate hepatic clearance (other clearance option(s) will still be included in the snapshot)
- ☒ **Substitute BQL?** Replaces BQL values with LLOQ/sqrt(2).

### Simulation options

- ☒ Human (1mpk IV 3mpk PO single dose)
- ☐ Mouse (1mpk IV 3mpk PO single dose)
- ☐ Rat (1mpk IV 3mpk PO single dose)
- ☐ Dog (0.1mpk IV 0.3mpk PO single dose)
- ☐ Monkey (0.1mpk IV 0.3mpk IV single dose)

## NVP codes

NVP Codes

propranolol

You can enter multiple NVP codes, separated by commas.

Example: NVP-XYZ123, NVP-XYZ123, NVP-XYZ123


Submit NVP Codes

PKSim snapshot generated successfully!

PKSim Snapshot

```
{
  "Version": 79,
  "Description": "PKSim snapshot created for propranolol from CDF on 2025-09-23 10:49:48\r\nScript version 1.5.0",
  "Individuals": [
    {
      "Name": "Rat",
      "OriginData": {
        "CalculationMethods": {
          "SurfaceAreaPlsInt_VAR1": {}
        },
        "Species": "Rat",
        "Weight": {
          "Value": 0.312,
          "Unit": "kg",
          "ValueOrigin": {
            "Source": "Database",
            "Method": "InVivo",
            "Description": "Average of 2 animals from 1 study",
            "ExpressionProfiles": {}
          },
          "Name": "Mouse",
          "OriginData": {
            "CalculationMethods": {
              "SurfaceAreaPlsInt_VAR1": {}
            },
            "Species": "Mouse",
            "ExpressionProfiles": {}
          },
          "Name": "Human",
          "OriginData": {
            "CalculationMethods": {
              "SurfaceAreaPlsInt_VAR1": {}
            },
            "Body surface area - Mosteller",
            "Species": "Human",
            "Population": "European_ICRP_2002",
            "Gender": "MALE",
            "Age": {
              "Value": 30.0,
              "Unit": "year(s)",
              "Weight": {
                "Value": 70.0,
                "Unit": "kg",
                "ExpressionProfiles": {}
              },
              "Compounds": [
                {
                  "Name": "propranolol",
                  "IsSmallMolecule": true,
                  "PlasmaProteinBindingPartner": "Albumin",
                  "Lipophilicity": {
                    "Name": "Lipophilicity determination in 1-octanol/water",
                    "Parameters": {
                      "Name": "Lipophilicity",
                      "Value": 3.21,
                      "Unit": "Log Units",
                      "ValueOrigin": {
                        "Source": "Database",
                        "Method": "InVitro",
                        "Description": "Potentiometric titration, average of 3 experiments"
                      },
                      "Name": "Direct logP/D in 1-octanol/buffer",
                      "Parameters": {
                        "Name": "Lipophilicity",
                        "Value": 2.8,
                        "Unit": "Log Units",
                        "ValueOrigin": {
                          "Source": "Database",
                          "Method": "InVitro",
                          "Description": "Direct, from one experiment"
                        },
                        "IsDefault": false,
                        "Name": "High-Throughput Octanol/Water Partition Coefficient",
                        "Parameters": {
                          "Name": "Lipophilicity",
                          "Value": 3.4,
                          "Unit": "Log Units",
                          "ValueOrigin": {
                            "Source": "Database",
                            "Method": "InVitro",
                            "Description": "High-throughput, from one experiment"
                          },
                          "IsDefault": false,
                          "Name": "NIBR logP and logD model (multi-task)",
                          "Parameters": {
                            "Name": "Lipophilicity",
                            "Value": 3.06,
                            "Unit": "Log Units",
                            "ValueOrigin": {
                              "Source": "Database",
                              "Method": "Other",
                              "Description": "In silico"
                            },
                            "IsDefault": false,
                            "FractionUnbound": {
                              "Name": "Mouse Plasma Protein Binding (UC) (PPBUC)",
                              "Species": "Mouse",
                              "Parameters": {
                                "Name": "Fraction unbound (plasma, reference value)",
                                "Value": 0.1989,
                                "ValueOrigin": {
                                  "Source": "Database",
                                  "Method": "InVitro",
                                  "Description": "Mouse, ultracentrifugation, average of 85 experiments"
                                },
                                "Name": "Mouse NIBR-PPB",
                                "Species": "Mouse",
                                "Parameters": {
                                  "Name": "Fraction unbound (plasma, reference value)",
                                  "Value": 0.1735,
                                  "ValueOrigin": {
                                    "Source": "Database",
                                    "Method": "Other",
                                    "Description": "Mouse, in silico"
                                  },
                                  "IsDefault": false,
                                  "Name": "Rat HT-Dialysis Plasma Protein Binding",
                                  "Species": "Rat",
                                  "Parameters": {
                                    "Name": "Fraction unbound (plasma, reference value)",
                                    "Value": 0.153,
                                    "ValueOrigin": {
                                      "Source": "Database",
                                      "Method": "InVitro",
                                      "Description": "Rat, HT-Dialysis, from one experiment"
                                    },
                                    "Name": "Rat Plasma Protein Binding (UC) (PPBUC)",
                                    "Species": "Rat",
                                    "Parameters": {
                                      "Name": "Fraction unbound (plasma, reference value)",
                                      "Value": 0.2201,
                                      "ValueOrigin": {
                                        "Source": "Database",
                                        "Method": "InVitro",
                                        "Description": "Rat,
```

Download PKSim Snapshot





# PK-Sim Snapshots: “High-throughput” version

Building Blocks

Expression Profiles

Individuals

Populations

Compounds

Formulations

Administration Protocols

Events

Observers

Observed Data

propranolol

Mouse

IV

propranolol hydrochloride-2 - MAP-USCA-14-0261 - Mouse (Male) - Blood IV 1 mg/kg Solution

propranolol hydrochloride-2 - MAP-USCA-14-0262 - Mouse (Male) - Blood IV 1 mg/kg Solution

Simulations

IV

PO

propranolol - Rat - PO 1 mg/kg Suspension

4Comp

Rat

propranolol

PO 1 mg/kg

Default particle dissolution

propranolol hydrochloride-2 - MAP-CHBS-14-G611 - Rat (Male) - Blood PO 1 mg/kg Single

propranolol - Rat - PO 10 mg/kg Suspension

History

Comparison

Journal Diagram

Compound: 'propranolol' - PK-Sim® 12

Basic Physico-chemistry

ADME

Advanced Parameters

☒ Is small molecule

Lipophilicity:

Experiment	Lipophilicity	Value Origin
Lipophilicity determination in 1-octanol/water	3.21 Log Units	Database-In Vitro-Potentiometric titration, average of 3 experiments

Fraction unbound (plasma, reference value):

Bind to: ☒ Albumin ☐ α1-acid glycoprotein ☐ Unknown

Experiment	Fraction Unbound	Species	Value Origin	Default
Mouse Plasma Protein Binding (UC) (PPBUC)	0.20	Mouse	Database-In Vitro-Mouse, ultracentrifugation, a...	<input checked="" type="checkbox"/>
Rat HT-Dialysis Plasma Protein Binding	0.15	Rat	Database-In Vitro-Rat, HT-Dialysis, from one ex...	<input checked="" type="checkbox"/>

Molweight:

Molecular weight	259.35 g/mol
Has halogens	No
Effective molecular weight	259.35 g/mol
Value origin	Database-Other-SMR

Compound type / pKa:

Base	9.20
Acid	10.37
Neutral	<None>
Value origin	Database-Other-In silico (NIBR pKa Prediction)

Solubility:

Experiment	Solubility at Ref-pH	Ref-pH	Solubility gain per charge	pH-dependent Solubility	Value Origin
High Throughput Equilibrium Solu...	Edit Table			Show Graph	Database-In Vitro-Equilibrium so...

Project: Propranolol PKSim snapshot    Journal: Undefined

12 - Build 440

FileModelingParameter Identification & SensitivityWorking JournalImport/ExportUtilitiesViews

Expression ProfileIndividualsPopulationsCompoundsFormulationsAdministration ProtocolsEventsObserversObserved Data

Building Blocks

Simulationssimulations

HistoryComparisonJournal Diagram

Compound: 'propranolol' - PK-Sim® 12

PK-Sim Snapshots: “Precision” version

Compound: 'propranolol'

Basic Physico-chemistryADMEAdvanced Parameters

Is small molecule

Lipophilicity:

Experiment	Lipophilicity	Value Origin	Default
Lipophilicity determination in 1-octanol/water	3.21 Log Units	Database-In Vitro-Potentiometric titration, average of 3 experiments	<input checked="" type="checkbox"/>
Direct logP/D in 1-octanol/buffer	2.80 Log Units	Database-In Vitro-Direct, from one experiment	<input type="checkbox"/>
High-Throughput Octanol/Water Partition Coefficient	3.40 Log Units	Database-In Vitro-High-throughput, from one experiment	<input type="checkbox"/>
NIBR logP and logD model (multi-task)	3.06 Log Units	Database-Other-In silico	<input type="checkbox"/>

Fraction unbound (plasma, reference value):

Binds to:Albuminα1-acid glycoproteinUnknown

Experiment	Fraction Unbound	Species	Value Origin	Default
Mouse Plasma Protein Binding (UC) (PPBUC)	0.20	Mouse	Database-In Vitro-Mouse, ultracentrifugation, average of 85 experiments	<input checked="" type="checkbox"/>
Mouse NIBR-PPB	0.18	Mouse	Database-Other-Mouse, in silico	<input type="checkbox"/>
Rat HT-Dialysis Plasma Protein Binding	0.15	Rat	Database-In Vitro-Rat, HT-Dialysis, from one experiment	<input checked="" type="checkbox"/>
Rat Plasma Protein Binding (UC) (PPBUC)	0.22	Rat	Database-In Vitro-Rat, ultracentrifugation, average of 19 experiments	<input type="checkbox"/>
Rat MAP RED Plasma Protein Binding	0.16	Rat	Database-In Vitro-Rat, rapid equilibrium dialysis, average of 6 experiments	<input type="checkbox"/>
Rat MAP RED Protein Binding: Non-standard Assay	0.24	Rat	Database-In Vitro-Rat, non-standard assay, average of 2 experiments	<input type="checkbox"/>
Rat NIBR-PPB	0.19	Rat	Database-Other-Rat, in silico	<input type="checkbox"/>

Molweight:

Molecular weight259.35 g/mol

Has halogensNo

Effective molecular weight259.35 g/mol

Value originDatabase-Other-SMR

Compound type / pKa:

Base9.20

Acid10.37

Neutral<None>

Value originDatabase-Other-In silico (NIBR pKa Prediction)

Solubility:

Experiment	Solubility at Ref-pH	Ref-pH	Solubility gain per charge	pH-dependent Solubility	Value Origin	Default
High Throughput Equilibrium Solubility	Edit Table			Show Graph	Database-In Vitro-Equilibrium solubility	<input checked="" type="checkbox"/>
MELLODDY solubility models	Edit Table			Show Graph	Database-Other-in silico (High Throughput Equilibri...	<input type="checkbox"/>

propranolol hydrochloride-2 - MAP-USCA-14-0261 - Mouse (Male) - Blood IV 1 mg/kg Solution

propranolol hydrochloride-2 - MAP-USCA-14-0262 - Mouse (Male) - Blood IV 1 mg/kg Solution

propranolol - Rat - PO 1 mg/kg Suspension

4Comp

Rat

propranolol

PO 1 mg/kg

Default particle dissolution

propranolol hydrochloride-2 - MAP-CHBS-14-G611 - Rat (Male) - Blood PO 1 mg/kg Single

propranolol - Rat - PO 10 mg/kg Suspension

Project: Propranolol PKSim snapshotJournal: Undefined12 - Build 440



# Data and application






# Small/medium-scale use cases

On a project basis, our workflow allows to perform quick checks of whether compounds are “well-behaved”



Can be predicted well by bottom-up Vss & scaled LM CL<sub>int</sub>




Requires further investigation in PK-Sim GUI (alternative CL & Vss mechanisms, transporters)


# Large-scale use case

- Bottom-up PBPK simulations during early discovery (pre-synthesis) can be done using *in silico* predicted PhysChem and ADME properties
- We performed a large-scale assessments across our portfolio:  
how well can we assess Vss across compounds and species when we have IV data?
- Vss was well predicted for ECCS class 2 compounds, but class 3A/B were under-predicted

For rat IV studies: we surveyed ~575 compounds




For monkey IV studies: we surveyed ~150 compounds



# PhysChem and ADME in silico models as input parameters for PK-Sim

The simulations require these input parameters:

- Lipophilicity → multi-task (LogP / LogD) LASSO, RDKit 2D descriptors + Morgan FP (r=2, 4k bits)
- fu,p → multi-task (5 species + 7 auxiliary tasks) GNN (chemprop)
- Solubility → MELLODDY multi-task FF-NN (SparseChem), ECFP6, 32k bits
- Passive permeability → multi-task (5 end-points: MDCK-LE, PAMPA, Caco-2, MDR1) GNN (chemprop)
- Ionization → Molecular Discovery's MoKa, trained on in-house data
- Blood/plasma ratio → no models (yet!), using instead a default value per ionization class, based on an analysis of our in-house data
- Intrinsic clearance → multi-task (6 species) GNN (chemprop)







# Machine learning models to predict PhysChem/ADME/PK properties

- Machine learning (ML) models can relate molecular structures to properties  
→ Quantitative or qualitative structure-property relationship (Q)SPR

1. **Molecular features** to represent them numerically, e.g. descriptors, fingerprints, graph neural networks



2. **ML algorithm** that finds structural patterns that correlate with ADME properties



# Predicting metabolic clearance

- Multi-task learning strategy where each species was considered as a separate task<sup>[1]</sup>
- A graph neural network (GNN) was utilized for representation learning<sup>[2]</sup>
- A multi-species GNN model was generated to predict metabolic clearance<sup>[1]</sup>


~ # Compounds

Rat	Human	Mouse	Dog	Monkey	Minipig
180,000	112,000	85,000	10,000	8,000	700

[1] Rodriguez-Perez et al. 2022 Mol Pharm, 20, 1, 383-394

[2] Yang et al. 2019 JCI, 59, 3370-3388

Example: Multi-species metabolic clearance model  
(Multi-task graph neural network, MT-GNN)



# Predicting metabolic clearance

- Multi-task learning strategy where each species was considered as a separate task<sup>[1]</sup>
- A graph neural network (GNN) was utilized for representation learning<sup>[2]</sup>
- A multi-species GNN model was generated to predict metabolic clearance<sup>[1]</sup>


~ # Compounds

Rat	Human	Mouse	Dog	Monkey	Minipig
180,000	112,000	85,000	10,000	8,000	700

[1] Rodriguez-Perez et al. 2022 Mol Pharm, 20, 1, 383-394

[2] Yang et al. 2019 JCI, 59, 3370-3388

## Example: Multi-species metabolic clearance model (Multi-task graph neural network, MT-GNN)



**Especially for species with less data points, e.g. dog and monkey, results were superior with MT-GNN**




# Metabolic clearance:

## Model precision vs. Experimental reproducibility

- **Model precision:** From the compounds predicted “Low”, how many are measured “Low”?
- **Experimental reproducibility:** From the ones measured “Low”, how many were always measured “Low”?
- **With the limitation of having some ‘inconclusive’ predictions (25-32%), model’s precision is comparable to experimental reproducibility**


Species	Low CL <sub>int</sub> (≤ 100μL/min/mg)		High CL <sub>int</sub> (≥ 300μL/min/mg)	
	Model precision	Experim. reproduc.	Model precision	Experim. reproduc.
Rat	84	83	75	69
Human	85	88	75	64
Mouse	80	84	78	72
Dog	92	89	97	58
Monkey	94	87	86	71

# Outlook



# What is coming next?

- Still some plumbing to do:
  - Storing of simulations
  - Centralizing the decision trees into a snowflake-based process, which will improve the performance (currently: ~3.8s/compound for data pre-processing and ~2.5s/compound to run a single species, single dose, single arm simulation)
- Automating parameter identification
- Learning from the data! E.g.
  - Which method works best for which type of compounds (depending on ECCS class, ionization, permeability, modality, ...)
  - Investigating protein binding, transporters, clearance...
  - Assessing performance of unbound parameters
- Democratizing the process
  - Onboarding more projects
  - Enabling simulations in our design & analytics tools
























7 ■

? ⚙ ×



	Common Name ↑	Structure	HT-PBPK	Species	Matrix	Route	Vehicle	Dose	In vivo CL	Vss	Cmax	AUCinf	AUClast	%F
1	clonazepam			Rat Legacy	Plasma	i.v.	20%PG, 50(20%)Solutol, WFI	1.00	71.00	3.10		766.00	759.00	
2	etoposide			Mouse	Plasma	i.v.	PBS	10.00	8.97	0.16		31700.00	31700.00	
3	gliquidone			Rat	Blood	i.v.	NMP:PEG 200 (10:90)	0.30	6.42	0.38		1610.00	1600.00	
4	tofacitinib			Rat	Blood	i.v.	NMP:PEG200 (30:70)	1.00	53.00	1.70		1004.00	987.00	
5	tofacitinib			Mouse	Plasma	p.o.	0.5% MC/0.1% Tween	10.00			1409.00	1985.00		19.00
6	trametinib			Mouse	Plasma	p.o.	0.5% MC/0.5% Tween; suspension	10.00			635.00	13178.00		18.00
7	trametinib			Mouse	Plasma	i.v.	PEG300/D5W, 3:1; solution	2.00	3.63	1.19	3085.00	14938.00		18.00
8	vismodegib			Rat Legacy	Plasma	p.o.	10% NMP, 15% PEG300, 5 Solutol and WFI	3.00			7399.23	39911.27	32466.09	100.00

HT-PBPK

2 S

10 of 836 rows

SELECT



# Reimagining medicine, together.

## **Novartis:**

- Jan Schlender
- Sarah McFann
- Gloria Ha
- Churni Gupta
- Jimmy Kromann
- Peter Ashcroft
- Raquel Rodríguez-Pérez

## **ESQLabs:**

- Stephan Schaller
- Pavel Balazki
- Laura Villain
- Diane Lefaudeux
- Susana Proença
- Mariana Guimaraes Sa Correia