OpenFOAM® Journal

Volume 2 [Enter correct Journal Section here], Pages 1–999
ISSN: 2753-8168

OpenFOAM® Journal publication

DOI: TBD

Results with version(s): OpenFOAM® v20xx

Repository:

Abstract. This is the place for an abstract.

1. Introduction

This is the place for introduction.

1.1. Subsection. Example text:

We shall consider the specific transport property ϕ and note that its spatial and temporal variation is governed by a second-order partical differential equation (PDE), viz.

$$\frac{\partial}{\partial t}(\rho\phi) + \nabla\bullet(\rho\phi\mathbf{U}) - \Gamma_\phi\nabla^2\phi - S_\phi(\phi) = 0. \tag{1}$$

Herein, $\phi = \phi(\mathbf{x}, t)$ is an arbitrary general intensive physical quantitity, e.g., a fluid property (scalar or tensor of any rank). Thus, Eqn. 1 is often referred to as generic transport equation.

OpenFOAM® (Open Field Operation And Manipulation) is a flexible and mature C++ Class Library for Computational Continuum Mechanics (CCM) and Multiphysics. Its Object-Oriented-Programming (OOP) paradigm enables to *mimic data types and basic operations* of CCM using top-level syntax as close as possible to the conventional mathematical notation *for tensors and partial differential equations*:

```
solve
(
  fvm::ddt(rho,Phi)
  + fvm::div(phi, Phi)
  - fvm::laplacian(Gamma, Phi)
 ==
  Sphi
);
```

* Corresponding author

Received: 14 Juntember 2022, Accepted: 29 Januaril 2022, Published: 17 Decemby 2022

©2022 The Authors. This work is an open access article under the CC BY-SA 4.0 license

Table 1. Finite Volume Notation

implicit differential operators	
rate of change	$\left[\!\left[\frac{\partial[\rho\phi]}{\partial t}\right]\!\right]$
convection term	$\left[\!\left[\nabla\bullet\left(F[\phi]_{f(F,S,\gamma)}\right)\right]\!\right]$
diffusion term	$[\![\nabla\bullet(\Gamma\nabla[\phi])]\!]$
linear part of source term	$[\![S_p\,[\phi]]\!]$
explicit differential operators	
temporal term	$\frac{\partial\rho\phi}{\partial t}$
divergence term	$\nabla\bullet(\rho\mathbf{U}\phi_{f(\rho\mathbf{U},S,\gamma)})$
laplacian term	$\nabla\bullet(\Gamma\nabla\phi)$
constant part of source term	S_u




Figure 1. Examplary figure

Beside providing OpenFOAM® code itself, spatial and temporal discretisation of Eqn. 1 can be also described in a precise and concise manner using the finite-volume notation [1] - see Tab. 1.

2. Theoretical backgroud

Text in this section. Here is an examplary figure (Fig. 1).

3. Conclusion

This is a conclusion.

References

[1] H. G. Weller, G. Tabor, H. Jasak, and C. Fureby, "A tensorial approach to computational continuum mechanics using object orientated techniques," *Computers in Physics*, vol. 12, pp. 620–631, 1998.