OpenFOAM® Journal

Volume 2 [Enter correct Journal Section here], Pages 1–999
ISSN: 2753-8168

OpenFOAM® Journal publication

Maria Jersey[1,*], John Smith[2], and Philip Murphy[2]

[1]Address1
Email address: Emailaddress1

[2]Address2

DOI: TBD
Results with version(s): OpenFOAM® v20xx
Repository: https://github.com/xxx

Abstract. This is the place for an abstract.

1. Introduction

This is the place for introduction.

1.1. Subsection.

Example text:

We shall consider the specific transport property ϕ and note that its spatial and temporal variation is governed by a second-order partical differential equation (PDE), viz.

$$\frac{\partial}{\partial t}(\rho\phi) + \nabla\bullet(\rho\phi\mathbf{U}) - \Gamma_\phi\nabla^2\phi - S_\phi(\phi) = 0. \tag{1}$$

Herein, $\phi = \phi(\mathbf{x}, t)$ is an arbitrary general intensive physical quantitity, e.g., a fluid property (scalar or tensor of any rank). Thus, Eqn. 1 is often referred to as generic transport equation.

OpenFOAM® (Open Field Operation And Manipulation) is a flexible and mature C++ Class Library for Computational Continuum Mechanics (CCM) and Multiphysics. Its Object-Oriented-Programming (OOP) paradigm enables to *mimic data types and basic operations* of CCM using top-level syntax as close as possible to the conventional mathematical notation *for tensors and partial differential equations*:

```
solve
(
  fvm::ddt(rho,Phi)
  + fvm::div(phi, Phi)
  - fvm::laplacian(Gamma, Phi)
 ==
  Sphi
);
```

Beside providing OpenFOAM® code itself, spatial and temporal discretisation of Eqn. 1 can be also described in a precise and concise manner using the finite-volume notation [1] - see Tab. 1.

2. Theoretical backgroud

Text in this section. Here is an examplary figure (Fig. 1).

3. Conclusion

This is a conclusion.

* Corresponding author

Received: 14 Juntember 2022, Accepted: 29 Januaril 2022, Published: 17 Decemby 2022

©2022 The Authors. This work is an open access article under the CC BY-SA 4.0 license

Table 1. Finite Volume Notation

implicit differential operators	
rate of change	$\left[\!\left[\frac{\partial[\rho\phi]}{\partial t}\right]\!\right]$
convection term	$\left[\!\left[\nabla\bullet\left(F[\phi]_{f(F,S,\gamma)}\right)\right]\!\right]$
diffusion term	$[\![\nabla\bullet(\Gamma\nabla[\phi])]\!]$
linear part of source term	$[\![S_p\,[\phi]]\!]$
explicit differential operators	
temporal term	$\frac{\partial\rho\phi}{\partial t}$
divergence term	$\nabla\bullet(\rho\mathbf{U}\phi_{f(\rho\mathbf{U},S,\gamma)})$
laplacian term	$\nabla\bullet(\Gamma\nabla\phi)$
constant part of source term	S_u




Figure 1. Examplary figure

Acknowledgements

This is an example acknowledgements section and should only be included !after! the review process and before publication.

The OpenFOAM Journal requires the contributions of all authors to be explicitly stated As noted below, please see CRediT taxonomy for the term explanation. Replace the example authors initials (F.A., S.A) below as appropriate

Please turn to http://img.mdpi.org/data/contributor-role-instruction.pdf (CRediT taxonomy) for the term explanation.

Author Contributions: Conceptualisation, J.S.; methodology, J.S.; software, J.S; validation, J.S. and P.M.; formal analysis, J.S. and P.M.; investigation, J.S.; resources, P.M.; data curation, P.M.; writing—original draft preparation, J.S.; writing—review and editing, J.S. and P.M.; visualisation, J.S.; supervision, P.M.; project administration, P.M.; funding acquisition, P.M. All authors have read and agreed to the published version of the manuscript.

References

[1] H. G. Weller, G. Tabor, H. Jasak, and C. Fureby, "A tensorial approach to computational continuum mechanics using object orientated techniques," *Computers in Physics*, vol. 12, pp. 620–631, 1998.