State Cancer Profiles Demo

Dave Stinchcomb, Westat
Zaria Tatalovich, NCI

November 18, 2016
FCSM GIG Workshop

Cancer Statistics and other geographically- based data

- Incidence - how many people get cancer in a given period

- Mortality - how many people died from cancer in a given period

- Prevalence - how many people have ever had a cancer and are still alive

- Survival – how many people stay alive within a certain period after diagnosis

Screening
Risk factors
Demographics
Cancer Knowledge

Cancer Control

- **Goal: to reduce the impact of cancer on people's lives**

- **The cancer control continuum:**

THE CANCER CONTROL CONTINUUM				
Focus				
PREVENTION	**DETECTION**	**DIAGNOSIS**	**TREATMENT**	**SURVIVORSHIP**
Tobacco control	Pap/HPV testing	Shared and informed decision making	Health care delivery and outcomes research	Coping
Diet	Mammography			Health promotion for survivors
Physical activity	Fecal occult blood test			
Sun protection	Colonoscopy			
HPV vaccine	Lung cancer screening			
Limited alcohol use				
Chemoprevention				

- **Cancer control planning**
  - Interventions to reduce the impact of cancer

Quick Profiles: New Mexico

▶ Choose a Different State:

State: [New Mexico ▼]   **View Profile**

|				
👥 Demographics	❤ Screening & Risk Factors	〽 Incidence	◉ Prevalence	◔ Mortality

Age-Adjusted Incidence Rates by Cancer Site (2008-2012)	New Mexico Rate	USA Rate	Map	Table	Historical Trends Graph	5-Year Rate Change Graph
All Cancer Sites	388.1	453.8				
Bladder	14.9	20.8				
Brain & ONS	5.5	6.6				
Breast	112.1	123.0				
Breast (in situ)	20.1	31.0				
Cervix	7.5	7.7				
Childhood (Ages <15, All Sites)	13.7	16.0				
Childhood (Ages <20, All Sites)	15.0	17.4				
Colon & Rectum	35.8	41.9				
Esophagus	3.8	4.7				
Kidney & Renal Pelvis	13.7	16.0				
Leukemia	12.6	13.2				

Incidence Rates[†] for United States, 2008 - 2012
Breast
All Races (includes Hispanic), Female, All Ages

Age-Adjusted

Annual Incidence Rate

(Cases per 100,000)

Quantile Interval

- 128.6 to 141.7
- 125.3 to 128.6
- 122.2 to 125.3
- 118.9 to 122.2
- 107.9 to 118.9

Data Not Available ◇

US (SEER + NPCR)
Rate (95% C.I.)
123.0 (122.8 - 123.2)

DC

Notes:
Created by statecancerprofiles.cancer.gov on 10/15/2015 9:46 am.
Data for the United States does not include data from Nevada.
State Cancer Registries may provide more current or more local data.
Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries (for more information).
† Incidence rates (cases per 100,000 population per year) are age-adjusted to the 2000 US standard population (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The 1969-2013 US Population Data File is used for SEER and NPCR incidence rates.
◇ Data not available for this combination of geography, statistic, age and race/ethnicity.

Incidence Rates[†] for New Mexico, 2008 - 2012
Breast
All Races (includes Hispanic), Female, All Ages

Age-Adjusted

Annual Incidence Rate

(Cases per 100,000)

Quantile Interval

- 124.2 to 217.1
- 103.2 to 124.2
- 96.9 to 103.2
- 90.2 to 96.9
- 66.7 to 90.2

⬚ Suppressed */**

US (SEER + NPCR)
Rate (95% C.I.)
123.0 (122.8 - 123.2)

New Mexico
Rate (95% C.I.)
112.1 (109.3 - 114.9)

Notes:

Created by statecancerprofiles.cancer.gov on 10/08/2015 3:17 pm.

Data for the United States does not include data from Nevada.

State Cancer Registries may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries (for more information).

† Incidence rates (cases per 100,000 population per year) are age-adjusted to the 2000 US standard population (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The 1969-2013 US Population Data File is used for SEER and NPCR incidence rates.

* Data have been suppressed to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-race category.

** Data have been suppressed for states with a population below 50,000 per sex combination for American Indian/Alaska Native or Asian/Pacific Islanders because of concerns regarding the relatively small size of these populations in some states.

Interactive Maps

	193.4 to 209.5
	184.6 to 193.3
	177.7 to 184.5
	170.8 to 177.6
	164.9 to 170.7
	131.3 to 164.8

United States
Rate (95% C.I.)
176.4 (176.2 - 176.6)

Maps of mortality data can be created at the state and county level.

Tables

California	272,100	277,800
Florida	217,200	224,600
Texas	175,100	180,100
New York	155,200	156,000
Pennsylvania	122,000	124,200
Illinois	97,600	99,000
Ohio	96,000	97,500
Michigan	88,800	90,200
New Jersey	76,800	77,200
North Carolina	72,100	74,000

Tables of mortality data can be created at the state and county level.

Historical Trends

Compare trends in cancer mortality and incidence by user selectable criteria.

5-Year Rate Changes

Rate changes in cancer mortality or incidence for all major cancer sites.

Rate/Trend Comparison by Cancer

Set higher priority for cancer control when rates are high or rising.

Rate/Trend Comparison by Area

Set higher priority for cancer control when rates are high or rising.

Data Topics Across the Cancer Control Continuum

Cancer statistics, charts, and maps by data topic across the cancer control continuum.

- Demographics
- Screening & Risk Factors
- Cancer Knowledge
- Incidence
- Prevalence
- Mortality

Interactive Maps at the state and county level

Interactive Maps at the national level

Interactive Maps

DATA OPTIONS

Area: --- choose area ---

Data Group: --- choose data group ---

Cancer: --- choose cancer ---

Statistic: --- choose data type ---

Race/Ethnicity: --- choose race/ethnicity ---

Sex: --- choose sex ---

Age: --- choose age group ---

Year(s): --- choose year range ---

MAP OPTIONS

Interval Type

Color Scheme

Interval Type: Quantiles (rankings)

Number of Intervals: 5

Color Scheme: Red-Yellow-Blue (Div)

Generate Map

Data Table   Export Data   Interpret   Save Map Image

Tip: Hover on a state to see its rate (and 95% C.I.). Click a state to show its counties.

Death Rates for United States
All Cancer Sites, 2009 - 2013
All Races (includes Hispanic), Both Sexes, All Ages

Age-Adjusted Annual Death Rate (Deaths per 100,000)

Quantile Interval

- 128.1 to 159.0
- 159.0 to 168.2
- 168.2 to 171.6
- 171.6 to 182.3
- 182.3 to 202.6

DC

United States
Rate (95% C.I.)
168.5 (168.3 - 168.7)

Healthy People 2020
Goal C-1
160.6

Notes:
Created by statecancerprofiles.cancer.gov on 04/19/2016 8:38 am.
State Cancer Registries may provide more current or more local data.
Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries (for more information).
Source: Death data provided by the National Vital Statistics System public use data file. Death rates calculated by the National Cancer Institute using SEER*Stat. Death rates (deaths per 100,000 population per year) are age-adjusted to the 2000 US standard population (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). The Healthy People 2020 goals are based on rates adjusted using different methods but the differences should be minimal. Population counts for denominators are based on the Census 1969-2014 US Population Data File as modified by NCI.
Healthy People 2020 Goal C-1 : Reduce the overall cancer death rate to 160.6.
Healthy People 2020 Objectives provided by the Centers for Disease Control and Prevention.

Interactive Maps

DATA OPTIONS

Area:	--- choose area ---
Data Group:	--- choose data group ---
Cancer:	--- choose cancer ---
Statistic:	--- choose data type ---
Race/Ethnicity:	--- choose race/ethnicity ---
Sex:	--- choose sex ---
Age:	--- choose age group ---
Year(s):	--- choose year range ---

About this Map
Quick Reference Guide
Tutorial
Data Use Restrictions

MAP OPTIONS

Interval Type:	Quantiles (rankings)
Number of Intervals:	5
Color Scheme:	Red-Yellow-Blue (Div)

Interval Type
Color Scheme

Generate Map

DATA OPTIONS

Area:

Data Group:

Cancer:

Statistic:

Race/Ethnicity:

Sex:

Age:

Year(s):

--- choose area ---
US by state
US by county
Alabama counties
Alaska counties
Arizona counties
Arkansas counties
California counties
Colorado counties
Connecticut counties
Delaware counties
District of Columbia
Florida counties
Georgia counties
Hawaii counties
Idaho counties
Illinois counties
Indiana counties
Iowa counties
Kansas counties
Kentucky counties
Louisiana counties
Maine counties
Maryland counties
Massachusetts counties
Michigan counties
Minnesota counties
Mississippi counties
Missouri counties
Montana counties

--- choose data group ---
Cancer Rates
Demographic Data
Screening & Risk Factors

Data Group:	Cancer Rates ∨
Cancer:	--- choose cancer --- ∨
Statistic:	--- choose data type --- ∨

--- choose cancer ---
All Cancer Sites
Bladder
Brain & ONS
Breast (Female)
Breast (Female in situ)
Cervix
Childhood (Ages <15, All Sites)
Childhood (Ages <20, All Sites)
Colon & Rectum
Esophagus
Kidney & Renal Pelvis
Leukemia
Liver & Bile Duct
Lung & Bronchus
Melanoma of the Skin
Non-Hodgkin Lymphoma
Oral Cavity & Pharynx
Ovary
Pancreas
Prostate
Stomach
Thyroid
Uterus

--- choose data type ---
Incidence
Mortality

Data Group:	Demographic Data
Topic:	--- choose demographic topic ---
Variable:	--- choose data type ---

Topic dropdown:
- --- choose demographic topic ---
- Crowding
- Education
- Income
- Mobility
- Insurance
- Non-English Language
- Population
- Poverty
- Workforce

Education
- --- choose demographic variable ---
- Less than 9th grade
- Less than HS
- At least bachelors degree

Income
- --- choose demographic variable ---
- Median family income ($)
- Median household income ($)

Poverty
- --- choose demographic variable ---
- Families below poverty
- Persons below poverty
- Persons <150% of poverty

Population
- --- choose demographic variable ---
- Ages under 18
- Ages 40 and over
- Ages 50 and over
- Ages 65 and over
- Foreign born
- Black
- American Indian/Alaska Native
- Asian/Pacific Islander
- Hispanic
- White

Mobility
- --- choose demographic variable ---
- Moved from outside US (in last 5 yrs)
- Moved, different state (in last 5 yrs)
- Moved, different county, same state (in last 5 yrs)
- Moved, same county (in last 5 yrs)
- Haven't moved (in last 5 yrs)

| Area: | US by state | ▼ |

| Data Group: | Screening & Risk Factors | ▼ |

| Topic: | --- choose topic --- | ▼ |

| Screening/Risk Factor: | --- choose screening or risk factor --- | ▼ |

| Data Type: | Direct Estimates | ▼ |

--- choose topic ---
Colorectal Screening
Diet & Exercise
Smoking
Women's Health

Diet & Exercise

--- choose screening or risk factor ---
BMI is Healthy, Ages 20+
BMI is Obese, Ages 20+
BMI is Obese, High School Survey
BMI is Overweight, High School Survey
Consumed 1 or More Fruits per Day
Consumed 1 or More Vegetables per Day
No Leisure Time Physical Activity, All Ages

Smoking

--- choose screening or risk factor ---
Smokers (Current); Ages 18+
Smokers (Ever); Ages 18+
Smokers (Stopped for 1 Day or Longer)
Smoking Not Allowed at Work (All People)
Smoking Not Allowed at Work (Current Smokers)
Smoking Not Allowed at Work (Former/Never Smokers)
Smoking Not Allowed in Home (All People)
Smoking Not Allowed in Home (Current Smokers)
Smoking Not Allowed in Home (Former/Never Smokers)

Women's Health

--- choose screening or risk factor ---
Mammogram in Past 2 Years, Ages 40+
Mammogram in Past 2 Years, Ages 50-74
Pap Smear in Past 3 Years, No Hysterectomy, Ages 18+
Pap Smear in Past 3 Years, No Hysterectomy, Ages 21-65
Percent who received 3+ doses of HPV vaccine, Ages 13-15
Percent who received 3+ doses of HPV vaccine, Ages 13-17

Colorectal Screening ▼

--- choose screening or risk factor ---
Ever Had Colorectal Endoscopy (sigmoidoscopy or colonoscopy), Ages 50+
FOBT in Last Year and/or Flex Sig in Past 5 Years and FOBT in Last 3 Years and/or Colonoscopy in Past 10 Years
Home-Based Fecal Occult Blood Test (FOBT) in Past Two Years
Home-based FOBT in the past two years or ever had a colorectal endoscopy

Area:	US by county	∨
Data Group:	Screening & Risk Factors	∨
Topic:	--- choose topic ---	∨
Screening/Risk Factor:	--- choose screening or risk factor ---	∨
Data Type:	County Level Modeled Estimates	∨

--- choose topic ---
Colorectal Screening
Smoking
Women's Health

Colorectal Screening ∨

--- choose screening or risk factor ---
Ever Had Colorectal Endoscopy (sigmoidoscopy or colonoscopy)
Home-based FOBT in the past two years or ever had a colorectal endoscopy
Home-based Fecal occult blood test (FOBT) in Past Two Years

Smoking

--- choose screening or risk factor ---
Smokers (Current), Ages 18+
Smokers (Ever); Ages 18+

Women's Health

--- choose screening or risk factor ---
Mammogram in Past 2 Years, Ages 40+
Pap Smear in Past 3 Years, No Hysterectomy, Ages 18+

Data Topics Across the Cancer Control Continuum

Cancer statistics, charts, and maps by data topic across the cancer control continuum.

- Demographics
- Screening & Risk Factors
- Cancer Knowledge
- Incidence
- Prevalence
- Mortality

Interactive Maps at the state and county level

Interactive Maps at the national level

Cancer Knowledge Map

DATA OPTIONS

Topic: Lung Cancer

Question: Does smoking increase chances of cancer? (2003)

Does smoking increase chances of cancer?

HINTS 2003 CK-13a

Responded "Yes"

High : 96%

Low : 79%

Created by statecancerprofiles.cancer.gov on 11/16/2016 1:30 pm.
For more information, visit the HINTS web site or the data used to generate the map.

A cancer control planning example:
a smoking cessation program in Kentucky

Lung cancer incidence

A cancer control planning example:
a smoking cessation program in Kentucky

Lung cancer mortality

Age-Adjusted Annual Death Rate (Deaths per 100,000)

Quantile Interval

- 47.8 to 64.2
- 64.2 to 70.5
- 70.5 to 79.5
- 79.5 to 88.9
- 88.9 to 125.6
- Suppressed *

United States Rate (95% C.I.) 46.0 (45.9 - 46.1)

Kentucky Rate (95% C.I.) 70.0 (68.9 - 71.0)

Healthy People 2020 Goal C-2 45.5

A cancer control planning example: a smoking cessation program in Kentucky

Current smoking rates

Current Smokers
(Percent of Respondents)

Quantile Interval

- 18.5 to 26.9
- 26.9 to 29.3
- 29.3 to 31.1
- 31.1 to 33.6
- 33.6 to 38.6

Data Not Available ◇

Kentucky
Rate (95% C.I.)
Data not available.

Healthy People 2020
Goal TU-1.1
12.0%

A cancer control planning example:
a smoking cessation program in Kentucky

Now It's Your Turn!

- Construct a series of 2-3 maps to support a cancer control initiative
- You pick the state
- You pick the type of cancer
- A few possibilities:
    - Education programs in areas with high cancer rates
    - Cancer screening programs
    - Areas with vulnerable populations (poverty, minorities)
- Watch out for data suppression:
    - Counties with low populations
    - Rare cancers

Questions?

DavidStinchcomb@westat.com
TatalovichZP@mail.nih.gov