



Original Articles

Effect of Hyperinsulinemia on Urea Pool Size and Substrate Oxidation Rates

Luc Tappy, Oliver E Owen **and** Guenther Boden

Author Affiliations

Address correspondence to Guenther Boden, MD, Temple University Health Sciences Center, 3401 North Broad Street, Philadelphia, PA 19140.

Diabetes 1988 Sep; 37(9): 1212-1216.

<https://doi.org/10.2337/diab.37.9.1212>



Previous

Next

Article

Info & Metrics

PDF

Abstract

Recently, indirect calorimetry has frequently been used together with hyperinsulinemic clamps. With few exceptions, however, no attention was paid in these studies to the possible effects of hyperinsulinemia on urea nitrogen (*M*) pool size and the consequences of such changes on the calculated rates of protein, lipid, and carbohydrate (CHO) oxidation. We have determined the effects of euglycemic-hyperinsulinemic clamps on urea N pool size, urinary N excretion, and rates of protein, lipid, and CHO oxidation (measured by indirect calorimetry) in six normal men. Insulin infusion ($1 \text{ mU} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) increased peripheral venous insulin concentration from 7 ± 1.2 (mean \pm SE) to $51 \pm 4 \text{ } \mu\text{U/ml}$. Glucose concentration was clamped at $84 \pm 1.1 \text{ mg/dl}$. Between 0 (preclamp) and 360 min (end of clamp), blood urea N concentration decreased from 17.2 ± 1.1 to $11 \pm 0.8 \text{ mg/dl}$ ($P < .001$), and the urea N pool decreased from 604 ± 41 to $388 \pm 30 \text{ mmol}$ ($P < .001$). The urea N

production rate decreased from 461 ± 91 (preclamp) to 91 ± 63 $\mu\text{mol}/\text{min}$ during the last 4 h of the clamp ($P < .05$). Urinary N excretion remained unchanged (705 ± 113 vs. 905 ± 125 $\mu\text{mol}/\text{min}$, NS). Correction of urinary N excretion for insulin-induced reductions in the urea N pool resulted in the following changes in substrate oxidation rates (calculated for the last 4 h of the clamp). Protein oxidation rate decreased by $72 \pm 8\%$ (from 82.8 ± 10.9 to 25.7 ± 9 mg/min); lipid oxidation rate increased by $66 \pm 16\%$ (from 43.6 ± 11.9 to 61.7 ± 12.1 mg/min); and CHO oxidation rate increased by $12 \pm 2\%$ (from 167.8 ± 20.4 to 187.7 ± 21.6 mg/min). We conclude that euglycemic hyperinsulinemia decreases the urea N pool. If uncorrected, this will result in substantial overestimation of protein oxidation and underestimation of lipid and CHO oxidation rates.

Received December 14, 1987.

Revision received April 4, 1988.

Accepted April 4, 1988.

Copyright © 1988 by the American Diabetes Association

Log in using your username and password

<input type="text" value="Username"/>	<input type="text" value="Password"/>
<input type="button" value="Log in"/>	

[Forgot](#) your user name or password?

[Pay Per Article](#) - You may access this article (from the computer you are currently using) for 1 day for US\$35.00

[Regain Access](#) - You can regain access to a recent Pay per Article purchase if your access period has not yet expired.

We recommend

Effects of insulin on cholesterol synthesis in type II diabetes patients.

A Scoppola et al., *Diabetes Care*

Effects of physical training and diet therapy on carbohydrate metabolism in patients with glucose intolerance and non-insulin-dependent diabetes mellitus.

C Bogardus et al., *Diabetes*

Similar insulin sensitivity in NIDDM patients with normo- and microalbuminuria.

S Nielsen et al., *Diabetes Care*

Link Between Insulin Sensitivity and Albuminuria
PracticeUpdate

Morning Hyperinsulinemia Enhances Glucose Uptake and Glycogen Storage
PracticeUpdate

Insulin resistance of protein anabolism accompanies that of glucose metabolism in lean, glucose-tolerant offspring of persons with type 2 diabetes
Sergio A Burgos et al., *BMJ Open Diab Res Care*