Weather, daylight, snow and bicycle traffic on rue de Terrebonne

Five count series, August 2024 to November 2026

Terrebonne-open-sources, version 2026-11-10-10-15-33. Author: Rick Hoge, PhD (see the repository's « About the author »). Scripts, tables and figures: `https://github.com/OpenMobilityData/Terrebonne-open-sources/tree/main/2026-weather-and-bike-traffic`.

DRAFT for the author's review, November 10, 2026. Not yet checked line by line; figures may change.

Lay summary

How much does the weather change cycling on rue de Terrebonne? This note uses every count available for the street: citizen-operated camera sensors at King Edward and Draper, the borough's counter at Kensington, and trips at the BIXI station at Grand Boulevard, with City counters elsewhere in Montréal for comparison. Weather is Environment Canada's record at Trudeau airport; the counts run from August 2024 to November 2026.

Temperature matters most. Each degree warmer brings about 5 to 6 per cent more bicycles, up to about 20 °C, where the numbers stop rising. A day at 0 °C sees 30 to 34 per cent of the bicycles of the same day at 20 °C; a day at −10 °C, 17 to 20 per cent. The decline is steady: there is no temperature below which cycling stops.

Rain and snow come next. A day with 2 to 10 mm of rain has 14 to 22 per cent fewer bicycles than a dry day at the same temperature; snow on the ground goes with 31 to 47 per cent fewer. Wind chill describes the counts about as well as temperature and wind do. Darkness matters too: at the same evening hour and temperature, the borough's counter records about 27 per cent fewer bicycles after sunset.

Some questions the counts cannot settle. Cold, snow and short days arrive together, so much of the winter drop could be either weather or season. On days when the City loaded snow, the counts resemble those of other winter days, but such days are too few to be sure.

The cameras count only in daylight (read here from 8 a.m. to 4 p.m.); the borough's counter runs day and night. Their numbers are never compared directly, only their percentage changes. The note makes no recommendation.

Abstract

This note measures how the number of bicycles counted on rue de Terrebonne varies with the weather, with daylight and with snow, using every count series available for the street: two citizen-operated Telraam camera sensors (King Edward, 691 valid days from August 16, 2024; Draper, 509 days), the borough's loop counter at Kensington (284 days, eastbound, hourly release of June 4, 2026), trips at the BIXI station on the street, and, as a reference, twelve City counters elsewhere in Montréal. Weather is Environment and Climate Change Canada's record at Montréal-Trudeau. Each series is modelled separately by negative binomial regression, so that every effect is a percentage and no levels are compared between series; the Telraam sensors, which count only in daylight, are analysed in a fixed window (08:00–16:00) that is daylight all year. The note states what was measured, with intervals, and what the data cannot show. It makes no recommendation.

What the analysis finds

1. **Temperature.** On the three Terrebonne count series, each additional degree of daytime temperature goes with 5 to 6 % more bicycles, up to roughly 20 °C; above that the count stops rising. Holding rain, snow, wind, day of the week and time of year fixed, a day at 0 °C has 30 to 34 % of the count of the same day at 20 °C, and a day at −10 °C has 17 to 20 %. The count falls steadily with the cold; it does not fall to zero.
2. **No threshold in the cold.** In percentage terms the slope below 0 °C is not distinguishable from the slope between 0 °C and 20 °C on any Terrebonne series. The change of slope that the earlier analyses placed near −1 or −5 °C belongs to a straight-line model of the counts themselves; the only change of slope the data support is the levelling-off in warm weather, and its position is poorly determined (intervals of ten degrees or more).
3. **Wind chill.** Replacing air temperature and wind by the wind-chill index fits about as well (the two are correlated at 0.996); the data cannot separate them.
4. **Rain and snow.** At a given temperature and time of year, a day with 2 to 10 mm of rain has 14 to 22 % fewer bicycles on the Terrebonne series, a day with more than 10 mm 25 to 46 %, and a day with at least 1 cm of snow on the ground 31 to 47 %. Snow on the ground is the largest single weather effect after temperature.
5. **Season or weather?** The two cannot be fully separated: 65 to 81 % of the day-to-day variation on the Terrebonne series can be attributed to either. Weather effects estimated only from the variation inside each calendar month are nonetheless close to those estimated over the whole record.
6. **Darkness.** On the loop counter, the same clock hour in the evening carries 27 % (19 to 34) fewer bicycles when the sun has set than when it is up, at the same temperature and in the same month.
7. **Snow loading.** On the days the City loaded snow in the sector, the counts are not distinguishable from other winter days once temperature and snow are accounted for; the intervals are wide (36 to 44 loading days per series).
8. **Agreement.** The two camera sensors, the loop counter, the BIXI station on the street and the City's counters elsewhere respond to weather in the same direction and with similar size, although they count different things.




Figure 1: Daytime temperature and the daily count of each series. Each panel has its own scale; the series are not compared in level.

Contents

1 Questions **5**

2 Data **5**

2.1 The count series 5

2.2 Weather, daylight, snow 6

3 Method **6**

4 Temperature **7**

5 Wind chill **9**

6 Rain, snow and wind **9**

7 Day length, season and darkness **10**

8 Snowfall, the days after, and snow loading **12**

9 Agreement between the series **13**

10 Year to year, after weather **13**

11 How well the models predict **13**

12 Limits **14**

A Valid days by month **15**

B Model comparison **15**

C Where the slope changes: deviance profile **16**

D The earlier analyses and what was kept **16**

E Sources **17**

1 Questions

1. How does the daily count of bicycles vary with temperature, and is there a temperature at which the response changes?
2. Does wind chill describe the response better than air temperature?
3. What do rain, snowfall, snow on the ground and wind each add at a given temperature?
4. How much of the seasonal swing can be attributed to weather, and how much to day length and the calendar?
5. Does a snowfall depress the counts on the following days, and do the days on which the City loads snow differ from other winter days?
6. Do the series agree?
7. After weather is accounted for, does the count differ from one year to the next?

2 Data

2.1 The count series

Series	What is counted	Days used in the models	Days	Mean a day
King Edward	Bicycles, both directions, 08:00–16:00 (Telraam S2 camera sensor, west of Cavendish)	August 16, 2024 to November 8, 2026	691	163
Draper	The same, east of Cavendish	September 25, 2024 to October 21, 2026	509	198
Kensington loop	Bicycles eastbound, 24 h (the borough's Eco-Counter loop)	July 27, 2025 to May 10, 2026	284	157
BIXI station	Trips starting or ending at the station « Grand Boulevard / de Terrebonne », April 15 to November 15	August 2, 2024 to August 31, 2026	441	63
City counters	Bicycles, 24 h, sum of twelve City loop counters elsewhere in Montréal	January 1, 2025 to November 6, 2026	661	14,821

The means are given to describe each series, not to compare them: a camera sensor counting eight daytime hours in both directions, one direction of a loop counting 24 hours, and trips at a bike-share dock are different quantities.

Telraam sensors (King Edward, Draper). Installed by Terrebonne residents in August and September 2024; the author was tasked with the statistical analysis of their data. Hourly counts from the Telraam API (pull of November 9, 2026). A Telraam sensor counts only in daylight. Its daily total therefore covers about nine hours in December and sixteen in June, and would rise and fall with the length of the day even if nobody changed their habits (Figure 6, left). To remove that artefact every analysis here uses the count in a *fixed clock window, 08:00 to 16:00*, which is daylight on every day of the year at Montréal's latitude (the latest sunrise is about 07:35 and the earliest sunset about 16:10). The price is that the evening peak is left out. A day is valid when the sensor reported at least 8 hours with at least 6 hours of summed uptime, and all eight window hours with an uptime of at least 0.5. Invalid days are left out and nothing is interpolated. King Edward has no valid day in February 2026; Draper has none from February to April 2026 (Appendix A). Whether the API's hourly counts are already scaled for uptime could not be confirmed from Telraam's documentation; they are used as delivered.

Kensington loop. The borough's Eco-Counter (inductive loops, 24 hours, by direction). The hourly workbook released on June 4, 2026 under access request 2026-300 runs from July 26, 2025 to May 11, 2026. The westbound channel under-counts from about March 1, 2026, so the models use the eastbound channel alone. The daily table released on October 9, 2026 under request 2026-581 serves to identify the days set aside (May 29–31, 2026, zeros; June 7–9, 2026). One site, less than one full year, one direction.

BIXI station. The station « Grand Boulevard / de Terrebonne » stands 6 m from the lane. BIXI Montréal's open trip files give each trip's start and end station and time; the series is the number of trips starting or ending there each day. A trip is not a bicycle passing a point, and a day without a trip is absent from the files rather than recorded as zero, so the models use the regular season only (April 15 to November 15). The station recorded trips in the winter of 2025–26 (3 to 6 a day on the days with trips) and none in the winter of 2024–25.

City counters. Twelve of the City's permanent loop counters, none on Terrebonne, kept because they report on at least 97 % of days from January 1, 2025 to November 6, 2026; the series is their sum on the days when all twelve report. It is a reference for how bicycle traffic elsewhere in Montréal responds to the same weather.

2.2 Weather, daylight, snow

Environment and Climate Change Canada, station Montréal-Trudeau (climate ID 7025251), about 9 km from the street. From the daily file: rain (mm), snowfall (cm), snow on the ground (cm). From the hourly file: temperature and wind, averaged over 08:00–16:00 local time (« daytime »), and the wind-chill index computed from them by Environment Canada's formula. The hourly file carries no precipitation amount, so rain and snowfall are 24-hour totals: rain that fell at night counts as a rainy day. Sunrise and sunset are computed for the street's position (NOAA solar equations). Snow loading is the City's record of loads by day for sector CDN-301, which contains the lane from its west end to Royal (open data, snow-removal transactions).

3 Method

One model per series, on a percentage scale. Daily counts are modelled by negative binomial regression with a log link. Counts are non-negative, their spread grows with their level, and a weather effect is plausibly proportional (rain removes a share of the riders, not a fixed number); a straight-line model of the counts themselves, as in the earlier analyses, forces a rainy day to remove as many bicycles in January as in June. On the log scale every coefficient is a percentage change, which is also what allows the series to be set side by side without comparing their levels.

Terms. Day type (weekday, Saturday, Sunday or public holiday); December 24 to January 2; the counting year (from August 15); two annual harmonics for the time of year; daytime temperature; rain of the day in classes (none, under 2 mm, 2 to 10 mm, over 10 mm); snowfall of the day in classes (none, under 2 cm, 2 cm or more); snow on the ground (1 cm or more); daytime wind; snowfall of the day before. A class with fewer than five days is merged with its neighbour.

Temperature. Two forms are fitted: a natural cubic spline (five degrees of freedom), which lets the data draw the curve, and a line with one change of slope at a temperature that is itself estimated, by profiling the deviance over a grid from -8 to 26 °C. The second is the form of the earlier analyses, here on the percentage scale and with the two segments joined.

Uncertainty. Consecutive days resemble each other, so standard errors that assume independent days are too small. The intervals reported are robust to heteroskedasticity and autocorrelation (Newey–West, seven lags); the interval for the change of slope comes from a moving-block bootstrap (blocks

of 14 consecutive days, 300 replicates). The dispersion parameter is set so that the Pearson statistic equals its degrees of freedom.

Season against weather. Temperature, day length and the calendar move together over the year. Three devices are used: the deviance accounted for by calendar-and-season terms alone, by weather terms alone and by both; a model in which the season terms are replaced by one constant per calendar month, so that weather effects are estimated only from days of the same month; and, for the loop, an hourly model in which the same clock hour is sometimes light and sometimes dark.

What the models cannot show. They describe association in the record, not what any person would do. They cannot separate temperature from other things that change with the season (school terms, holidays abroad, habits). Weather is measured at the airport. The loop's record covers one winter.

4 Temperature



Figure 2: Daily count against daytime temperature, days marked by precipitation. Black line: median of the days without precipitation, by 2.5-degree band.

Figure 2 shows the raw relation. Figure 3 shows the fitted response with everything else held fixed, each series relative to its own level at 20 °C.




Figure 3: Fitted response to temperature (spline model), relative to 20 °C.

Daytime temperature (°C)	King Edward	Draper	Kensington loop	BIXI station	City counters
−15	13	16	17	––	28
−10	17	20	19	––	29
−5	22	26	24	––	33
0	30	34	34	––	41
5	41	47	51	69	55
10	59	66	70	76	74
15	81	86	88	86	91
20	100	100	100	100	100
25	107	102	105	107	99

Count relative to the same day at 20 °C (= 100), spline model. BIXI: regular season only.

Slopes. With temperature as a line with one change of slope:

Series	Change of slope, °C (bootstrap 95 %)	Below it: % per degree	Above it: % per degree	Below it, within calendar months
King Edward	20.0 (10 to 26)	+6.1 (+4.9 to +7.4)	+0.4 (−1.6 to +2.4)	+6.6 (+5.3 to +8.0)
Draper	24.5 (8 to 25)	+5.0 (+4.1 to +5.8)	−7.9 (−10.2 to −5.5)	+5.0 (+4.1 to +5.9)
Kensington loop	18.0 (-8 to 26)	+5.6 (+4.5 to +6.8)	0.0 (−2.1 to +2.2)	+6.3 (+5.1 to +7.4)
BIXI station	23.5 (1 to 26)	+2.7 (+1.9 to +3.4)	−2.2 (−5.5 to +1.2)	+3.2 (+2.5 to +3.9)
City counters	20.5 (-4 to 24)	+3.7 (+3.1 to +4.4)	−1.7 (−3.0 to −0.4)	+4.4 (+3.8 to +5.0)

On the three Terrebonne series each degree below the change of slope goes with 5 to 6 % more bicycles, and the estimate is the same when only days of the same calendar month are compared (last column). The BIXI station and the City counters show smaller slopes (2.7 and 3.7 %). Above the change of slope the count is flat or falls slightly.

Where the slope changes. The lowest deviance is at 20, 24.5 and 18 °C for King Edward, Draper and the loop, but the bootstrap intervals are wide (Appendix C): the response bends gradually, as the spline

shows, rather than at a point. The spline and the two-segment line fit about equally well (Appendix B); a single straight line on the log scale fits worse on every series.

Is there a second change in the cold? Adding a change of slope at 0 °C to the model:

Series	Days below 0 °C	Below 0 °C: % per degree	0 °C to the change of slope: % per degree	p (difference)
King Edward	135	+5.2 (+1.9 to +8.6)	+6.5 (+5.5 to +7.6)	0.47
Draper	109	+4.6 (+2.5 to +6.8)	+5.1 (+3.9 to +6.3)	0.74
Kensington loop	108	+5.0 (+3.5 to +6.4)	+6.3 (+4.7 to +8.0)	0.22
City counters	169	+2.4 (+1.4 to +3.5)	+4.8 (+3.9 to +5.6)	0.00

On the Terrebonne series the two slopes are not distinguishable. The earlier analyses, which fitted two straight lines to the counts themselves, placed a change of slope at −1 °C (air temperature) or −5 °C (feels-like temperature): a count that falls by a constant percentage per degree traces a curve that flattens towards zero, and two straight lines fitted to that curve meet near where it flattens. The City counters do show a flatter slope below 0 °C.

5 Wind chill

The wind-chill index and the daytime temperature are almost the same variable over a year (correlation 0.996). A model with the wind-chill index in place of temperature and wind has an AIC lower by 2.8 to 3.9 on the Terrebonne series (Appendix B): a small difference in its favour, not enough to say that riders respond to wind chill rather than to temperature. Wind as a separate term goes with −7 (−10 to −3) % per 10 km/h at King Edward and is not distinguishable from zero at Draper and on the loop.

6 Rain, snow and wind

Condition	King Edward	Draper	Kensington loop	BIXI station	City counters
Rain, under 2 mm	−6 (−14 to +3)	0 (−7 to +8)	−23 (−33 to −12)	−2 (−8 to +3)	−13 (−18 to −8)
Rain, 2 to 10 mm	−22 (−29 to −14)	−14 (−21 to −7)	−22 (−34 to −8)	−20 (−24 to −15)	−26 (−31 to −21)
Rain, over 10 mm	−40 (−48 to −30)	−25 (−32 to −17)	−46 (−56 to −34)	−31 (−39 to −22)	−45 (−49 to −40)
Snowfall, under 2 cm	−27 (−40 to −9)	−10 (−25 to +9)	−7 (−22 to +10)	−25 (−39 to −9)	−11 (−19 to −2)
Snowfall, 2 cm or more	−36 (−52 to −14)	−24 (−41 to −2)	−18 (−34 to +2)	–	−21 (−28 to −13)
Snow on the ground, 1 cm or more	−47 (−61 to −28)	−37 (−53 to −18)	−31 (−47 to −11)	–	−18 (−31 to −2)
Wind, per 10 km/h	−7 (−10 to −3)	0 (−5 to +4)	−3 (−9 to +3)	−1 (−5 to +3)	−4 (−7 to −1)
Snowfall the day before, per cm	−1 (−4 to +1)	−1 (−2 to +1)	−1 (−5 to +2)	−4 (−8 to 0)	−1 (−2 to 0)
Saturday	−8 (−16 to +1)	0 (−9 to +9)	−34 (−42 to −24)	−11 (−16 to −5)	−36 (−41 to −31)
Sunday or holiday	−2 (−9 to +6)	+7 (−2 to +16)	−32 (−39 to −25)	−20 (−25 to −15)	−42 (−45 to −38)
December 24 to January 2	−22 (−39 to 0)	−49 (−66 to −25)	−49 (−72 to −7)	–	−62 (−68 to −55)

Percentage change in the daily count (95 % interval), compared with a day without the condition (for the day types, with a weekday). –: fewer than five such days, or not applicable.

Number of days behind each estimate:

Condition	King Edward	Draper	Kensington loop	BIXI station	City counters
Rain, under 2 mm	81	61	25	58	77
Rain, 2 to 10 mm	110	76	37	79	92
Rain, over 10 mm	45	36	15	39	47
Snowfall, under 2 cm	56	48	38	9	64
Snowfall, 2 cm or more	46	30	30	–	46
Snow on the ground, 1 cm or more	120	93	105	–	161

Light rain (under 2 mm in 24 hours) is not distinguishable from none on the camera sensors; the loop, which counts day and night, shows a reduction. The effects are multiplicative: on a January day that




Figure 4: Change in the daily count associated with each condition, temperature and time of year held fixed.

would see 20 bicycles in the window, 2 to 10 mm of rain removes about four; on a June day that would see 300, about sixty. Snow on the ground carries the persistence of a snowfall: once it is in the model, the snowfall of the day before adds little.

7 Day length, season and darkness

	Deviance accounted for (%)			of which (%)		
Series	Calendar and season alone	Weather alone	Both	Either	Weather only	Calendar and season only
King Edward	67	77	80	65	13	2
Draper	78	81	86	73	8	5
Kensington loop	87	86	92	81	5	5
BIXI station	49	53	63	38	14	10
City counters	86	81	92	75	6	11

Most of what the models account for can be credited to either set of terms: the time of year predicts the weather, and the weather predicts the time of year. The record cannot say how much of the winter low is cold and snow and how much is the season as such. What it does say is that weather matters *within* the season: the weather-only share is 5 to 13 % on the Terrebonne series, and the temperature slope is unchanged when estimated inside calendar months.

What a daylight-only sensor sees. In June the 08:00–16:00 window holds a little over half of a Telraam sensor's daily total; in December it holds more than 80 %, because there is little daylight outside it. A year-round comparison of Telraam daily totals therefore mixes a change in riding with a change in counting hours. On the loop, which counts day and night, the same window holds 46 to 57 % of the 24-hour total in every month of the hourly record, so the fixed window is a stable part of the day.




Figure 5: Share of the deviance of daily counts accounted for by the two sets of terms.




Figure 6: Left: the share of a Telraam sensor's daylight total that falls in the fixed window. Right: the share of the loop's 24-hour total in the same window.

Darkness at the same clock hour. The loop's hourly record (4,592 hours between 06:00 and 22:00 on 287 days, 1,116 of them after sunset and 231 before sunrise) allows a direct comparison: the same hour of the same type of day, in the same month and at the same temperature, when the sun is up and when it is down. An evening hour after sunset carries 27 % (19 to 34) fewer bicycles, a morning hour before sunrise 17 % (3 to 30). In the same model an hour with rain observed carries 47 % (38 to 55) and an hour with snow observed 26 % (13 to 37). This is one site and one direction; with month constants in the model, the comparison rests mostly on the weeks around the clock changes and the equinoxes.

8 Snowfall, the days after, and snow loading




Figure 7: Winter days: change in the count per 5 cm of snowfall on the day and on each of the seven days before.

On winter days (November 16 to March 31), with temperature, rain, wind, day type and snow on the ground in the model:

Days after the snowfall	King Edward	Draper	Kensington loop
0	−12 (−24 to +2)	−17 (−23 to −10)	−14 (−27 to +2)
1	+1 (−14 to +18)	−3 (−10 to +6)	+2 (−13 to +18)
2	+6 (−8 to +22)	−2 (−17 to +15)	−16 (−29 to −1)
3	−6 (−21 to +11)	−3 (−15 to +10)	−11 (−26 to +8)
4	−14 (−28 to +2)	−11 (−22 to 0)	−7 (−24 to +13)
5	−13 (−23 to −2)	−6 (−19 to +10)	+3 (−12 to +21)
6	+8 (−1 to +18)	−14 (−20 to −8)	−6 (−21 to +12)
7	+9 (−2 to +22)	−12 (−19 to −4)	−11 (−24 to +4)

Percentage change per 5 cm of snow (95 % interval).

The day of a snowfall shows the clearest reduction (12 to 17 % per 5 cm; the interval excludes zero at Draper only). On the following days the estimates scatter around zero and a few negative values, with wide intervals and no pattern common to the three series; winter counts are small (a few dozen a day), each series has 132 to 191 winter days, and snow on the ground already carries part of the after-effect. An average over snowfall episodes was also computed (`snow_episodes_summary.csv`); with fewer than ten episodes of 10 cm or more per series it is too noisy to read.

Snow-loading days.

Series	Winter days	of which loading days (sector CDN-301)	Loading day: change (%)	Temperature: % per gree (winter only)
King Edward	191	44	−4 (−29 to +30)	+7 (+5 to +9)
Draper	154	37	−21 (−48 to +21)	+7 (+5 to +9)
Kensington loop	132	36	+2 (−25 to +39)	+7 (+5 to +10)

On the days the City loaded snow in the sector, the counts are not distinguishable from those of other winter days with the same temperature and snow. The intervals are wide enough to contain a reduction of a quarter or more as well as no change; the record does not settle the question. In winter alone, each degree goes with about 7 % more bicycles on all three series.

9 Agreement between the series

The five series count different things with different instruments. Their relative responses agree in direction everywhere and broadly in size: the temperature curves of the three Terrebonne series lie within each other's intervals (Figure 3); the rain and snow effects overlap (Figure 4). The differences that stand out: the BIXI station and the City counters respond less steeply to temperature than the Terrebonne counters; the loop shows a weekend reduction of about a third that the camera sensors, counting 08:00–16:00, do not show; and the loop responds to light rain where the camera sensors do not. The first may reflect who rides where; the others are what one would expect from the difference between a 24-hour count and a daytime window. None of this was tested formally: the series cover different periods.

10 Year to year, after weather

The counting-year terms of the models compare the same series across years with weather, day type and time of year held fixed:

Series	Counting years (from August 15)	Days in the later year	Change (%, 95 % interval)
King Edward	2025–26 against 2024–25	272	−1 (−11 to +9)
King Edward	2026–27 against 2024–25	86	+27 (+12 to +45)
Draper	2025–26 against 2024–25	245	−12 (−20 to −5)
BIXI station	2025–26 against 2024–25	229	−10 (−16 to −4)
City counters	2025–26 against 2024–25	361	−7 (−14 to 0)

These figures are to be read with care. A year term absorbs anything that differs between the years besides the weather: the sensors' gaps fall in different months each year, a camera sensor's field of view can change when it is remounted, and the 2026–27 counting year so far holds only August 15 to early November. King Edward's 2026–27 figure compares that autumn with the autumn of 2024. Draper's and the BIXI station's later years are merged (too few days after August 15, 2026), and the City counters' 2026–27 term is not shown (its first year holds no autumn to compare with). The loop has a single year and no such term.

11 How well the models predict

Each calendar month in turn was left out, the model fitted on the rest, and the month predicted:

		Median absolute error (%)		R^2 on the log scale	
Series	Days predicted	Fitted	Month left out	Fitted	Month left out
King Edward	691	22	23	0.84	0.82
Draper	505	19	20	0.88	0.86
Kensington loop	284	20	25	0.92	0.87
BIXI station	441	15	17	0.59	0.50
City counters	661	15	18	0.94	0.93

The typical day is predicted to within about a fifth to a quarter of its count, and little accuracy is lost on a month the model has not seen. A fifth of a small winter count is a few bicycles.

12 Limits

- **Daylight-only sensors.** The Telraam figures are 08:00–16:00 counts. They say nothing about evening and night riding, and their levels are never to be compared with the loop's.
- **Gaps.** King Edward has no valid day in February 2026 and few in December 2025 and July 2026; Draper has none from February to April 2026. The winter of 2025–26 is therefore thinly covered by the camera sensors and carried mostly by the loop.
- **One loop site, one winter, one direction** in the models; the westbound channel is not used after the fault of March 2026, and for consistency not before.
- **Weather at the airport,** 9 km away; precipitation as 24-hour totals; the state of the lane itself (cleared or not, icy or not) is not observed. Snow on the ground at the airport stands in for it.
- **Season and weather are confounded** and no model can fully separate them from this record.
- **Association, not cause.** A percentage per degree describes days in the record; it does not predict what would happen under conditions the record does not contain.
- **Uptime.** Telraam hours with an uptime under 0.5 are dropped; whether the remaining counts are already scaled for uptime is not documented.

A Valid days by month

Month	King Edward	Draper	Kensington loop	BIXI station	City counters
2024-08	14	0	0	31	0
2024-09	26	4	0	30	0
2024-10	28	18	0	31	0
2024-11	29	21	0	21	0
2024-12	19	21	0	0	0
2025-01	26	24	0	0	30
2025-02	27	22	0	0	21
2025-03	30	22	0	0	30
2025-04	30	30	0	20	30
2025-05	31	29	0	31	31
2025-06	30	30	0	30	30
2025-07	31	30	6	29	31
2025-08	30	27	31	14	31
2025-09	28	28	30	30	30
2025-10	31	31	31	31	31
2025-11	30	27	30	30	30
2025-12	14	26	31	31	31
2026-01	25	12	29	25	29
2026-02	0	0	28	23	28
2026-03	22	0	29	28	29
2026-04	30	0	30	30	30
2026-05	31	18	28	31	31
2026-06	29	25	27	30	30
2026-07	10	19	31	31	31
2026-08	22	16	31	31	31
2026-09	30	17	0	0	30
2026-10	31	13	0	0	31
2026-11	8	0	0	0	6

Loop: days valid in the daily table (both directions); the models use the eastbound channel, to May 10, 2026. BIXI: days with at least one trip.

B Model comparison

	AIC, temperature as			AIC, without the wind term		
Series	one line	two segments	spline	temperature	wind chill	Correlation, temperature and wind chill
King Edward	7286.8	7269.8	7273.0	7276.2	7265.9	0.996
Draper	5413.0	5384.9	5389.1	5383.0	5381.2	0.996
Kensington loop	2743.3	2734.9	2734.5	2733.8	2732.2	0.995
BIXI station	3687.1	3676.5	3683.0	3674.8	3674.3	0.995
City counters	12286.9	12242.5	12218.5	12249.8	12237.4	0.996

Lower is better; differences of less than about 2 are not meaningful. AIC values are comparable only within a row.

C Where the slope changes: deviance profile




Figure 8: Deviance of the two-segment model as the change of slope is moved, each series relative to its own minimum.

D The earlier analyses and what was kept

Between September 2024 and May 2026 the author explored the Telraam counts in MATLAB. Those scripts were read before this work; none of their code is used here. In summary:

Earlier analysis	What it did	Here
Regressions of February 2025	Least-squares fits of daily counts on temperature or feels-like temperature, squared terms, precipitation; two straight lines with a change of slope found by search (-1 and $-5\,^{\circ}$C); King Edward only, 290 days	Count model on the percentage scale; change of slope estimated with an interval, segments joined; the cold change of slope tested and not found
Sunset adjustment	Counts up to 15:00 multiplied by a fixed ratio (1.65 on weekdays, 1.37 on weekends) to stand for the whole day	Replaced by the fixed 08:00–16:00 window, with no scaling
Filling of dark hours	Five ways to fill the hours after sunset from a reference daily profile	Not kept: no count is estimated for hours the sensor did not see
Daily weather regression of November 2025	Least squares on before-15:00 counts with temperature, precipitation, wind, weekend and a linear trend; ten interaction tests; weather classes compared by t-tests	Kept as the main model, with a count distribution, season terms in place of the trend and autocorrelation-robust intervals
Weekly season-and-anomaly split	Share of variance from a seasonal baseline and from weather anomalies, weekly	Kept, daily, as the deviance shares and the within-month model
Lagged-snow regression and snow-event averages, winter 2024–25	Ten lags of a snow proxy (precipitation on days below 0 °C); average response by snowfall class	Kept with measured snowfall and snow on the ground, two winters, three series, and the City's loading record
Winter retention, storm recovery, composite scores	Ratios of winter to summer counts by mode; one storm; ad hoc scores	Not kept
Weather source	A commercial city-level feed	Environment and Climate Change Canada, Montréal-Trudeau

E Sources

Environment and Climate Change Canada, historical climate data, MONTREAL INTL A (7025251), downloaded November 10, 2026. Telraam (citizen-operated sensors; the counts belong to Telraam). Arrondissement de Côte-des-Neiges–Notre-Dame-de-Grâce, loop counter at Terrebonne/Kensington, access requests 2026-300 (released June 4, 2026) and 2026-581 (released October 9, 2026). Ville de Montréal open data: permanent bicycle counters (files of November 7, 2026), snow-removal transactions; CC BY 4.0. BIXI Montréal open data, trip history 2024–2026 (downloaded November 10, 2026; listed by the City's portal under CC BY 4.0, attribution to BIXI Montréal). Details and licences: the folder's README.