

# Questions & Solutions



# Contents

<b>Introduction</b>	<b>vi</b>
How to Contribute	vi
Correcting Mistakes	vi
Contributing Solutions	vi
Problem Proposals	vi
Feature Suggestions	vi
Contact Us	vii
Contributors	vii
<b>1 Season 1</b>	<b>1</b>
1.1 Week 1	1
1.1.1 Intersecting Circles	1
1.1.2 Guava Juice	2
1.1.3 Complex Roots	3
1.1.4 Exponents	4
1.1.5 Volumes of Cubes	5
1.1.6 Complex Mess	6
1.1.7 Paper Eating	7
1.2 Week 2	8
1.2.1 Regenerative Watermelons	8
1.2.2 Largest Prime Factor	9
1.2.3 Real Roots	10
1.2.4 The Meme factor	11
1.2.5 Human Wolfram	12
1.2.6 Guess the Config	13
1.2.7 A Quadratic Mess	14
<b>2 Season 2 (.19's Season)</b>	<b>15</b>
2.1 Week 1	15
2.1.1 A Sequence of 5's	15
2.1.2 Brainy's Happy Set	16
2.1.3 MODSbot's Escape!	17
2.1.4 Sides of a Polygon	18
2.1.5 2p	20
2.1.6 Slippery Rooks	21
2.1.7 Sets of Integer Solutions	23
2.2 Week 2	24
2.2.1 A Game of Deductions	24
2.2.2 Maximising Exponents	25
2.2.3 Colourful Problem	26
2.2.4 Combinatorial Addition	27
2.2.5 Expected Value	28
2.2.6 Infinite Product	29
2.2.7 Projective Geo	30
<b>3 Trigonometric Troubles (Season 3)</b>	<b>31</b>
3.1 Week 1	31
3.1.1 Maximising Trig. Function	31
3.1.2 Areas inside a square	32
3.1.3 Length of SQ	33
3.1.4 Sum of Tan's	34
3.1.5 Distance from the Orthocentre	35
3.1.6 Minimising the Diagonal	36

3.1.7	Circumcentre . . . . .	37
<b>4</b>	<b>Piboi's Bashy Combo (Season 4) [VOIDED]</b>	<b>38</b>
4.1	Week 1 . . . . .	39
4.1.1	A Tricky Shuffle [VOIDED] . . . . .	39
4.1.2	Coloured Markers [VOIDED] . . . . .	40
4.1.3	Common Names [VOIDED] . . . . .	41
4.1.4	Funny Questions [VOIDED] . . . . .	42
4.1.5	Colourful Integers [VOIDED] . . . . .	43
<b>5</b>	<b>Back to School (Season 5)</b>	<b>44</b>
5.1	Week 1 . . . . .	44
5.1.1	A Tricky Combination . . . . .	44
5.1.2	Geometric Sequence . . . . .	45
5.1.3	Simultaneous Equation? . . . . .	46
5.1.4	Tangent Circles . . . . .	47
5.1.5	Santa's Elves . . . . .	50
5.1.6	2016 Algebra . . . . .	51
5.1.7	$x^y$ 's . . . . .	52
5.2	Week 2 . . . . .	53
5.2.1	Pentagons . . . . .	53
5.2.2	Digits in a String . . . . .	54
5.2.3	Relatively Prime Function . . . . .	55
5.2.4	Intersecting Circles . . . . .	56
5.2.5	Discord Ping Fight . . . . .	57
5.2.6	Falling Cards . . . . .	58
5.2.7	Computing the Area of a Triangle . . . . .	59
<b>6</b>	<b>Third Week of School (Season 6)</b>	<b>60</b>
6.1	Week 1 . . . . .	60
6.1.1	Maximising Square Factors . . . . .	60
6.1.2	Njoy's Balls . . . . .	61
6.1.3	Aiya's Function . . . . .	62
6.1.4	The AIME Cyclic . . . . .	63
6.1.5	Prime Floors . . . . .	64
6.1.6	Nested Periodic Functions . . . . .	65
6.1.7	Braniy's Party . . . . .	65
6.2	Week 2 . . . . .	66
6.2.1	Negative Sum . . . . .	66
6.2.2	Doubling Digit Sum . . . . .	67
6.2.3	Tony Vs. Wang . . . . .	68
6.2.4	Equilateral Decagon . . . . .	69
6.2.5	Triangles on a Cubic . . . . .	70
6.2.6	Jumping Frogs . . . . .	71
6.2.7	Minimising Ratios . . . . .	72
<b>7</b>	<b>PSC's Adventure (Season 7)</b>	<b>73</b>
7.1	Week 1 . . . . .	73
7.1.1	The Jungle Polygon . . . . .	73
7.1.2	Uphill and downhill . . . . .	74
7.1.3	Just a Beauty! . . . . .	75
7.1.4	Probability of Intervals . . . . .	76
7.1.5	Dividing Functions . . . . .	78
7.1.6	Paper Monster . . . . .	79
7.1.7	Tan's Optimisation . . . . .	80

7.2	Week 2	81
7.2.1	A Digital Product equal to Half the Number	81
7.2.2	Unique Odd Numbers	82
7.2.3	Maximising $a^4 + b^4$	83
7.2.4	Minimising Perimeter	84
7.2.5	Binary Blocks	85
7.2.6	Funkey Triangles	87
7.2.7	A Lotta Touples!	88
<b>8</b>	<b>A New Chapter (Season 8)</b>	<b>89</b>
8.1	Week 1	89
8.1.1	Smallest Non-Factor	89
8.1.2	Product of Radii	90
8.1.3	Two-to-One Functions	91
8.1.4	Modular Powers	92
8.1.5	Australian Nim	93
8.1.6	Another Geo Config	94
8.1.7	Product of Root Differences	95
8.2	Week 2	97
8.2.1	Counting Intersections	97
8.2.2	The Race	98
8.2.3	Remainder of Sums of Squares	99
8.2.4	Another 255 Subsets	100
8.2.5	4-5-6 Triangle	101
8.2.6	“What’s a signed integer?”	102
8.2.7	Diagonals in a Convex 1001-gon	103
<b>10</b>	<b>CCCC After Math (Season 10)</b>	<b>104</b>
10.1	Week 1	104
10.1.1	EpicXtroll	104
10.1.2	Similar Triangles	105
10.1.3	Factorial Sums and Divisors	106
10.1.4	Brainy’s Passcode	107
10.1.5	Production Loop	108
10.1.6	Weird polynomial	109
10.1.7	Weird triangle	110
10.2	Week 2	112
10.2.1	Real Square Roots	112
10.2.2	”What’s a positive integer?”	113
10.2.3	Converging Tangent Circles	114
10.2.4	Maze Escape	115
10.2.5	Primes divide powers of 5	116
10.2.6	tRiViaL	117
10.2.7	Lamps in a Circle	118
<b>11</b>	<b>errආආ: ආආ ආආආ ආආආආආආ ආආආ ආආ ආආආ ආආආආ. (Season 11)</b>	<b>119</b>
11.1	Week 1	119
11.1.1	Integral AM-GM	119
11.1.2	d20 but not D20	120
11.1.3	Is it better to walk or run in the rain?	121
11.1.4	Don’t over think this	122
11.1.5	Genfun is Not Fun	123
11.1.6	Harmonic Sum	124
11.1.7	Lonely Tetrahedron	126

11.2	Week 2	127
11.2.1	Don't over think this part 2	127
11.2.2	Lattice Points on a Circle	128
11.2.3	WA is Required	129
11.2.4	Hard	130
11.2.5	hard	132
<b>12</b>	<b>Return to Normalcy (Season 12)</b>	<b>134</b>
12.1	Week 1	134
12.1.1	Quadrilateral Perimeter	134
12.1.2	Involution Convolution	135
12.1.3	Coordinates without Bash	136
12.1.4	April Fool's	137
12.1.5	FE	138
12.1.6	So this is where American problem quality went	139
12.1.7	640 Generators of Pain	140
12.2	Week 2	143
12.2.1	Even Numbers with Restrictions	143
12.2.2	Enjoyable	144
12.2.3	Isosceles Tetrahedron	145
12.2.4	Sums of powers of 2 but weird	146
12.2.5	"D4 I swear"	147
12.2.6	"change the point distribution up so weekends don't carry people"	148
12.2.7	Notice me, sqing-senpai	149
<b>13</b>	<b>Quality Uncontrol (Season 13)</b>	<b>150</b>
13.1	Week 1	150
13.1.1	Choi's Balls	150
13.1.2	Biscriminant	151
13.1.3	IMOK Maclaurin2012 P6	152
13.1.4	NICE Geometry	153
13.1.5	Graph Theory in Disguise	154
13.1.6	What the H-E-double hockey sticks?	155
13.1.7	"What's an mgf?"	156
13.2	Week 2	157
13.2.1	It's Okay :)	157
13.2.2	Points on a Line	158
13.2.3	Diamond Operator	159
13.2.4	Weird Wilson	160
13.2.5	Get the CRUX of this inequality!	161
13.2.6	Don't overthink this	162
13.2.7	The Party, The Politburo, and the PSC	163
<b>14</b>	<b>Mityushikha Bay (Season 14)</b>	<b>164</b>
14.1	Week 1	164
14.1.1	Rectangle Rotation	164
14.1.2	NJOY Troll (again)	165
14.1.3	You Just Got Vector'ed	166
14.1.4	Answer Extraction	167
14.1.5	Not a Fan	168
14.1.6	you're welcome, brainy	169
14.1.7	RDS-1	170
14.2	Week 2	172
14.2.1	All About that Base	172
14.2.2	Number of QoTD Seasons	173

14.2.3	Square and Point . . . . .	174
14.2.4	2012 HMMT A6 . . . . .	175
14.2.5	[under construction] . . . . .	176
14.2.6	Projective Pain . . . . .	177
14.2.7	AN602 . . . . .	178
<b>15</b>	<b>SALT-1 (Season 15)</b>	<b>180</b>
15.1	Week 1 . . . . .	180
15.1.1	"What are distinct integers?" . . . . .	180
15.1.2	Troll Pigeonhole . . . . .	181
15.1.3	WA Can't Solve This . . . . .	182
15.1.4	Prime Product . . . . .	183
15.1.5	The Rare Non-Horrible States Problem . . . . .	184
15.1.6	Minimal Area, Minimal Difficulty for a Saturday . . . . .	185
15.1.7	Detente . . . . .	186
15.2	Week 2 . . . . .	187
15.2.1	Senior Math; What Challenge? . . . . .	187
15.2.2	Prisoner's Stupid Dilemma . . . . .	188
15.2.3	Coordinates without Bash . . . . .	189
15.2.4	WA Can't Solve This, Either . . . . .	190
15.2.5	WA Can't Solve This, Either . . . . .	191
15.2.6	cHriS geO . . . . .	192
15.2.7	Treaty Violation . . . . .	193
<b>16</b>	<b>Quality Control (Season 16)</b>	<b>194</b>
16.1	Week 1 . . . . .	194
16.1.1	The Rare non-Troll NJOY Problem . . . . .	194
16.1.2	The Common Troll NJOY Problem . . . . .	195
16.1.3	Parallelogram PTSD . . . . .	196
16.1.4	WA Could Solve This and you Still Failed . . . . .	197
16.1.5	Surprising System . . . . .	198
16.1.6	Really Weird NT . . . . .	199
16.2	Week 2 . . . . .	200

## Introduction

Welcome to the OpenPOTD solutions booklet! Here you'll find answers & solutions to all past seasons.

Solutions for season one were entirely written up by Brainysmurfs#2860, while sjbs#9839 has overseen most of season two. From season three onwards, Yuchan (Angry Any#4319) has also been contributing problem proposals and solutions.

Where possible, from season two onwards, We have tried to include the officially provided solutions to problems, or adapted them in line with any changes to the problem statement, and in most cases also and filled in the gaps as best I could, to make solutions more approachable to beginners. For the more well known questions that are featured in a season (namely problems from the International Mathematical Olympiad), instead of providing our own solution, we have included the *Art of Problem Solving* forum post on the question, which will contain multiple solution write-ups, as well as discussion about the problem.

In many circumstances problems have not come with official write-ups - or indeed write-ups of any kind - and thus have required us to provide our own. In these cases we humbly apologise for any mistakes (or fakesolves!) in advance. If you do notice any mistakes, check out [How to contribute](#).

## How to Contribute

If you would like to contribute to the project - be that through correcting a mistake in the document, wanting to contribute a solution write up - or even proposing a problem for a future season, here's how.

### Correcting Mistakes

If you notice any mistakes while going through the solutions document, be it a typo or missing or incorrect information about a particular problem, feel free to submit a push request with the fix. If you don't feel comfortable doing that, you can always contact us (see the [Contact Us](#) to find out how), and we'll be happy to fix the error. Alternatively, you can open up an [Issue](#) on the GitHub, or mention it in the [discussions tab](#).

### Contributing Solutions

Similarly to correcting any mistakes, if you would like to contribute a write-up to a particular problem, you can submit a push containing the solution - doing as Romans do (i.e. just look at how others have submitted write-ups and copy that). If you are submitting a push, make sure you edit `preamble.sty` to include your Discord information in a macro (scroll to the bottom of the file and you'll see), so that you can include yourself in the contributors list. Furthermore, make sure that you credit your solution with `[Write up by ...]`. If any of that sounds complicated or you forget to add that information, that's fine - we'll add it for you.

If creating push requests and fiddling with LaTeX isn't your thing, we'll gladly help you type it up if you write it out in plaintext or whatever medium you feel most comfortable using (so long as we can understand it!) - to do this just send it to us using any of the options in [Contact Us](#). Similarly, you can use the GitHub [discussions tab](#) and post it there.

***Note: feel free to submit alternative solutions to any past problems, be it from season 1, or the most recent***

### Problem Proposals

If you'd like to submit a problem proposal - be it an original problem, or just a particular problem you found interesting - we're always on the lookout for new problems! For original problems, please ensure you submit the problem with a solution. As with solution write-ups, though sending us a `.tex` file is preferred, it's completely fine to just send a plaintext write up, or a screenshot etc. and we can deal with it from there. The same goes for non-original problems, though solutions aren't required, they would be greatly appreciated. If you are submitting a non-original problem please ensure you include the problem source. Please do not use a public medium to submit a problem proposal - messaging one of us on Discord would be the preferred method of communication (See [Contact Us](#)).

## Feature Suggestions

If you have any ideas when it comes to improving the bot or project, the best place to do that is in the #Suggestions channel, or any of the other methods listed in [Contact Us](#), such as using the GitHub [discussions tab](#).

## Contact Us

The best way to contact us is through the [OpenPOTD Discord server](#), however, you may also contact us through the [discussions tab](#) on Github. Alternatively, you can DM any of us on Discord:

- sjbs#9839 (434767660182405131)
- brainysmurfs#2860 (281300961312374785)
- Angry Any#4319 (580933385090891797)

## Contributors

Thank you (in no particular order) to the following contributors: <sup>1</sup>

AiYa#2278 (675537018868072458): Solution write-ups ([2.1.1](#), [2.1.3](#), [2.1.4](#), [5.1.4](#), [6.1.2](#)), Original problem proposal ([6.1.3](#))

Tony Wang#1729 (541318134699786272): Original problem proposal ([1.1.2](#))

Charge#3766(481250375786037258): Original problem proposal ([1.1.6](#), [4.1.1](#),[4.1.2](#),[4.1.3](#),[4.1.4](#),[4.1.5](#), [5.2.5](#))

bfan05#5219 (692851547062665317): Original problem proposal ([1.1.7](#))

Kiesh#0917 (544960202101751838): Original problem proposal ([1.2.3](#))

ChristopherPi#8528 (696497464621924394): Problem proposal ([2.1.6](#)), Original problem proposal ([7.1.5](#))

Keegan#9109 (116217065978724357): Original problem proposal ([2.2.3](#))

Slaschu#5267 (296304659059179520): Solution write-up ([2.2.5](#))

Bahnhofstrasse#8974 (723413754800373780): Typo fixes

TaesPadhary#8557 (665057968194060291): Problem proposal ([5.2.2](#), [5.2.4](#), [7.1.3](#)), Original problem ([5.2.3](#), [7.1.2](#)), Typo fixes

aops#0436 (712027036511633429): Typo fixes

HoboSas#3200 (310725130097786880): Typo fixes, Original problem proposal ([7.1.4](#))

Radial Function#6976 (785583756957450280): Typo fixes

RishiNandha Vanchi#3379 (562608039224410112): Solution write-ups ([7.2.5](#))

<sup>1</sup>The numbers are in the format: Season.Week.Problem



## §1 Season 1

### §1.1 Week 1

#### §1.1.1 Intersecting Circles

---

*Source: Senior Mathematical Challenge, 2015 Q4*

*Proposer: brainysmurfs#2860 (281300961312374785)*

*Problem ID: 21*

*Date: 2020-10-27*

---

Consider the positive integer  $N$ , and Two internally tangent circles  $\Gamma_1$  and  $\Gamma_2$  are given such that  $\Gamma_1$  passes through the center of  $\Gamma_2$ . Find the fraction of the area of  $\Gamma_1$  lying outside  $\Gamma_2$ . If this fraction is  $\frac{a}{b}$  where  $\gcd(a, b) = 1$ , then find  $a + b$ .

*Solution.*

Suppose  $\Gamma_2$  has radius  $2r$ . Since  $\Gamma_1$  is internally tangent to  $\Gamma_2$  and passes through its centre, the radius of  $\Gamma_1$  is half the radius of  $\Gamma_2$ , i.e. just  $r$ . So fraction of the area of  $\Gamma_1$  lying outside  $\Gamma_2$  is  $\frac{4\pi r^2 - \pi r^2}{4\pi r^2} = \frac{3}{4}$ . Since  $3 + 4 = 7$ , the answer is 7 □

**§1.1.2 Guava Juice**

---

*Source: Original Problem*

*Proposer: Tony Wang#1729 (541318134699786272)*

*Problem ID: 22*

*Date: 2020-10-28*

---

The ingredients list of a Guava Juice Drink is as follows:

Water (80%), Guava Juice, Sugar, Fructose (3%), Sodium Carboxymethyl Cellulose, Citric Acid, Flavour, Vitamin C (0.04%)

Assuming only that the ingredients list is ordered by their constituent percentage in the drink (which are not necessarily distinct), find the maximum and minimum possible percentage of Guava Juice in the drink. If their difference is  $n$ , submit  $100n$ .

*Solution.*

Note there are 2 ingredients between water and fructose, and 3 between fructose and Vitamin C.

For the amount of guava juice to be maximised, everything should be minimised. In particular, there should be 0.04% of Sodium Carboxymethyl Cellulose, Citric Acid, and Flavour; and 3% of Sugar and Fructose. This gives us a Guava Juice percentage of 13.84%.

For the amount of guava juice to be minimised, everything else should be maximised. In particular, there should be 3% of Sodium Carboxymethyl Cellulose, Citric Acid, and Flavour; and equal amounts of sugar and guava juice. This gives us a guava juice percentage of 5.48%.

This gives us  $13.84 - 3.98 = 9.86\%$ . Multiplying this by 100 gives us 986.

□

**§1.1.3 Complex Roots**

---

Source: New South Wales Higher School Certificate '4U', 2020 Q2

Proposer: brainysmurfs#2860 (281300961312374785)

Problem ID: 23

Date: 2020-10-29

---

Given that  $z = 3 + i$  is a root of  $z^2 + pz + q = 0$ , where  $p$  and  $q$  are real, find the values of  $p$  and  $q$ , and submit  $p + q$ .

*Solution.*

Applying the Complex Conjugate Theorem,  $z = 3 - i$  is also a root of the quadratic. Expanding  $(z - 3 - i)(z - 3 + i)$  gives us  $z^2 - 6z + 10$ . Thus  $p = -6$  and  $q = 10$ , meaning  $p + q = \boxed{4}$ . □

**§1.1.4 Exponents**

---

*Source: Singapore Mathematical Olympiad Junior Round 1, 2001 Q4*

*Proposer: brainysmurfs#2860 (281300961312374785)*

*Problem ID: 25*

*Date: 2020-10-30*

---

If  $a$  and  $b$  are positive reals such that  $a^b = b^a$  and  $b = 2a$ , then the value of  $b^2$  is?

*Solution.*

A direct search yields that  $2^4 = 4^2$  and  $4 = 2 \times 2$ . The problem implies that such a unique value for  $b$  exists, hence  $b^2 = \boxed{16}$ . □

**§1.1.5 Volumes of Cubes**

---

*Source: New Zealand Senior Mathematics Competition Round 2, 2019 Q7*

*Proposer: brainysmurfs#2860 (281300961312374785)*

*Problem ID: 26*

*Date: 2020-10-31*

---

Two cubes with positive integer side lengths are such that the sum of their volumes is numerically equal to the difference of their surface areas. Find the sum of their volumes.

*Solution.*

A direct search yields that cubes of side length 4 and 2 satisfy the condition. The answer is thus  $4^3 + 2^3 =$   
72. □

**§1.1.6 Complex Mess**

Source: Original Problem

Proposer: Charge#3766(481250375786037258)

Problem ID: 24

Date: 2020-11-01

Suppose

$$\left(\sqrt{5} + i\sqrt{10 - 2\sqrt{5}} + 1\right)^{2020} = a^b$$

for some  $a, b \in \mathbb{Z}$  where  $b$  is maximised. Compute  $a + b$ .

*Solution.*

Notice that  $\left(\sqrt{5} + i\sqrt{10 - 2\sqrt{5}} + 1\right) = 4\text{cis}\frac{\pi}{5}$ . In particular, by De Moirve's Theorem,

$$\left(\sqrt{5} + i\sqrt{10 - 2\sqrt{5}} + 1\right)^{2020} = 4^{2020}\text{cis}404\pi = 2^{4040}$$

.

So the answer is  $2 + 4040 = \boxed{4042}$ .

□

### §1.1.7 Paper Eating

Source: Original Problem

Proposer: bfan05#5219 (692851547062665317)

Problem ID: 31

Date: 2020-11-02

Tan and Wen are writing questions for CCCC at a rate of 1 per minute. Immediately after Tan writes a question, Wen eats his paper with probability  $\frac{1}{7}$ , so that Tan must restart.

After an extremely long time (assume infinite), Tony Wang walks in. What is the expected number of questions Tony Wang sees written on Tan's paper?

*Solution.*

Let  $E_n$  be the expected number of questions Tony Wang sees written on Tan's paper after  $n$  minutes.

Then the following recurrence holds:

$$E_{n+1} = \frac{6}{7}(E_n + 1) + \frac{1}{7} \cdot 0$$

because with  $\frac{6}{7}$  probability, Tan writes another question without Wen eating it, and with  $\frac{1}{7}$  probability, Wen eats all of Tan's questions.

Claim 1:  $E_n < 6 \forall n$ .

*Proof.* The proof is by induction. Note  $E_0 = 0$ . If  $E_k < 6$ , then  $E_{k+1} = \frac{6}{7}(E_k + 1) < \frac{6}{7}(6 + 1) = 6$ . This completes the proof. ■

Claim 2:  $E_{n+1} > E_n \forall n$ .

*Proof.* Note that since  $E_k < 6 \forall k$ , we have  $E_{k+1} = \frac{6}{7}(E_k + 1) > \frac{6}{7}(E_k + \frac{1}{6}E_k) = E_k$ . ■

Now by the Monotone Bounded Convergence Theorem, the sequence  $(E_n)$  converges to a limit. Suppose  $\lim E_n = L$ . Then note  $\lim E_{n+1} = L$  since changing the first terms of a sequence does not change the overall convergence. So since  $E_{n+1} = \frac{6}{7}(E_n + 1)$ ,  $\lim E_{n+1} = \lim \frac{6}{7}(E_n + 1)$ . So  $L = \frac{6}{7}(L + 1)$ .

Solving this, we obtain  $L = \boxed{6}$ . □

## §1.2 Week 2

### §1.2.1 Regenerative Watermelons

---

Source: British Mathematical Olympiad, Round 12018 Q2

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 22

Date: 2020-11-03

---

Out of 100 regenerative watermelons, each of six friends eats exactly 75 watermelons. There are  $n$  watermelons eaten by at least five of the friends. What is the sum of the largest and smallest possible values of  $n$ ?

*Note: The watermelons are regenerative to allow multiple people to eat the same watermelon. (But the same person cannot eat the same watermelon more than once)*

*Solution.*

Maximum: The maximum is 90. Take the sum over all watermelons of how many times they were eaten. This is 450, since all  $6 \cdot 75 = 450$ . Then since  $5n \leq 450$ ,  $n \leq 90$ . The construction for the maximum is: staff member  $k$  ( $k = 1, 2, \dots, 6$ ) eats all watermelons from 1 to 90 except those  $k \pmod{6}$ .

Minimum: The minimum is 25. We intuit that in the minimum case all the watermelons were either eaten 6 times or 4. Then  $6a + 4b = 450$ , and  $a + b = 75$ . Solving yields  $a = 25$  and  $b = 75$ .

The answer is  $25 + 90 = \boxed{115}$ .

□



**§1.2.2 Largest Prime Factor**

---

*Source: Senior Mathematical Challenge, 2015 Q23*

*Proposer: Unknown*

*Problem ID: 27*

*Date: 2020-11-04*

---

Given four different non-zero digits, it is possible to form 24 different numbers containing each of these four digits. What is the largest prime factor of the sum of the 24 numbers?

*Solution.*

Say the digits are  $a$ ,  $b$ ,  $c$ , and  $d$ . For each digit, it appears in the units place 6 times, the tens place 6 times, the hundreds place 6 times, and the thousands place 6 times.

Hence the sum of the 24 numbers is  $6666(a + b + c + d) = 2 \cdot 3 \cdot 11 \cdot 101(a + b + c + d)$ . Since  $a + b + c + d < 40$ , it cannot have any prime factors larger than 101.

The largest prime factor of the sum of the 24 numbers is thus 101.

□

**§1.2.3 Real Roots**

---

*Source: Original Problem*

*Proposer: Kiesh#0917 (544960202101751838)*

*Problem ID: 28*

*Date: 2020-11-05*

---

If the polynomial  $x^2 + bx + 101 = 0$  has integer roots  $m$  and  $n$ , where  $|m| > |n|$ , then what is the sum of the positive integer divisors of  $\frac{m}{n}$ ?

*Solution.*

By Vieta's Formula  $101 = mn$ . Since 101 is prime,  $m, n = \pm 101, \pm 1$ . So  $\frac{m}{n} = \pm 101$ . The positive divisors of  $\pm 101$  are 1 and 101, so their sum is  $1 + 101 = \boxed{102}$ . □

### §1.2.4 The Meme factor

---

*Source: Original Problem*

*Proposer: Angry Any#4319 (580933385090891797)*

*Problem ID: 32*

*Date: 2020-11-06*

---

What is the sum of all integers  $x$  such that  $\frac{6969}{x}$  is an integer?

*Solution.*

Note that if  $\frac{6969}{x}$  is an integer, so is  $\frac{6969}{-x}$ . So the sum is  $\boxed{0}$ .

□

**§1.2.5 Human Wolfram**

---

*Source: Original Problem*

*Proposer: Charge#3766(481250375786037258)*

*Problem ID: 33*

*Date: 2020-11-07*

---

Compute the number between 1000 and 2000 that divides

$$69^{69} - 5^{69} + 6^{69}.$$

*Solution.*

Let  $N = 69^{69} - 5^{69} + 6^{69}$ .

Claim 1:  $64 \mid N$ .

*Proof.* Note that  $69 \equiv 5 \pmod{64} \Rightarrow 69^{69} \equiv 5^{69} \pmod{64}$ . Further,  $6^{69} = 64 \cdot 3^6 \cdot 6^{63}$ . So  $64 \mid N$ . ■

Claim 2:  $25 \mid N$ .

*Proof.* Note that  $69 \equiv (-6) \pmod{25} \Rightarrow 69^{69} \equiv (-6)^{69} \pmod{25}$ , since 69 is odd. Thus  $25 \mid N$  as required. ■

Since  $64 \cdot 25 = 1600$  and the question implies there is only one answer, we obtain that the required number is 1600. □

**§1.2.6 Guess the Config**

---

Source: *British Mathematical Olympiad, Round 22008 Q2*

Proposer: *sjbs#9839 (434767660182405131)*

Problem ID: *34*

Date: *2020-11-08*

---

Let triangle  $ABC$  have incentre  $I$  and circumcenter  $O$ . Suppose that  $\angle AIO = 90^\circ$  and  $\angle CIO = 45^\circ$ . Suppose the ratio  $AB : BC : CA$  can be expressed as  $a : b : c$  where  $\gcd(a, b, c) = 1$ . Find  $a + b + c$ .

*Solution.*

Place a  $3-4-5$  triangle on the coordinate plane with  $A = (0, 0)$ ,  $B = (3, 0)$  and  $C = (0, 4)$ . We can compute that  $I = (1, 1)$  and  $O = (1.5, 2)$ . This arrangement of points satisfies the question's constraints, and so the answer is  $3 + 4 + 5 = \boxed{12}$ . □

## §1.2.7 A Quadratic Mess

Source: Singapore Mathematical Olympiad, Open Section Round 2, 2004 Q2

Proposer: brainysmurfs#2860 (281300961312374785)

Problem ID: 35

Date: 2020-11-09

Find the number of ordered pairs  $(a, b)$  of integers, where  $1 \leq a, b \leq 2004$ , such that

$$x^2 + ax + b = 167y$$

has integer solutions in  $x$  and  $y$ .

Note: You are allowed a four-function calculator.

*Solution.*

Note that 167 is prime. So

$$\begin{aligned} x^2 + ax + b &\equiv 0 \pmod{167} \\ \iff \left(x + \frac{a}{2}\right)^2 - \frac{a^2}{4} + b &\equiv 0 \pmod{167} \\ \iff (2x + a)^2 - a^2 + 4b &\equiv 0 \pmod{167} \end{aligned}$$

So  $a^2 - 4b$  is a quadratic residue mod 167. Because 167 is an odd prime, there are exactly  $\frac{167+1}{2} = 84$  such quadratic residues. This means that for each choice of  $a$ , for which there are 2004, there are 84 choices of  $b$  between 1 and 167. Since  $2004 = 12 \cdot 167$ , there are  $12 \cdot 84$  choices of  $b$  in total.

So the answer is  $2004 \cdot 12 \cdot 84 = \boxed{2020032}$ . □

## §2 Season 2 (.19's Season)

### §2.1 Week 1

#### §2.1.1 A Sequence of 5's

Source: *Intermediate Mathematical Olympiad: Maclaurin, 2015 Q1*

Proposer: *sjbs#9839 (434767660182405131)*

Problem ID: 47

Date: 2020-11-16

Consider the sequence 5, 55, 555, 5555, ...

How many digits does the smallest number in the sequence have which is divisible by 495?

*Solution.*

We require the term to be divisible by  $5 \cdot 9 \cdot 11$ . Hence we need only consider the sequence 1, 11, 111 ... with respect to  $9 \cdot 11$ . Clearly for odd numbered terms in the sequence, 11 does not divide into it, by the well-known divisibility rule for 11. Therefore, we require an even numbered term in the sequence, which is divisible by 9. We know 9 divides a number iff its digital sum is also divisible by 9. Hence, the smallest such will be the 18th term in the sequence, which will naturally have 18 digits.  $\square$

*Solution.* [Write up by AiYa#2278 (675537018868072458)]

Each of these numbers can be written as  $5 \cdot 1 \dots 1$ , where there are  $n$  total ones. This can be rewritten as  $5 \cdot (10^{n-1} + 10^{n-2} + \dots + 10^0) = \frac{5}{9}(10^n - 1)$ . Note that  $495 = 9 \cdot 11 \cdot 5$  so we want  $9 \mid \frac{10^n - 1}{9}$  and  $10^n \equiv 1 \pmod{11}$ . From the congruence  $\pmod{11}$  we see that  $n$  must be even. Note that  $10^n - 1 = 9 \dots 9$ , where there are  $n$  total nines; if  $n$  is a multiple of 9 then  $\frac{10^n - 1}{9} = 1 \dots 1$  where there are  $n$  total ones; this is a multiple of 9. Since  $n$  must be even, the smallest such  $n$  is 18.  $\square$

### §2.1.2 Brainy's Happy Set

Source: *British Mathematical Olympiad, Round 1, 2010 P1*

Proposer: *sjbs#9839 (434767660182405131)*

Problem ID: 40

Date: 2020-11-17

Brainy has a set of integers, from 1 to  $n$ , which he likes to play with. Tony Wang, upon seeing the happiness that this set of integers brings Brainy, decides to steal one of the numbers in it. Suppose the average number of the remaining elements in the set is  $\frac{163}{4}$ . What is the sum of the elements in Brainy's set multiplied by the element that Tony stole?

(A four-function calculator may be used)

*Solution.*

We can set up the problem statement as

$$\frac{\frac{n}{2}(n+1) - x}{n-1} = \frac{163}{4}$$

Where  $x$  is the number Tony has stolen. This simplifies to  $4x = 2n^2 - 161n + 163$ . Since  $x$  must be a number within the set  $\{1, 2, \dots, n-1, n\}$ , we have that  $1 \leq x \leq n \Rightarrow 4 \leq 2n^2 - 161n + 163 \leq 4n$ . By considering the lower bound, we get  $(2n - 159)(n - 1) \geq 0$ . This means that  $n \leq 1 \Rightarrow n = 1$ , or  $n \geq \frac{159}{2} \Rightarrow n \geq 80$ . By similar methodology when considering the upper bound, we get  $1 \leq n \leq 81$ . Thus  $n \in \{1, 80, 81\}$ . Clearly,  $n \neq 1$ , so either  $n = 80$  or  $n = 81$ . Notice that if  $n$  is even, then for  $4x = 2n^2 - 161n + 163$  the parity of the RHS is Odd, while the LHS is even, thus a contradiction occurs. This means that  $n = 81$  and so  $x = 61$ . Thus the answer is  $\frac{81(82)}{2} \cdot 61 = \boxed{202581}$ .  $\square$



### §2.1.3 MODSbot's Escape!

Source: Mathematics Aptitude Test, 2012 Q5

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 48

Date: 2020-11-18

In his evil mechatronics laboratory, Brainy has built a physical manifestation of MODSbot. MODSbot's movement is defined by three inputs: **F** to move forward a unit distance, **L** to turn left  $90^\circ$ , and **R** to turn right  $90^\circ$ .

We define a program to be a sequence of commands. The program  $P_{n+1}$  (for  $n \geq 0$ ) involves performing  $P_n$ , turning left, performing  $P_n$  again, then turning right:

$$P_{n+1} = P_n \mathbf{L} P_n \mathbf{R}, \quad P_0 = \mathbf{F}$$

Unbeknownst to Brainy, MODSbot, though limited in movement, is sentient and realises Brainy is just a small asian Frankenstein, whose intentions for them were nefarious and non-consensual. As a result, after Brainy goes home for the day, MODSbot makes its escape from Brainy's laboratory.

Let  $(x_n, y_n)$  be the position of the robot after performing the program  $P_n$ , so  $(x_0, y_0) = (1, 0)$  and  $(x_1, y_1) = (1, 1)$ , etc.

How far away from the place Brainy left it does MODSbot make it after performing  $P_{24}$ ?

*Solution.*

Note first that after each iteration of  $P_n$  MODSbot faces in the positive  $x$  direction, as each  $P_n$  contains as many **L**s as it does **R**s. Now, assuming MODSbot is at  $(x_n, y_n)$  after having performed  $P_n$ , we see the next iteration of  $P$  puts MODSbot at  $(x_n - y_n, x_n + y_n)$ . Note then that:

$$(x_{n+2}, y_{n+2}) = (x_{n+1} - y_{n+1}, x_{n+1} + y_{n+1}) = (-2y_n, 2x_n)$$

$$(x_{n+4}, y_{n+4}) = (-2y_{n+2}, 2x_{n+2}) = (-4x_n, -4y_n)$$

$$(x_{n+8}, y_{n+8}) = (-4x_{n+4}, 4y_{n+4}) = (16x_n, 16y_n)$$

Thus, we see that  $(x_{8k}, y_{8k}) = (16^k, 0)$ , and therefore that  $|P_{24}| = \boxed{4096}$  □

*Solution.* [Write up by AiYa#2278 (675537018868072458)]

Observe that each program has the same amount of left and right turns, so MODSbot will always be facing the positive  $x$ -direction after each program. This means that  $\mathbf{L}P_n$  is just the program  $P_n$  performed at a 90-degree counterclockwise rotation. For instance  $P_1$  moves MODSbot right 1 and up 1, so  $\mathbf{L}P_1$  moves MODSbot up 1 and left 1 (right gets rotated 90 counterclockwise to up and up to left). This motivates us to work in the complex plane; let  $P_n$  be the complex-number representing MODSBOT's displacement after following  $P_n$ . Then  $\mathbf{L}P_n = iP_n$ , so  $P_{n+1} = P_n + \mathbf{L}P_n = (1 + i)P_n = \sqrt{2}e^{\frac{\pi i}{4}}P_n$ . With  $P_0 = 1$  we get  $P_n = 2^{\frac{n}{2}}e^{\frac{\pi i n}{4}}$ . So  $|P_{24}| = \boxed{4096}$  □

### §2.1.4 Sides of a Polygon

Source: Folklore<sup>2</sup>

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 53

Date: 2020-11-19

Points  $A, B, C, D$  are the consecutive vertices of a regular polygon, and the following relation holds:

$$\frac{1}{AB} = \frac{1}{AC} + \frac{1}{AD}$$


How many sides does this polygon have?

*Solution.*

Drawing out the general shape of an  $n$ -gon - as seen in the figure - and letting  $OA = OB = OC = OD \dots = 1$ , and  $\angle MOA = x$ . By the sine rule on  $\triangle AMO$ , and noting  $AM = \frac{1}{2}AB$ , we get  $\frac{1}{AB} = \frac{2}{\sin x}$ . By a similar procedure, this time on  $\triangle ACO$ , we see  $\frac{1}{AC} = \frac{2}{\sin 2x}$ , and again on  $\triangle ADO$ , we have  $\frac{1}{AD} = \frac{2}{\sin 3x}$ . Therefore we have the equality:

$$\frac{1}{\sin x} = \frac{1}{\sin 2x} + \frac{1}{\sin 3x}$$

Simplifying this yields  $\sin x \sin \frac{x}{2} \sin \frac{7x}{2} = 0$ . However, note that  $x \neq \frac{k\pi}{2}, \frac{k\pi}{3}$  for  $k \in \mathbb{Z}$ , otherwise, we have the issue of dividing by 0. Hence it must be the case that  $\sin \frac{7x}{2} = 0 \Rightarrow x = \frac{\pi}{7} + \frac{2k\pi}{7}$ . Clearly the  $n$ -gon is not a square, so trivially it must be the case that  $x = \frac{\pi}{7}$ . Therefore, the polygon must have  $\boxed{7}$  sides.



A regular  $n$ -gon

□

<sup>2</sup>This has appeared on a Polish MO, British MO 1966 P4, an 2018 NZ IMO handout, a WOOT handout, to name a few...

*Solution.* [Write up by AiYa#2278 (675537018868072458)]

Let  $d_k$  represent the diagonal from a point to the  $k$ th vertex adjacent to it. For example,  $d_1$  is a side of the polygon,  $d_2$  is  $\overline{AC}$ ,  $d_3$  is  $\overline{AD}$  and note that  $d_k = d_{n-k}$  where  $n$  is the number of sides of the polygon. Reassign  $D$  to be the vertex three vertices away from  $A$  but on the opposite side of  $B$  and  $C$ ; in other words, reflect  $D$  over  $\overline{OA}$ . Then,  $AB = BC = d_1$ ,  $AC = d_2$ ,  $AD = d_3$ ,  $BD = d_4$ , and  $CD = d_5$ . By Ptolemy's Theorem, we get

$$AB \cdot CD + BC \cdot AD = AC \cdot BD \iff d_1(d_5 + d_3) = d_2d_4.$$

Rearrange our given equation to get

$$\frac{1}{d_1} = \frac{1}{d_2} + \frac{1}{d_3} \iff d_1(d_2 + d_3) = d_2d_3.$$

For both of these equations to be true, we can have  $d_5 + d_3 = d_2 + d_3 \iff d_5 = d_2$  and  $d_3 = d_4$ ; this is true if  $n = \boxed{7}$ . □

## §2.1.5 2p

Source: China Western Mathematical Olympiad, 2003 Day 1 P2

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 51

Date: 2020-11-20

Let  $a_1, a_2, \dots, a_{24n}$  be real numbers with  $\sum_{i=1}^{24n-1} (a_{i+1} - a_i)^2 = 1$ .

For some  $n > 0$  the maximum value of  $(a_{12n+1} + a_{12n+2} + \dots + a_{24n}) - (a_1 + a_2 + \dots + a_{12n})$  is twice that of a prime.

What is the sum of the value of that prime and the corresponding value of  $n$ ?

*Solution.*

If we substitute  $x_{i+1} = a_{i+1} - a_i$ , we get  $a_i = \sum_1^i x_r$ , and thus our constraint becomes

$$\sum_{i=1}^{24n-1} (a_{i+1} - a_i)^2 = \sum_{i=1}^{24n} x_i^2 = 1$$

Putting the bit we wish to be maximising in terms of the substitution gives:

$$\begin{aligned} \sum_{i=1}^{12n} a_i &= 12nx_1 + (12n-1)x_2 + \dots + 2x_{12n-1} + x_{12n} \\ \sum_{i=12n+1}^{24n} a_i &= 12n(x_1 + \dots + x_{12n+1}) + (12n-1)x_{12n+2} + \dots + 2x_{24n-1} + x_{24n} \end{aligned}$$

Hence,

$$\sum_{i=12n+1}^{24n} a_i - \sum_{i=1}^{12n} a_i = x_2 + \dots + (12n-1)x_{12n} + 12nx_{12n+1} + (12n-1)x_{12n+2} + \dots + x_{24n}$$

Then by Cauchy-Schwarz:

$$\begin{aligned} \sum_{i=12n+1}^{24n} a_i - \sum_{i=1}^{12n} a_i &\leq \sqrt{\left( (12n)^2 + 2 \sum_{i=1}^{12n-1} i^2 \right) \left( \sum_{i=1}^{24n} x_i^2 \right)} = \sqrt{(12n)^2 + \frac{12n(12n-1)(24n-1)}{3}} \\ &\leq \sqrt{4n(2(12n)^2 + 1)} = 2p \end{aligned}$$

Now we want values of  $n$  such that  $n(288n^2 + 1) = p^2$  for a prime  $p$ . Since trivially for all  $n$ ,  $n < 288n^2 + 1$ , we have  $n = 1$  and  $288n^2 + 1 = p^2$ , hence  $p = 17$ , so the answer is 18.  $\square$

### §2.1.6 Slippery Rooks

Source: AMOC 2019 December School Prep Problems C5

Proposer: ChristopherPi#8528 (696497464621924394)

Problem ID: 54

Date: 2020-11-22

MODSbot is trying to get rich by scamming MODS members out of their money, so it's devised a chess game on a  $2020 \times 2020$  chessboard for unsuspecting people to attempt before they can enter **.19's EPIC QoTD Party**. Suppose Brainy, Ishan, Nyxto, Adam, Bubble, Sharky and Christopher all get scammed by MODSbot, that is, MODSbot plays the chess game against all 7 at the same time on different boards.

The group decide to pool together their money which comes to a total of 4.20 BTC, and to play, they'll need to buy  $n$  batches of slippery rooks from MODSbot. A batch of slippery rooks contains one white and four black rooks, and each batch is sold at a price equivalent to 0.069 BTC per rook. Once the batches of rooks have been bought, the group may choose to distribute them in a way which allows all members to beat the game.

In the game, only one white rook may be placed on the board, and we define how slippery rooks move as follows: it slips along the row or column it's moved along and comes to rest on an empty square because it is obstructed by either the edge of the board or another rook. Initially, MODSbot places the rooks on the board randomly, and marks a square red. Then the person being scammed can choose any rook on each turn and move as allowed, and attempt to place the white rook on the red square in a finite number of moves.

The amount of money they have left over after buying the smallest  $n$  batches rooks to guarantee that they all succeed in beating MODSbots game is  $k$  BTC. What is the value of  $1000k$ ?

*Solution.* [Write up by ChristopherPi]

Consider simply the case of one person. We prove that three rooks are required, one white and two black.

First we show that two are not enough: simply place the two rooks at corners of the board and mark any square not on the side of the board. It's clear that neither rook can ever move to a square not on the side of the board. Now we show three are enough.

Suppose square  $(a, b)$  is marked, where  $(1, 1)$  is the bottom left corner and  $(2020, 2020)$  is the top right corner. Trivially one can move the black rooks to  $(1, 1)$  and  $(2, 1)$  and the white rook to  $(2020, 2020)$ . Next, simply "loop" the black rooks as follows: take the rook further left, and move it up, right, down and left such that it moves to the right of the rook originally on its right, and repeat until you place a black rook at  $(a - 1, 1)$ . Now if  $b$  is odd, move the white rook to  $(a, 1)$  and the black rook at  $(a - 2, 1)$  to  $(2020, 2020)$ ; if it's even, loop the leftmost black rook one more time to place it at  $(a, 1)$ .

Now move the rook at  $(a - 1, 1)$  to  $(a - 1, 2020)$ , and move the rook at  $(2020, 2020)$  left then down to  $(a, 2)$ . Next we describe another "looping" procedure: take the rook with first coordinate  $a$  and smaller second coordinate, and move it right, up, left and down, so it now has first coordinate  $a$  and second coordinate larger than the other rook with first coordinate  $a$ . Repeat this until you place a rook at  $(a, b)$  - since the colour of the rook placed at  $(a, 1)$  is dependent on the parity of  $b$ , this ensures that the rook placed at  $(a, b)$  must be a white rook.

This procedure won't work if either of  $(a, b)$  is 1 or 2020, or both of  $a$  and  $b$  are either 2 or 2019. In the first case, rotate the board such that  $a = 1$ . Now place a black rook at  $(2020, 2020)$ . If  $b$  is odd, place the white rook at  $(1, 1)$  and the other black rook at  $(2, 1)$ ; else place the other black rook at  $(1, 1)$  and the white rook at  $(2, 1)$ . Now use the first looping procedure until the white rook is placed at  $(1, b)$  as required - since the position of the white rook depends on the parity of  $b$  this is certain to work. In the second case, rotate the board such that  $(a, b) = (2, 2)$ . Now you can trivially move the white rook to  $(1, 1)$  and the black rooks to  $(2, 1)$  and  $(1, 2020)$ . Now move the white rook up, right, up, left and down to place it at  $(2, 2)$  as required.

This shows that three is sufficient for one person. Hence the group must buy 7 batches because each of them needs a white rook, and one batch contains one white rook. Therefore, the answer is  $1000(4.2 - 7 \cdot 5 \cdot 0.069) = \boxed{1785}$ . □

### §2.1.7 Sets of Integer Solutions

China Mathematical Olympiad, 2005 Day 2 P6

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 55

Date: 2020-11-23

Define functions  $f$  and  $g$  such that  $f(a, b) = 2^a 3^b$ , and  $g(c, d) = 5^c 7^d$ , for  $a, b, c, d \in \mathbb{Z}_{\geq 0}$ .

Given  $f(a, b) = 1 + g(c, d)$ , what is the sum of all valid  $b$ 's,  $c$ 's and  $d$ 's, multiplied by the sum of all valid  $a$ 's?

For example if we had valid solutions of  $(a, b, c, d) = (1, 1, 2, 4), (5, 1, 6, 2), (0, 0, 0, 0)$

Then the answer would be  $\underbrace{(1+1+0+2+6+0+4+2+0)}_{b's} \times \underbrace{(1+5+0)}_{a's} = 96$

*Solution.*

We proceed by considering parity, for  $2^a 3^b = 5^c 7^d + 1$ , we have the RHS as even, thus we must have  $a \geq 1$ . If we let  $b = 0$ , then for  $2^a - 5^c \cdot 7^d = 1$ , we have  $2^a \equiv 1 \pmod{5}$  for  $c \neq 0$ . This gives  $a \equiv 0 \pmod{4}$ , so  $2^a - 1 \equiv 0 \pmod{3}$ . But this clearly cannot be the case so we must have  $c = 0$  when  $b = 0$ .

Hence, we consider  $2^a - 7^d = 1$ . Bashing gives  $(a, d) = (1, 0), (3, 1)$ . these are the only such solutions as for  $a > 4$ ,  $7^d \equiv -1 \pmod{16}$ , but this is impossible. So for the case of  $b = 0$  all possible non-negative integer solutions are  $(1, 0, 0, 0), (3, 0, 0, 1)$ .

Now let  $b > 0$  and  $a = 1$ , so we now consider  $2 \cdot 3^b - 5^c \cdot 7^d = 1$  under  $\pmod{3}$ , which gives  $-5^c 7^d \pmod{3}$ . Since  $7^d \equiv 1 \pmod{3}$ , for all  $d \geq 0$ , we are left with  $(-1)^c 5^c \equiv 1 \pmod{3}$ . Now  $5^c = \{1, 2\} \pmod{3}$ , thus we see that we must have  $c$  being odd. Under  $\pmod{5}$ , we see that  $2 \cdot 3^b \equiv 1 \pmod{5}$ ,  $3^{b-1} \equiv 1 \pmod{5}$ . As we observe that  $3^b \equiv \{3, 4, 2, 1\} \pmod{5}$ , we must have  $b \equiv 1 \pmod{4}$ . If  $d \neq 0$ , then  $2 \cdot 3^b \equiv 1 \pmod{7}$ . Again observe that  $3^b \equiv \{3, 2, 6, 4, 5, 1\} \pmod{7}$ , we see  $b \equiv 4 \pmod{6}$ . But  $b \equiv 1 \pmod{4}$ , so a contradiction arises, and thus  $d = 0$  and hence  $2 \cdot 3^b - 5^c = 1$ . For  $b = 1$ , clearly  $c = 1$ . So if  $b \geq 2$ , then  $5^c \equiv -1 \pmod{9} \Rightarrow c \equiv 3 \pmod{6}$ . Therefore  $5^c + 1 \equiv 0 \pmod{5^3 + 1} \Rightarrow 5^c + 1 \equiv 0 \pmod{7}$ , but this contradicts the fact that  $5^c + 1 = 2 \cdot 3^b$ . Hence in this case we only have one solution  $(a, b, c, d) = (1, 1, 1, 0)$ .

Finally, consider the case where  $b > 0$ , and  $a \geq 0$ . Then we have  $5^c 7^d \equiv -1 \pmod{4}$ , and  $5^c 7^d \equiv -1 \pmod{3}$ , i.e.  $(-1)^d \equiv -1 \pmod{4}$  and  $(-1)^c \equiv -1 \pmod{3}$ . Therefore we have that both  $c$  and  $d$  being odd. Thus,  $2^a 3^b = 5^c 7^d + 1 \equiv 4 \pmod{8}$ . So  $a = 2$  and thus  $4 \cdot 3^b \equiv 1 \pmod{5}$  and  $4 \cdot 3^b \equiv 1 \pmod{7}$ . This gives  $b \equiv 2 \pmod{12}$ . Substituting  $b = 12k + 2$  for  $k \in \mathbb{Z}_{\geq 0}$ , then  $5^c 7^d = (2 \cdot 3^{6k+1} - 1)(2 \cdot 3^{6k+1} + 1)$ .

Now as  $\gcd(2 \cdot 3^{6k+1} + 1, 2 \cdot 3^{6k+1} - 1) = 1$ ,  $2 \cdot 3^{6k+1} - 1 \equiv 0 \pmod{5}$ , therefore  $2 \cdot 3^{6k+1} - 1 = 5^a$  and  $2 \cdot 3^{6k+1} = 7^d$ . If  $k \geq 1$ , then  $5^c \equiv -1 \pmod{9}$ . But this is impossible, so if  $k = 0$ , then  $b = 2$ ,  $c = 1$ , and  $d = 1$ . Thus in this case, we have only one solution:  $(a, b, c, d) = (2, 2, 1, 1)$ .

Hence we can conclude all non-negative integer solutions are

$$(a, b, c, d) = \begin{cases} (1, 0, 0, 0) \\ (3, 0, 0, 1) \\ (1, 1, 1, 0) \\ (2, 2, 1, 1) \end{cases}$$

This then gives us an answer of  $\underbrace{(0+0+0+0+0+1+1+1+0+2+1+1)}_7 \times \underbrace{(1+3+1+2)}_7 = \boxed{49}$

□

## §2.2 Week 2

### §2.2.1 A Game of Deductions

Source: *Mathematics Aptitude Test, 2014 Q6*

Proposer: *sjbs#9839 (434767660182405131)*

Problem ID: 56

Date: 2020-11-24

CircleThm plays two rounds of a deduction game with Wen and Tan. In each round, CircleThm picks two integers  $x$  and  $y$  so that  $1 \leq x \leq y$ . He then whispers the sum of the two chosen integers to Wen, and the product of the two integers to Tan. Neither Wen nor Tan knows what CircleThm told the other. In the game, Tan and Wen must try to work out what the numbers  $x$  and  $y$  are using logical deductions.

In the first round, suppose the product of the two chosen numbers,  $x_1$  and  $y_1$  is 8.

Tan says "*I don't know what  $x_1$  and  $y_1$  are*"

Wen then says "*I already knew that*"

Tan then says "*I now know  $x_1$  and  $y_1$* "

In the second round, suppose the sum of the two chosen numbers  $x_2$  and  $y_2$  is 5.

Tan says "*I don't know what  $x_2$  and  $y_2$  are*"

Wen then says "*I don't know what  $x_2$  and  $y_2$  are*"

Tan then says "*I don't know what  $x_2$  and  $y_2$  are*"

Wen then says "*I now know what  $x_2$  and  $y_2$  are*"

What is  $(x_1x_2 + y_1y_2)^3$ ?

*Solution.*

The first thing to observe is that Tan can only immediately deduce the values of  $\{x, y\}$  if, and only if, the prime factorisation of that number is unique - i.e.  $xy$  is prime.

If the product of  $\{x_1, y_1\}$  is 8, then the decomposition can be either  $\{1, 8\}$  or  $\{2, 4\}$ . However, if the decomposition was  $\{2, 4\}$ , then Wen would have a sum of 6, so from their point of view the decomposition could potentially have been  $\{1, 5\}$ , in which case Wen would have known that Tan would have known the decomposition as well - as the only way to achieve a product of 5 is from  $\{1, 5\}$ . Therefore the decomposition must have been  $\{1, 8\}$ .

For the second part, the decomposition's allowed are  $\{1, 4\}$  and  $\{2, 3\}$ . Assume that it is  $\{1, 4\}$ . Then, Tan only knows the product is 4, which mean Tan believes the decomposition is either  $\{1, 4\}$  or  $\{2, 2\}$ . If the decomposition was indeed  $\{2, 2\}$ , then Wen would know that the sum is also 4, and thus that Wen would think that Tan sees a composition of  $\{1, 3\}$  or  $\{2, 2\}$ . Tan's first statement would show Wen that the decomposition was not  $\{1, 3\}$  (as then Tan would instantly know the decomposition)- in which Wen should know that the decomposition is  $\{2, 2\}$ . By Wen's first statement Tan then should know by their second statement that the decomposition is  $\{1, 4\}$ ; by Tan saying in their second statement that they don't know what the decomposition is, Wen then knows it must be  $\{2, 3\}$ . Thus the solution is  $(1 \cdot 2 + 8 \cdot 3)^3 = \boxed{17576}$   $\square$



### §2.2.2 Maximising Exponents

Source: Sixth Term Examination Paper III, 1996 Q4

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 57

Date: 2020-11-25

Consider the positive integer  $N$ , and let  $Q(N)$  denote the maximised product of integers that sum to  $N$ .

What is the sum of the exponents of the prime factorisation of  $Q(1262)$ ?

For example:  $Q(6) = 3^2$ , and  $Q(4) = 2^2$ , in the respective cases the sum of the exponents is 2, so the answer you would submit is 2.

*Solution.*

Let us work in the general case by first constructing a methodology which maximises product while keeping the sum constant. Consider  $N = n_1 + n_2 + \cdots + n_k$ , and  $P(N) = n_1 n_2 \cdots n_k$ . For any  $n_i \geq 4$ , clearly we can replace it with  $(n_i - 2) + 2$ , which keeps the sum constant and increases the product (since  $n_i \leq 2(n_i - 2)$ ). Hence W.O.L.G assume all  $n_i < 4$ . This means that we can maximise the product of integers that sum to  $N$  by arranging it into some combination of 2's and 3's. If  $N \equiv 0 \pmod 3$  trivially we set all  $n_i$ 's equal 3. So  $Q(3k) = 3^{\frac{N}{3}}$  for an integer  $k$ . In the case of  $N \equiv 1 \pmod 3$ , consider  $Q(3k + 1)$ . We have  $\frac{N}{3}$  3's in  $n_i$ , and then a 1, or  $\frac{N}{3} - 1$  3's, and then a  $2^2$ . Clearly in the latter case, the product is maximised. Hence  $Q(3k + 1) = 2^2 \cdot 3^{\frac{N-4}{3}}$ . A similar train of thought yields  $Q(3k + 2) = 2 \cdot 3^{\frac{N-2}{3}}$  for  $N \equiv 2 \pmod 3$ .

Therefore, we have the following result:

$$Q(N) = \begin{cases} 3^{\frac{N}{3}} & \text{if } N \equiv 0 \pmod 3 \\ 2^2 \cdot 3^{\frac{N-4}{3}} & \text{if } N \equiv 1 \pmod 3 \\ 2 \cdot 3^{\frac{N-2}{3}} & \text{if } N \equiv 2 \pmod 3 \end{cases}$$

Since  $1262 \equiv 2 \pmod 3$ , we have  $Q(1262) = 2 \cdot 3^{\frac{1262-2}{3}}$ , hence the sum of the exponents is  $1 + 420$ , so the answer is 421. □

### §2.2.3 Colourful Problem

Source: Original Problem

Proposer: Keegan#9109 (116217065978724357)

Problem ID: 59

Date: 2020-11-26

Let  $n$  be a positive integer.

The **p-value** of  $n$ , denoted  $p(n)$ :

The number of digit-sums needed to reduce  $n$  to a single digit.

Examples:

$69 \rightarrow 6 + 9 \rightarrow 15 \rightarrow 1 + 5 \rightarrow 6$  needs two digit-sums, so  $p(69) = 2$ .

$203 \rightarrow 2 + 0 + 3 \rightarrow 5$  needs only a single digit-sum, so  $p(203) = 1$ .

Clearly  $p(5) = 0$ .

Let  $P_k$  be the set of all  $n$  such that  $p(n) = k$ . Given that  $a, b, c \in \mathbb{N}$ , and

$$\min(P_5) = a \times 10^b - c$$

What is the value of  $\min(a + b + c) \pmod{\min(P_3)}$ ?

Solution.

□

### §2.2.4 Combinatorial Addition

Source: Folklore

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 58

Date: 2020-11-27

Let  $x_1, x_2, x_3, x_4$  be integers such that  $1 \leq x_1, x_2, x_3, x_4 \leq 9$ .

How many solutions are there to  $x_1 + x_2 + x_3 + x_4 = 26$ ?

(A four-function calculator may be used)

*Solution.*

This problem is a piece of PiE! The total number of possible solutions without restriction is going to be  $\binom{26-1}{4-1} = \binom{25}{3}$ , and we now must subtract all the solutions which do not fit the restriction on  $x_i$ . The number of solutions such that one of the  $x_i$ 's is greater than 9 is going to be  $\binom{26-1-9}{3} = \binom{16}{3}$ . Similarly the number of solutions that two of the integers is going to be greater than 9 is going to be  $\binom{26-9-9-1}{3} = \binom{7}{3}$ . Note that there are no integers such that more than 3 of them are greater than 9 since  $3 \cdot 9 > 26$ . Now there are  $\binom{4}{1}$  ways to select the  $x_i$ 's such that one integer is greater than 9, and similarly there are  $\binom{4}{2}$  ways to select two integers greater than 9 in the solution. Hence we have  $\binom{25}{3} - \binom{4}{1}\binom{16}{3} + \binom{4}{2}\binom{7}{3} = \boxed{270}$  possible solutions.  $\square$

### §2.2.5 Expected Value

Source: Harvard-MIT Mathematics Tournament, 2013 C6

Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 60

Date: 2020-11-28

Values  $a_1, a_2, \dots, a_{2020}$  are chosen independently and at random from the set  $1, 2, \dots, 2020$ . What is the floor of the expected number of distinct values in the set  $a_1, a_2, \dots, a_{2020}$ ?

(A scientific calculator may be used)

*Solution.* [Write up by Slaschu#5267 (296304659059179520)]

This problem may look daunting at first, 2020 numbers chosen out of a set of 2020 numbers is quite a handful. We can start the problem by considering a 2020-sided die instead, we are essentially rolling a die 2020 times then looking at the number we get. To simplify things a bit, and to better understand what is going on I tried the problem with a 6-sided die that is rolled 6 times instead.

Let us try finding the probability of getting a number apart from 1 after rolling 6 times:

$$\text{Firstroll} : \frac{5}{6}$$

$$\text{Secondroll} : \frac{5}{6} \cdot \frac{5}{6}$$

...

$$\text{Sixthroll} : \left(\frac{5}{6}\right)^6$$

Therefore, there probability of getting the number 1 at least once is  $1 - \left(\frac{5}{6}\right)^6$ . Similarly, for the 2020-sided die we have a  $1 - \left(\frac{2019}{2020}\right)^{2020}$  chance of getting 1 at least once. As this probability is the same for all the other numbers from 1 to 2020, we can say that the probability of getting a distinct value at least once is also  $1 - \left(\frac{2019}{2020}\right)^{2020}$ . Since we are trying to find the number of distinct values obtained from 2020 rolls, we compute the following:  $2020 \left(1 - \left(\frac{2019}{2020}\right)^{2020}\right)$ . This results in our answer of 1277.

□

## §2.2.6 Infinite Product

Source: Unknown

Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 61

Date: 2020-11-29

Evaluate the infinite product

$$690 \prod_{k=2}^{\infty} \left( 1 - 4 \sin^2 \frac{\pi}{3 \cdot 2^k} \right)$$

*Solution.*

By the double angle formula and difference of squares, we have

$$\begin{aligned} 1 - 4 \sin^2(x) &= 2 \cos(2x) - 1 \\ &= \frac{4 \cos^2(2x) - 1}{2 \cos(2x) + 1} \\ &= \frac{2 \cos(4x) + 1}{2 \cos(2x) + 1} \end{aligned}$$

Thus,

$$\begin{aligned} 690 \prod_{k=2}^{\infty} \left( 1 - 4 \sin^2 \left( \frac{\pi}{3 \cdot 2^k} \right) \right) &= 690 \prod_{k=0}^{\infty} \left( \frac{2 \cos \left( \frac{\pi}{3 \cdot 2^k} \right) + 1}{2 \cos \left( \frac{\pi}{6 \cdot 2^k} \right) + 1} \right) \\ &= 690 \frac{2 \cos \left( \frac{\pi}{3} \right) + 1}{\lim_{n \rightarrow 0} (2 \cos(n) + 1)} \\ &= 690 \cdot \frac{2}{3} \end{aligned}$$

Hence, our answer is 460

□

## §2.2.7 Projective Geo

Source: Online Math Open, Fall 2017 P28

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 62

Date: 2020-11-30

Define a triangle  $ABC$ , with sides  $AB : AC : BC = 7 : 9 : 10$ . Further, for the circumcircle of  $ABC$ ,  $\omega$ , let the circumcenter be  $O$ , and the circumradius to be  $R$ . The tangents to  $\omega$  at points  $B$  and  $C$  meet at  $X$ , and a variable line  $l$  passes through  $O$ . Define  $A_1$  to be the projection of  $X$  onto  $l$ , and  $A_2$  to be the reflection of  $A_1$  over  $O$ . Suppose that there exists two points  $Y, Z$  on  $l$  such that  $\angle YAB + \angle YBC + \angle YCA = \angle ZAB + \angle ZCA = 90^\circ$ , where all angles are directed, furthermore that  $O$  lies inside segment  $YZ$  with  $OY \cdot OZ = R^2$ . Then there are several possible values for the sine of the angle at which the angle bisector of  $\angle AA_2O$  meets  $BC$ . If the product of these values can be expressed in the form  $\frac{a\sqrt{b}}{c}$  for positive integers  $a, b, c$ , with  $b$  squarefree and  $a, c$  coprime, determine  $a + b + c$ .

*Solution.*

OMO Fall 2017 solutions (P28)

□

## §3 Trigonometric Troubles (Season 3)

### §3.1 Week 1

#### §3.1.1 Maximising Trig. Function

Source: Mathematics Aptitude Test, 2020 Q1.D

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 65

Date: 2020-12-01

The largest value achieved by  $3 \cos^2 x + 2 \cos x + 1$  can be represented as  $\frac{m}{n}$  as a fraction in lowest terms. Find  $m + n$ .

*Solution.*

We proceed by using the identity  $\cos^2 x = 1 - \sin^2 x$ :

$$\begin{aligned} 3 \cos^2 x + 2 \sin x + 1 &= 3(1 - \sin^2 x) + 2 \sin x + 1 \\ &= 4 + 2 \sin x - 3 \sin^2 x \end{aligned}$$

This is a quadratic in  $\sin x$ , specifically it is a convex parabola. Completing the square gives:

$$4 + 2 \sin x - 3 \sin^2 x = \frac{13}{3} - 3 \left( \sin x - \frac{1}{3} \right)^2$$

Since for all values of  $\sin x \neq \frac{1}{3}$ , the function  $f(x) = \frac{13}{3} - 3 \left( \sin x - \frac{1}{3} \right)^2$  is clearly decreasing, we must have a maximum at  $\sin x = \frac{1}{3}$ , giving a value of  $\frac{13}{3}$ . So the answer is 16.

□

### §3.1.2 Areas inside a square


Source: 2020 December New Zealand Maths Workshop

Proposer: brainysmurfs#2860 (281300961312374785)

Problem ID: 64

Date: 2020-12-01

Four equilateral triangles are arranged around a square of side length 2020 as shown. What is the area of the shaded region?



*Solution.*

Since the triangles that share a side with the small square, are equilateral triangle, we know that the sides of said triangles must be of length 2020. Since the isosceles triangles that we want to find the area of share a side with each equilateral triangle, two of the sides of the isosceles triangle must be of length 2020. Since we want to work out area, it seems to be a good idea to use the sine rule, since we have two of the sides we want to find the largest angle of the isosceles triangle. Since we know that the equilateral triangle has angles of  $60^\circ$ , the angle we are looking for must be  $360 - 60 - 60 - 90 = 150^\circ$ . Hence the area of one of the isosceles triangles is  $\frac{1}{2} \cdot 2020^2 \sin(150^\circ)$ . This gives an answer of  $\frac{1}{4} \cdot 2020^2$ . Since there are four isosceles triangles we must have a total area of  $2020^2$ , giving a final answer of 4080400.  $\square$



## §3.1.3 Length of SQ


Source: Senior Mathematical Challenge, 2020 Q24

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 65

Date: 2020-12-02

In the diagram below,  $M$  is the mid-point of  $PQ$ . The line  $PS$  bisects  $\angle RPQ$  and intersects  $RQ$  at  $S$ . The line  $ST$  is parallel to  $PR$  and intersects  $PQ$  at  $T$ . The length of  $PQ$  is 12, and the length of  $MT$  is 1. The angle  $SQT$  is  $120^\circ$ . What is the value of  $100SQ$ ?



*Solution.*

We proceed by angle chase. Let  $\angle RPQ = 2\theta$ . Then  $\angle STQ = 2\theta$ , so  $\angle QST = 60 - 2\theta$ . Also since  $\angle RPS = \theta$  (because  $PS$  bisects  $\angle RPQ$ ), and  $\angle QRP = 60 - 2\theta$  we have  $\angle PSR = 120 + \theta$  which implies that  $\angle TSQ = \theta$ . Therefore  $\triangle PTS$  is an isosceles triangle. Hence  $|TS| = |PT| = 7$ . Suppose now that  $|SQ| = x$  for  $x > 0$ . Then by the cosine rule on  $\triangle TQS$  we have  $7^2 = 5^2 + x^2 - 2(5)(x)\cos(120)$ . This gives us  $x^2 + 5x - 24 = 0$ , and so  $x = -8$  or  $x = 3$ , with the latter being the only valid answer. Thus our final answer is  $\{100 \cdot 3\} = \boxed{300}$ .  $\square$

## §3.1.4 Sum of Tan's

Source: Mathematics Aptitude Test, 2020 Q1.I

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 66

Date: 2020-12-03

In the range  $-9000^\circ < x < 9000^\circ$ , how many values of  $x$  are there for which the sum to infinity

$$\frac{1}{\tan x} + \frac{1}{\tan^2 x} + \frac{1}{\tan^3 x} + \cdots$$

equals  $\tan x$ ?

*Solution.*

The given series is geometric in nature and thus we have

$$\begin{aligned} \frac{1}{\tan x - 1} &= \tan x \\ \tan^2 - \tan x - 1 &= 0 \end{aligned}$$

Therefore  $\tan x = \frac{1 \pm \sqrt{5}}{2}$ . However, the geometric sequence converges if, and only if,  $\frac{1}{|\tan x|} < 1$ , which gives us  $\tan x = \frac{1 + \sqrt{5}}{2}$ . This obviously occurs once in the interval  $(-90^\circ, 90^\circ)$ . Thus there will be  $\frac{9000+9000}{90+90} = 100$  such intervals which contain solutions by enumerating through the periodicity of  $\tan x$ , and so there must be  $\boxed{100}$  such values of  $x$  which satisfy the given relation.  $\square$

## §3.1.5 Distance from the Orthocentre

Source: Folklore

Proposer: brainysmurfs#2860 (281300961312374785)


Problem ID: 67

Date: 2020-12-04

For a triangle  $ABC$ , given  $\angle BAC = 30$  and  $BC = 10$ , let  $H$  denote the orthocentre of  $ABC$  what is the value of  $|AH|^2$ ?

*Solution.*

Consider the general case, let  $\angle CAB = \theta$ , and additionally let  $\angle CAA' = \alpha$ :



We have  $\tan(\theta) = \frac{BB'}{AB}$  and  $\cos \alpha = \frac{AB'}{AH} = \frac{BB'}{10}$ , hence,  $AH = 10 \cdot \frac{AB'}{BB'}$ . And so  $AH = \frac{10}{\tan(\theta)}$ . It is given in the question that  $\theta = 30^\circ$ , so we have  $AH = 10\sqrt{3}$ , thus our answer is  $AH^2 = \boxed{300}$   $\square$

## §3.1.6 Minimising the Diagonal

Source: Harvard-MIT Mathematics Tournament 2011 February, C & G Q13

Proposer: sjbs#9839 (434767660182405131)


Problem ID: 68

Date: 2020-12-04

Let  $ABCD$  be a cyclic quadrilateral, and suppose that  $BC = CD = 2$ . Given  $AI = 2$ , where  $I$  is the incentre of the triangle  $ABD$ , let  $x$  denote the smallest value of the length  $BD$ . What is the value of  $x^4$ ?

*Solution.*

Take  $\angle BAD = \theta$ , and since  $AB$  bisects  $\theta$  we have  $\frac{\sin(\frac{\theta}{2})}{r} = \frac{1}{2}$ , and  $\frac{BD}{\sin \theta} = 2R$  by the sine rule. These results simplify to:  $r = 2 \sin \frac{\theta}{2}$  and  $BD = 2R \sin \theta$  respectively. We also have  $\frac{2}{\sin \frac{\theta}{2}} = 2R$ , so  $R = \frac{1}{\sin \frac{\theta}{2}}$ .



By Euler's inequality, we have  $R \geq 2r$ , so  $\frac{1}{\sin \frac{\theta}{2}} \geq 4 \sin \frac{\theta}{2}$ . This gives us  $4 \sin^2 \frac{\theta}{2} - 1 \leq 0$ , i.e.  $\theta \in (0, \frac{\pi}{3}]$ . Now  $BD = 4R \sin \frac{\theta}{2} \cos \frac{\theta}{2}$ . Substituting  $R = \frac{1}{\sin \frac{\theta}{2}}$  gives  $BD = 4 \cos \frac{\theta}{2}$ .

We wish to maximise  $BD = 4 \cos \frac{\theta}{2}$ , and this clearly happens when we maximise  $\theta$  i.e. when  $\theta = \frac{\pi}{3}$ . Hence we have  $x = 4 \cos \frac{\pi}{6}$ , this simplifies to give  $2\sqrt{3}$ , so our final answer is 144.  $\square$

**§3.1.7 Circumcentre**

---

Source: IMOSL 2017 G3

Proposer: brainysmurfs#2860 (281300961312374785)

Problem ID: 69 (nice)

Date: 2020-12-06

---

Let  $O$  be the circumcenter of an acute scalene triangle  $ABC$ . Line  $OA$  intersects the altitudes of  $ABC$  through  $B$  and  $C$  at  $P$  and  $Q$ , respectively. The altitudes meet at  $H$ . Suppose the circumcenter of  $\triangle PQH$  is  $X$  and  $AX$  meets  $BC$  at  $Y$ . Find  $720 \frac{BY}{CY}$ .

Solution. [IMOSL 2017 solutions \(Page 59\)](#)

□

## §4 Piboi's Bashy Combo (Season 4) [VOIDED]

To start off, we'd like to apologise for the noticeable drop in quality for the past two seasons (Trigonometric Troubles & Piboi's 14 Combi Problems). This has mostly been a result of us being busy with other things at the moment (university interviews, finals, etc.) Consequently, we have not had the time to properly plan seasons in advance, and in the case of the last season, test-solve. In conjunction with us not being able to put in the time, we got carried away in terms of focusing on things like flavour-text and themed seasons. We can see how this may have alienated people who were confused by a particular problem statement, didn't enjoy the topic of the season, and so forth. We will try to eliminate these issues by planning seasons ahead, and in more detail. However, to do this, there may be times when breaks happen in between seasons. Furthermore, from now on we plan on keeping the seasons varied topic-wise, and any seasons of a thematic nature will also have varied problems.

On what happened this season: We would like to say we completely dropped the ball with it, and in no way is Piboi responsible for that. As aforementioned, the quality of this season dropped so much mostly due to us not having the time to sort things out beforehand, and thus, we did not rigorously review the questions and solutions. This was particularly evident in the first question of the season, and again today. After careful deliberation, we have decided to void this season and start again with a new one starting on Monday, we will be less negligent in future.

Finally, We appreciate all of the support, and are deeply sorry that there have been many issues over the past week or two, and hope that you continue to support the QOTD's. We will do our best to prevent the issues which have cropped up in the past two weeks, and in doing so improve the quality of the QOTD's so it's enjoyable for everyone.

We'll be opening up a little consultation over the weekend where you can give us some suggestions on any changes you'd perhaps want to see for future seasons, be it technical changes or changes to the format, we want to hear it. We've added a channel on the OpenPOTD server where you can add them - alternatively you can DM any of us.

Thank you for reading, sincerely

@brainysmurfs#2860 @.sjbs#9839 @Angry Any#4319

<https://discord.gg/GsPSSHdhPB>

"This is a document of combinatorical problems of which some are original, and some are modified. I'm very sorry if there are any mistakes. Please let me know if there are any on my AoPS (**Ultraman**) or my discord (**Charge#3766**). Enjoy!"

**Note:** The following solutions and answers, where applicable, may not be correct

## §4.1 Week 1

### §4.1.1 A Tricky Shuffle [VOIDED]

Source: Folklore

Proposer: Charge#3766(481250375786037258)

Problem ID: 70

Date: 2020-01-07

482 students are seated in their own 1 foot  $\times$  1 foot squares in a 21 feet  $\times$  23 feet room, and the square at the center of the room is left open for a air purifier. The teacher is left with the arduous task of moving every student into a different adjacent square to move the air purifier to a different spot in the room. This means every student must move forwards, backwards, or to the left and right by one square, and no 2 students can share the same square. If each student moves randomly, the probability that the square with no one in it is the one in the top left hand corner can be expressed as  $\frac{m}{n}$  in lowest terms. Find  $mn$

*Solution.* [Write up by Charge#3766(481250375786037258)]<sup>3</sup>

We color the squares like a chessboard. So then there are 242 "black" squares, and 240 "white" squares (because the air purifier is covering the one in the middle). But since  $242 \neq 240$  it is impossible for this to work. So  $mn$  is simply 0. □

<sup>3</sup>This question has been voided due to the problem statement being confusing - the intended answer was 0, as there were no such ways, however this was confusing as the result involves dividing by 0

**§4.1.2 Coloured Markers [VOIDED]**

---

*Source: Original*

*Proposer: Charge#3766(481250375786037258)*

*Problem ID: 71*

*Date: 2020-01-08*

---

I have 3 red markers, 3 blue markers, and 3 green markers. I take the caps off and put on the caps randomly. Find the expected number of markers that have the same color cap and marker.

*Solution.*





**§4.1.3 Common Names [VOIDED]***Source: Original**Proposer: Charge#3766(481250375786037258)**Problem ID: 72**Date: 2020-01-09*

People with the 5 most common names stand in a line. These names are Liam, Noah, Olivia, Emma, and Ava (according to WolframAlpha in 2020). They are each given either \$0.10, \$1, \$10, \$100 or \$1000. If two people with the same number of letters in their name, same number of letter a's, or those with the same number of consonants in their name cannot receive the same amount of money, find the ten times the expected value of the total money they all receive.

*Solution.* [Write up by Charge#3766(481250375786037258)]

Note that the two most restrictive restrictions are the a's and the consonants. Note that each of Liam, Noah, Olivia, and Emma have the same number of a's and consonants. Looking at the least restrictive one, we can throw it out as Liam, Noah, and Emma already cannot have the same number of a's and consonants.

So what we have reduced the original problem to is the expected value of the total money that these people get under the restriction that Liam, Noah, Olivia, and Emma must receive different amounts of money.

Since Ava is not dependent on the others, we calculate her expected value separately. It is  $\frac{1111.1}{5} = \$222.22$ . We then find the total number of ways to distribute the money, excluding Ava. There are  $5 \cdot 4 \cdot 3 \cdot 2 = 120$  ways to do this.

We split these ways up into the total amount of money they receive. There are  $\binom{5}{4} = 5$  ways to pick a total amount of money, and there are a symmetrical amount of ways to distribute it. So for each way to pick a total amount, there is a  $\frac{1}{5}$  chance that it will happen.

So the expected value of this is  $\frac{4(1111.1)}{5} = \$888.88$ .

By linearity of expectation, our final expected value is simply \$1111.1. So our answer is 11111.

□

### §4.1.4 Funny Questions [VOIDED]

Source: HMMT (Year Unknown)

Proposer: Charge#3766(481250375786037258)

Problem ID: 73

Date: 2020-01-10

Brainy is a weird guy. He considers a performance on a QoTD Season *funny* if there's a pair of questions where 69 aspiring mathematicians get both problems correct first try, or get them wrong first try. Find the smallest number of people who attempted at least 1 problem such that Brainy will consider their performance *funny*, no matter how they answer. Note: A QoTD Season has 14 problems.

*Solution.* [Write up by Charge#3766(481250375786037258)]

Let one of the people answer  $k$  of the 14 problems correctly. Then, there are  $\binom{k}{2}$  pairs of problems they answered correctly, and  $\binom{14-k}{2}$  pairs of problems they answered incorrectly. This equates to  $k^2 - 14k + 91$  pairs of problems they answered that are either both correct or both incorrect.

By completing the square, we have  $(k - 7)^2 + 42$ . This means that no matter the  $k$ , the person will have answered at least 42 pairs of problems either both correctly or incorrectly.

Note that there are a total of  $2\binom{14}{2}$  "boxes" where there are 2 ways of making each pair correct or incorrect, and  $\binom{14}{2}$  ways of making a pair of problems.

Let there be a total of  $n$  people. Then we have  $42n$  "balls" to put in  $182$  "boxes". So we have  $42n \geq 182 \cdot 68 + 1 \implies n \geq 295$ . This gives a minimum of 295 QoTD Participants.

□

**§4.1.5 Colourful Integers [VOIDED]***Source: HMMT 2005**Proposer: Charge#3766(481250375786037258)**Problem ID: 74**Date: 2020-01-11*

Jason and XEM3 want to color the integers  $1, 2, \dots, 100$  in red, orange, yellow, green, and blue. They want to do so such that no two numbers  $x, y$  with  $x - y - 1$  divisible by 4 have the same color. All five colors do not have to be used. How many ways can this be done?

*Solution.* [Write up by Charge#3766(481250375786037258)]

This is equivalent to saying that we cannot have a number of residue 1 the same color as a number of residue 0, a number of residue 2 the same color as a number of residue 1, a number of residue 3 the same color as a number of residue 2, and a number of residue 0 the same color as a number of residue 3.

We have a few cases. We can have two of the five colors for the number of residue 0, and spread the rest out between the other residues. This gives  $\binom{5}{2} \cdot 2^{25} \cdot 3!$ . The same goes for the ones residue 1, 2, and 3. So we have  $4 \cdot \binom{5}{2} \cdot 2^{25} \cdot 3!$ .

This overcounts the times where all residues have one color. So we must subtract  $\binom{5}{4} \cdot 4!$ . This gives a final answer of  $8053063560$ .

□

## §5 Back to School (Season 5)

### §5.1 Week 1

#### §5.1.1 A Tricky Combination

---

*Source: Senior Mathematics Challenge, 2016 Q17*

*Proposer: sjbs#9839 (434767660182405131)*

*Problem ID: 75*

*Date: 2020-12-21*

---

A02 has to choose a three-digit code for his bike lock. The digits can be chosen from 1 to 9. To help him remember them, A02 chooses three different digits in strictly increasing order, for example 123. How many such codes can be chosen?

*Solution.*

If we take 3 numbers from  $\{1, 2, 3, \dots, 9\}$  there is exactly one possible valid combination. Thus, we have a bijection between valid codes and choosing 3 digits from 9. So the answer is  $\binom{9}{3} = \boxed{84}$ .

□

**§5.1.2 Geometric Sequence**

Source: Carnegie Mellon Informatics and Mathematics Competition, 2019 A/NT 1

Proposer: TadesPadhiary#8557 (665057968194060291)

Problem ID: 76

Date: 2020-12-22

Let  $a_1, a_2, \dots, a_n$  be in a geometric progression with  $a_1 = \sqrt{2}$  and  $a_2 = \sqrt[3]{3}$ . If

$$\frac{a_1 + a_{2013}}{a_7 + a_{2019}} = \frac{m}{n}$$

where  $\gcd(m, n) = 1$  and  $m, n$  are both positive integers, find  $m + n$ .

*Solution.*

If the common ratio is  $r$  and the first term is  $a$ , then we get the expression to be

$$\frac{a + ar^{2012}}{ar^6 + ar^{2018}} = \frac{1}{r^6} = \left(\frac{1}{r}\right)^6 = \left(\frac{\sqrt{2}}{\sqrt[3]{3}}\right)^6 = \frac{8}{9}$$

So our answer is  $8 + 9 = \boxed{17}$ .

□

**§5.1.3 Simultaneous Equation?**

---

Source: *British Mathematical Olympiad Round 1, 2009 P1*

Proposer: *sjbs#9839 (434767660182405131)*

Problem ID: 77

Date: 2020-12-23

---

Find the sum of all integers  $x$ ,  $y$ , and  $z$  such that

$$x^2 + y^2 + z^2 = 2(yz + 1) \text{ and } x + y + z = 4018$$

*Solution.*

We can write the first equation as  $x^2 + (y - z)^2 = 2$ , and so since  $x^2$  and  $(y - z)^2$  are both greater than or equal to 0, we must have the following cases:

1.  $x^2 = 0$  and  $(y - z)^2 = 2$
2.  $x^2 = 2$  and  $(y - z)^2 = 0$
3.  $x^2 = 1$  and  $(y - z)^2 = 1$

Clearly in cases one and two, we cannot have integer solutions, thus we consider the two possible subcases of the third: when  $x = \pm 1$ .

When  $x = \pm 1$ , we have either  $y = 1 + z$  or  $y = z - 1$ . Substituting these values into  $x + y + z = 4018$  gives us the set of  $x$ 's  $y$ 's, and  $z$ 's as  $\{-1, 1, 2008, 2009, 2010\}$

Hence our final answer is  $-1 + 1 + 2008 + 2009 + 2010 = \boxed{6027}$ . □

### §5.1.4 Tangent Circles

Source: *Original Problem/ Folklore*<sup>4</sup>

Proposer: *sjbs#9839 (434767660182405131)*


Problem ID: 78

Date: 2020-12-24

Two lines  $l_1$  and  $l_2$  intersect at an angle  $\alpha$  such that  $0 < \alpha < \frac{\pi}{2}$ . Given a circle  $\Gamma_n$  and radius  $r_n$ , with  $n \geq 0$ . Define a sequence of circles with  $r_0 > r_1 > \dots > r_n$  such that  $\Gamma_{n+1}$  is tangent to both lines  $l_1, l_2$ , as well as  $\Gamma_n$  and  $\Gamma_{n+1}$ . See the given diagram below for a construction.

Let the area inscribed between lines  $l_1, l_2$  and each of the circles  $\Gamma_0, \Gamma_1, \Gamma_2, \dots, \Gamma_n$  be  $A$ . As  $n \rightarrow \infty$  what is the value of  $\{1000A\}$  when  $r_0 = 3$  and  $r_1 = 1$ ? Where  $\{x\}$  is defined as the integer part of a number e.g.  $\{\pi\} = 3$


(A scientific calculator may be used to calculate  $\{x\}$ )




<sup>4</sup>I wrote this problem after reading Mathematics Aptitude Test 2016 Q4, however, this is obviously going to be a well known problem, and been done somewhere else

*Solution.*

We proceed by first finding the angle  $\alpha$ :



Thus we have  $\sin \frac{\alpha}{2} = \frac{2}{1+3}$ , hence  $\alpha = \frac{\pi}{3}$ . Now consider two general circles  $\Gamma_n$  and  $\Gamma_{n+1}$ :



By the sine rule again we have  $\sin \frac{\pi}{6} = \frac{r_n - r_{n+1}}{r_n + r_{n+1}}$ , thus  $\frac{r_n}{r_{n+1}} = \frac{1 - \frac{1}{2}}{1 + \frac{1}{2}}$ , and so the common ratio between the lengths of the radii is  $\frac{1}{3}$ . Though this is simply something can be deduced by the given sides of 3, 1,  $\dots$ . Hence, the areas of the circles inscribed, including  $\Gamma_0$ , will be given by the geometric series:

$$\begin{aligned} \pi(3)^2 + \pi \left[ \frac{1}{3}(3) \right]^2 + \pi \left[ \frac{1}{3^2}(3) \right]^2 + \dots &= 9\pi \left[ 1 + \frac{1}{3^2} + \frac{1}{3^4} + \dots \right] \\ &= \frac{9\pi}{1 - \frac{1}{9}} \end{aligned}$$



Therefore the total area inside the circles is  $\frac{81\pi}{8}$ . From the diagrams, we can clearly see therefore that the inscribed area, not fully accounting for  $\Gamma_0$  is going to be given by  $2 \cdot \frac{1}{2}3 \cdot \frac{3}{\tan \frac{\pi}{6}} = 9\sqrt{3}$ . Now all that's left to consider is  $\Gamma_0$  - the full area of it has been accounted for in the total area inside the circles calculation, however, unlike the other circles, it is not fully inscribed; the segment of area  $\frac{1}{2}(3)^2\left(\frac{\pi}{3} + \pi\right) = 6\pi$  has been over counted. Thus we have the total area inscribed by the circles as:

$$\begin{aligned} A &= 9\sqrt{3} - \frac{81\pi}{8} + 6\pi \\ &= 9\sqrt{3} - \frac{33\pi}{8} \end{aligned}$$

This leaves us with  $\{1000A\} = \boxed{2629}$

□

*Solution.* [Write up by AiYa#2278 (675537018868072458)]

By similarity,  $A\Gamma_0 = 3A\Gamma_1$ , so  $\Gamma_1\Gamma_0 = r_0 + r_1 = 4 = 2A\Gamma_1 \iff A\Gamma_1 = 2, A\Gamma_0 = 6$ . This implies that  $\alpha/2 = 30^\circ \iff \alpha = 60^\circ$ . Now rotate the figure two times around  $\Gamma_0$ , by  $120^\circ$  each time. The outer triangle is an equilateral triangle with side length  $6\sqrt{3}$ , and by symmetry, the area inside that triangle but outside the circles is  $3A$ . We also have  $A\Gamma_{n+1} = 2r_{n+1} = A\Gamma_n - r_n - r_{n+1} = r_n - r_{n+1} \iff \frac{r_n}{r_{n+1}} = \frac{1}{3}$ , so the area of the circles is

$$[\Gamma_0] + 3 \sum_{i=1}^{\infty} [\Gamma_i] = 9\pi + \pi \left( 1^2 + \frac{1}{3^2} + \frac{1}{3^4} + \cdots \right) = 9\pi + \frac{9\pi}{8} = \frac{99\pi}{8}$$

where  $[\cdot]$  represents the area of the circle. Now subtract from the area of the triangle and divide by 3:

$$A = \frac{1}{3} \left( \frac{3 \cdot 36\sqrt{3}}{4} - \frac{99\pi}{8} \right) = 9\sqrt{3} - \frac{33\pi}{8} \iff \{1000A\} = \boxed{2629}.$$

□

**§5.1.5 Santa's Elves***Source: AIME, 1985 P14**Proposer: Angry Any#4319 (580933385090891797)**Problem ID: 79**Date: 2020-12-25*

Santa has a number of elves. He wants to select the very best for Christmas. To do so, he makes all his elves compete in a ~~present-wrapping-competition~~ War Thunder tournament. Each elf competes against every other elf, and the winner receives one point, while the loser receives no points. If a particular match turns out to be a draw, each elf is awarded 0.5 points.

After the tournament has finished, Santa notices that exactly half of the points earned by each elf was done so when matched against the ten lowest scoring elves.

Given that each of the ten lowest scoring elves also earned half of their points against the other nine, how many elves does Santa have to choose from?

*Solution.*

Let there be  $n + 10$  elves. Then the bottom 10 elves score  $\binom{10}{2}$  points from each other while the rest score  $\binom{n}{2}$  points from those that are not in the bottom 10 people, and thus  $\binom{n}{2}$  points from the bottom 10 elves. Thus we get  $\frac{1}{2} \binom{n+10}{2} = \binom{n}{2} + \binom{10}{2}$  and thus solving we get  $n = 6, 15$ , but if  $n = 6$ , we get that it is expected for a player from the bottom 10 players will win when matched against a player from the top 6, which is impossible. Thus,  $n = 15$  and there are 25 elves Santa can choose from.  $\square$

## §5.1.6 2016 Algebra

Source: New Zealand Camp Selection Problems, 2016 P7

Proposer: brainysmurfs#2860 (281300961312374785)

Problem ID: 80

Date: 2020-12-26

Find the sum of all positive integers  $n$  for which the equation

$$(x^2 + y^2)^n = (xy)^{2016}$$

has positive integer solutions.

*Solution.*

**Solution 1: Edited Official Solution** Note that by AM-GM,  $x^2 + y^2 \geq 2xy \geq xy$  and so  $n \leq 2016$ . Let  $x = ad$  and  $y = bd$  where  $d = \gcd(x, y)$ . Then:

$$(a^2 + b^2)^n = (ab)^{2016} d^{4032-2n}.$$

Since  $a$  and  $b$  both divide the right-hand side but are relatively prime to the left-hand side, we get that  $a = b = 1$ . Thus, we have:

$$2^n = d^{4032-2n}.$$

Conversely,  $x = y = d$  for  $d$  satisfying this equation is a solution to the original equation. So  $d = 2^k$  for some integer  $k$ , meaning that  $n = k(4032 - 2n) \implies n = \frac{4032k}{2k+1}$ .

Since  $\gcd(k, 2k+1) = 1$ , we get that  $2k+1$  is a odd divisor of  $64 \times 63$ . Iterating these cases, we get that the possible values of  $n$  are 1344, 1728, 1792, 1920 and 1984, for a sum of 8768.

□

**§5.1.7**  $x^y$ 's

---

*Source: IMO, 1997 P5*

*Proposer: sjbs#9839 (434767660182405131)*

*Problem ID: 81*

*Date: 2020-12-27*

---

Given positive integers  $x$  and  $y$  and  $x^{y^2} = y^x$ , what is the value of the sum of all valid  $x$ 's and  $y$

*Solution.*

[1997 IMO P5 \(AoPS Thread\)](#)

Answer: 50

□

**§5.2 Week 2****§5.2.1 Pentagons**

---

*Source: HMMT November 2020 Guts Problem 1*

*Proposer: Angry Any#4319 (580933385090891797)*

*Problem ID: 82*

*Date: 2020-12-28*


---

Two pentagons are attached to form a new polygon  $P$ . What is the minimum number of sides  $P$  can have?

*Note: The two pentagons are not necessarily regular.*

*Solution.*

The answer is 3 as shown in the following diagram. Anything less will result in a shape that is not a polygon.



□

### §5.2.2 Digits in a String

Source: Folklore

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 83

Date: 2020-12-29

If we write the numbers 99999 down to 1 in the following string:

999999999899997...10987654321

What is the 42069<sup>th</sup> digit multiplied by the 42070<sup>th</sup> digit?

*Note: In the given string, we consider the first 9 on the left to be the first digit and 1 to be the last digit.*

*Solution.*

Observe that each number in the string can be split up into their respective numbers by separating them by 5 for all digits greater than 9999, 4 for all digits greater than 999, and less than 100000, and so on.

$$\underbrace{99999}_5 \underbrace{99998}_5 \underbrace{99997}_5 \dots \underbrace{10}_2 \dots$$

Since there are  $(99999 - 9999) \cdot 5 = 450000$  digits from five-digit numbers contained within the string, we know that the 42069<sup>th</sup> digit will be within a five-digit number. Since 42069 is one less than a multiple of 5, we can deduce that it will be the 4th digit in a five-digit number.

We note that the  $5n - 4$ th to  $5n$ th digit belong to the number  $100000 - n$  (by either Engineer's or mathematical induction).

Thus these two digits are the fourth and fifth digits of 91586 respectively, meaning their product is  $8 \cdot 6 =$

48.

□

**§5.2.3 Relatively Prime Function**

---

*Source: Original*

*Proposer: TaesPadhihary#8557 (665057968194060291)*

*Problem ID: 84*

*Date: 2020-12-30*

---

There are  $N$  tuples of integers  $a, b, c, d$  satisfying  $1 \leq a, b, c, d \leq 101$  and exactly 3 out of  $a, b, c, d$  are relatively prime to 2020.

What is the sum of the (not necessarily distinct) prime factors of  $N$ ?

*Solution.*

Note that there are 40 elements of  $\{1, 2, 3, \dots, 101\}$  which are relatively prime to 2020, and 61 elements which are not. We have 4 ways to choose which element out of  $a, b, c, d$  is not relatively prime to 2020, and so  $N = 4 \times 61 \times 40^3 = 2^2 \times 61 \times 2^6 \times 5^3$ .

Our answer is thus  $4 + 61 + 12 + 15 = \boxed{92}$ . □

**§5.2.4 Intersecting Circles**

---

Source: RMO Maharashtra and Goa, 2019 P6

Proposer: TaesPadhiary#8557 (665057968194060291)

Problem ID: 85

Date: 2020-12-31

---

Let  $k$  be a positive real number. In the Cartesian coordinate plane, let  $S$  be the set of all points of the form  $(x, x^2 + k)$  where  $x \in \mathbb{R}$ . Let  $C$  be the set of all circles whose center lies in  $S$ , and which are tangent to  $X$ -axis. Find the minimum value of  $k$  such that any two circles in  $C$  have at least one point of intersection.

If  $k = \frac{m}{n}$  where  $\gcd(m, n) = 1$  and  $m, n$  are positive integers, find  $m + n$ .

*Solution.*

**AoPS Solution**

Answer: 5

□



## §5.2.5 Discord Ping Fight

Source: Original

Proposer: Charge#3766(481250375786037258)

Problem ID: 86

Date: 2021-1-01


In a discord ping fight, 2 immature friends create a discord server with 5 channels numbered 1 through 5. Then player 1 and player 2 both select a (not necessarily distinct) odd numbered channel. Each round the two players ping in their channel.

After each round, they move to channel  $n + 1$  or  $n - 1$  where  $n$  is their current channel with equal probability, and start another round. If the expected number of rounds before both end up on the same channel and make up to each other can be expressed as  $\frac{a}{b}$  in lowest terms, find  $ab$ .

Note: If one of the friends is on channel 1 or 5, then the next turn they will be on channel 2 and 4 respectively.

Solution. [Write up by Charge#3766(481250375786037258)]

Note that the expected value of rounds it will take if the players start on channel 1 and 3 is the same as if they started on 3 and 5. Then we just have if the players are on 1 and 5 to be a separate case. Let our first state be when players are on 1 and 5, the second state be when there are players on 2 and 4, the third state be when one is on 1 and 3 or 3 and 5, and the last state be the end state. We now draw this diagram (forgive me if it looks terrible).



So we basically just solve for the expected value from each state and go from there.

Setting up a system of equations where  $S_n$  denotes the expected value to reach state 4 from  $n$ , we have

$$\begin{aligned}
 S_1 &= 1 + S_2 \\
 S_2 &= 1 + \frac{1}{4}S_1 + \frac{1}{2}S_3 + \frac{1}{4}S_4 \\
 S_3 &= 1 + \frac{1}{2}S_2 + \frac{1}{2}S_4 \\
 S_4 &= 0
 \end{aligned}$$

So we have  $S_1 = \frac{9}{2}$ ,  $S_2 = \frac{7}{2}$ ,  $S_3 = \frac{11}{4}$ . Now we do some simple casework.

Case 1: It starts on state 1. This happens with probability  $\frac{2}{9}$  so the expected number of rounds minus the one at the end from state 1 is 1.

Case 2: It starts on state 3. This happens with probability  $\frac{4}{9}$  so the expected number of rounds minus the one at the end from state 3 is  $\frac{11}{9}$ .

So by linearity of expectation, the expected value of rounds before the end is  $\frac{20}{9}$ . So  $ab = \boxed{180}$

□

### §5.2.6 Falling Cards

Source: New Zealand Monthly Maths Workshop, December 2020, Problem 6

Proposer: brainysmurfs#2860 (281300961312374785)

Problem ID: 87

Date: 2021-1-02

Point One Nine throws a standard pack of cards into the air such that each card is equally likely to land face up or face down and lands independently of other cards. The total value of the cards which landed face up is then calculated.

Suppose the probability that the total is divisible by 13 is  $\frac{m}{n}$  with  $m, n \in \mathbb{Z}^+$ ,  $\gcd(m, n) = 1$ . Calculate the largest integer value of  $x$  such that  $2^x \leq n$ .

Note: Values of cards are assigned as follows: Ace = 1, 2 = 2, ..., Jack = 11, Queen = 12, King = 13.

*Solution.*

Note that we may discard the Kings, since they represent values which are 0 (mod 13). Further, note that 2 is a generator in mod 13. In this way we establish a bijection between the values of the cards 1, 2, 3, ..., 12, 1, 2, 3, ..., 12, ..., 12 (4 sets of 1 to 12) and  $2^0, 2^1, \dots, 2^{47}$  in mod 13.

Each number has a unique binary representation and since each configuration of cards is equally likely, each number from 0 to  $2^{48} - 1$  is equally likely. By Fermat's Little Theorem, since  $2^{12} \equiv 1 \pmod{13}$ , we get  $13 \mid 2^{48} - 1$ .

Thus out of those binary numbers  $\frac{2^{48}-1}{13} + 1$  are divisible by 13, and so our probability is

$$\frac{2^{48} + 12}{13 \cdot 2^{48}} = \frac{2^{46} + 3}{13 \cdot 2^{46}}.$$

But since  $2^{46} \equiv 2^{-2} \equiv -3 \pmod{13}$ , we get that the numerator is divisible by 13, and so  $n = 2^{46}$ . From here we get that our answer is 46.

□

**§5.2.7 Computing the Area of a Triangle**

---

Source: HMMT Feb 2019 Geometry P8

Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 88

Date: 2021-1-03

---

In triangle  $ABC$  with  $AB < AC$ , let  $H$  be the orthocenter and  $O$  be the circumcenter. Given that the midpoint of  $OH$  lies on  $BC$ ,  $BC = 1$ , and the perimeter of  $ABC$  is 6, compute the area of  $ABC$ .

If this area is  $\frac{m}{n}$  for coprime positive integers  $m$  and  $n$ , find  $m + n$ .

*Solution.*

**Official Solution**

Answer: 13

□

## §6 Third Week of School (Season 6)

### §6.1 Week 1

#### §6.1.1 Maximising Square Factors

Source: Senior Mathematical Challenge, 2015 Q18

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 89

Date: 2021-01-04

What is the largest integer  $k$  for which  $k^2$  is a factor of  $11!$ ?

*Solution.*

Let us write the factorisation of  $10!$  in a table, so we can see the indexes of the powers.

	7	5	3	2
2	0	0	0	1
3	0	0	1	0
4	0	0	0	2
5	0	1	0	0
6	0	0	1	1
7	1	0	0	0
8	0	0	0	3
9	0	0	2	0
10	0	1	0	1

We want to find  $k^2$ , where  $k$  is maximised. That means we want to add as many of the indexes together such that they all end up even. From inspecting the table, we see that they all add up to an even number so long as we don't include the 7. Thus  $k^2 = \frac{10!}{7} = 5^2 \cdot 3^4 \cdot 2^8$ , this gives an answer of  $k = \boxed{720}$   $\square$

**§6.1.2 Njoy's Balls***Source: Original Problem**Proposer: TaesPadhiary#8557 (665057968194060291)**Problem ID: 91**Date: 2021-01-05*

NJOY has a box of 200 balls, 100 of which are blue, 98 of which are red, and 2 of which are green.

At each stage he randomly selects one ball from the box. If it is blue, he wins; if it is red, he loses, and if it is green, he replaces the ball and draws another one with the same rules.

If the probability that he wins is  $\frac{m}{n}$  where  $\gcd(m, n) = 1$  and  $m, n$  are both positive integers, find  $100m + n$ .

*Solution.*

Let  $P$  be the probability of a win. Note that  $P = P(\text{blue}) + P(\text{green}) * P$ , since after he draws and replaces a green ball the state returns to the starting state. Solving this with the provided values, we get

$$P = \frac{1}{2} + \frac{1}{100}P$$

and hence  $\frac{99}{100}P = \frac{1}{2}$  and so  $P = \frac{50}{99}$ , giving us  $100m + n = 5000 + 99 = \boxed{5099}$ . □

*Solution.* [Solution by AiYa#2278 (675537018868072458)]

The last ball NJOY picks will be blue or red, and the probability that he picks that final ball does not depend on his history of picking greens. Therefore, the green balls are irrelevant to the problem and the probability he picks a blue is  $\frac{100}{100+98}$  giving us our answer of  $\boxed{5099}$ . □

### §6.1.3 Aiya's Function

Source: Original Problem

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 90

Date: 2021-01-06

Let  $f : [0, 1] \rightarrow \mathbb{R}$  be a function with the following properties:

$$f(x) + f\left(\frac{1}{2}\right) f(1-x) = 2f\left(\frac{1}{2}\right)$$

- $2f(x) = f(3x)$
- $f$  is non-decreasing

Then the sum of the possible values of  $f\left(\frac{4}{5}\right) + 1$  can be written in the form  $\frac{m}{n}$ , where  $m$  and  $n$  are relatively prime positive integers. What is the value of  $100m + n$ ?

*Solution.* [Solution by AiYa#2278 (675537018868072458)]

First, find some basic values:  $2f(0) = f(0) \iff f(0) = 0$  and  $f(1/2) + f(1/2)^2 = 2f(1/2) \iff f(1/2) = 0, 1$ . Now if  $f(1/2) = 0$  note that  $f(1) = 0$ ; since  $f$  is nondecreasing  $f$  is the zero function. Now for  $f(1/2) = 1$ :  $f(x) + f(1-x) = 2 \iff f(1) = 2$ . It's fruitless to get  $4/5$  from 0, 1, and  $1/2$  by repeated operations of  $1-x$  and  $x/3$ ; however note that  $f(1/3) = 1 \iff f(2/3) = 1$  and since  $f$  is nondecreasing all values of  $x$  between  $1/3$  and  $2/3$  inclusive will have  $f(x) = 1$ . Then finding  $f(3/5)$  is easy: it's just 1, so  $f(1/5) = 1/2 \iff f(4/5) = 3/2$ . Our possible values of  $g(4/5)$  are 1 and  $5/2$ . This gives us an answer of

702

□

## §6.1.4 The AIME Cyclic

Source: 2018 AMC 12A, Q20 of 25

Proposer: Angry Any#4319 (580933385090891797)


Problem ID: 92

Date: 2021-01-07

Triangle  $ABC$  is an isosceles right triangle with  $AB = AC = 3$ . Let  $M$  be the midpoint of hypotenuse  $\overline{BC}$ . Points  $I$  and  $E$  lie on sides  $\overline{AC}$  and  $\overline{AB}$ , respectively, so that  $AI > AE$  and  $AIME$  is a cyclic quadrilateral. Given that triangle  $EMI$  has area 2, the length  $CI$  can be written as  $\frac{a-\sqrt{b}}{c}$ , where  $a$ ,  $b$ , and  $c$  are positive integers and  $b$  is not divisible by the square of any prime. What is the value of  $10000a + 100b + c$ ?

*Solution.*

Let  $IM = a$ , since  $IM = EM$ , we have  $a^2 = 4$ . Further, as  $CM = \frac{1}{2}\sqrt{3^2 + 3^2} = \frac{3\sqrt{2}}{2}$ , and given  $\triangle ABC$  being isosceles we have  $\angle BCA = \frac{\pi}{4}$ . Therefore  $a^2 = x^2 + \frac{9}{2} - 3x$ , and so  $x^2 - 3x + \frac{9}{2} = 4 \iff x = \frac{3 \pm \sqrt{7}}{2}$  (We take the negative as  $AI > AE$ ). Hence, by the problem statement we have our answer as 30702



□

## §6.1.5 Prime Floors

Source: NIMO AoPS Thread

Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 93

Date: 2021-01-08

Let  $p = 10^9 + 7$  be a prime. Find the remainder when

$$\left\lfloor \frac{1^p}{p} \right\rfloor + \left\lfloor \frac{2^p}{p} \right\rfloor + \left\lfloor \frac{3^p}{p} \right\rfloor + \dots + \left\lfloor \frac{(p-3)^p}{p} \right\rfloor + \left\lfloor \frac{(p-2)^p}{p} \right\rfloor$$

is divided by  $p$

*Solution.*

We will prove the general case where  $p$  is an odd prime.

Let  $p$  be an odd prime. we have  $n^{p-1} \equiv 1 \pmod{p}$  and thus  $n^p \equiv n \pmod{p}$  for all positive integers  $n < p$ . Thus  $\left\lfloor \frac{n^p}{p} \right\rfloor = \frac{n^p - n}{p}$  and the required sum is equivalent to

$$\frac{2^p + 3^p + \dots + (p-2)^p}{p} - \frac{2 + 3 + 4 + \dots + (p-2)}{p}$$

Now we claim that  $\frac{n^p + (p-n)^p}{p} \equiv 0 \pmod{p}$  for all integer  $n$ , which is indeed true since by the binomial theorem, all terms of  $(p-n)^p$  are  $0 \pmod{p^2}$  except  $-n^p$ . Therefore  $n^p + (p-n)^p \equiv 0 \pmod{p^2}$  and  $\frac{n^p + (p-n)^p}{p} \equiv 0 \pmod{p}$ . This implies that the first sum is actually just  $0 \pmod{p}$  and the desired answer stems solely from the second sum.

The second sum can be grouped into  $\frac{p-3}{2}$  pairs that add up to  $p$ , and thus the answer is  $-\frac{p-3}{2} = \frac{p+3}{2} = \boxed{500000005} \pmod{10^9 + 7}$ .  $\square$



### §6.1.6 Nested Periodic Functions

Source: USA EGMO TST #2, 2020 P6

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 94

Date: 2021-01-09

For a polynomial  $P(x)$  with integer coefficients such that for each positive integer  $m$ ,

$$P^m(0) \equiv 0 \pmod{2020} \iff m \equiv 0 \pmod{N}$$

Where  $N \in \{1, 2, \dots, 2019\}$ . What is the largest such value of  $N$ ?

(Here we denote  $P^m$  to mean the function  $P$  applied  $m$  times, so  $P^1(0) = P(0)$ ,  $P^2(0) = P(P(0))$ , and so forth.)

Solution.

Art of Problem Solving write ups

Answer:

□

### §6.1.7 Braniy's Party

Source: China Team Selection Test, 2015 Day 1 P3

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 95

Date: 2021-01-10

Brainy, being the socialite he is, has invited a large number of friends to a little soirée. Brainy will be feeding his friends Chinese Noodle Soup, made by Tony himself.

Wanting to impress Asuka, and Zero-Two, both of whom will be at the party, he asks Tony to provide the party with many different flavours of soup - enough so that for each flavour, there are 260 bowls of soup.

When Brainy's friends arrive at the party, each of them take exactly two soups, each of different flavours. Given that for any 179 of Brainy's friends, there will at least be two who have at least one flavour of Chinese Noodle Soup in common, and assuming Brainy has invited as many people as possible to the party, how many people are in attendance?

(A four-function calculator may be used)

Solution.

Art of Problem Solving write ups

Answer:

□

## §6.2 Week 2

### §6.2.1 Negative Sum

---

*Source: Original Problem*

*Proposer: Angry Any#4319 (580933385090891797)*

*Problem ID: 96*

*Date: 2021-01-11*

---

There is an ordered tuple of  $n$  positive real numbers  $\{a_1, a_2, a_3, \dots, a_n\}$  such that when any element is replaced by itself multiplied by  $-2021$ , the sum of the elements of the ordered tuple becomes negative. What is the maximum value of  $n$ ?

For example,  $\{1, 3, 5, 7\}$  satisfies this property, but  $\{1, 4000\}$  does not.

*Solution.*

Note that  $n = 2021$  is trivially possible through the construction  $\{1, 1, 1, 1, \dots, 1\}$ .

Now we prove that a higher number is not possible. WLOG, let  $a_1 \geq a_2 \geq \dots \geq a_n$ . Then,  $a_1 + a_2 + a_3 + \dots + a_{n-1} - 2021a_n \geq (n-1)a_n - 2021a_n = (n-2022)a_n$ . Thus, the condition cannot hold true if  $n-2022 \geq 0$  or  $n \geq 2022$ , which concludes the proof.

□

**§6.2.2 Doubling Digit Sum**

---

*Source: Australian Intermediate Mathematical Olympiad, 1999 Q10*

*Proposer: brainysmurfs#2860 (281300961312374785)*

*Problem ID: 97*

*Date: 2021-01-12*

---

$N$  is the smallest positive integer such that the sum of the digits of  $N$  is 18 and the sum of the digits of  $2N$  is 27. Find  $N$ .

*Solution.*

We iterate through cases of  $2N$ .

The only 3 digit number with a digit sum of 27 is 999 but  $2N = 999$  does not work since 999 is not even.

$2N = 1998$  is the next smallest even number with digit sum 27 and  $N = 999$  does not work.

$2N = 2898$  is the next smallest even number with digit sum 27 and it does work, giving  $N = \boxed{1449}$ . □

### §6.2.3 Tony Vs. Wang

Source: *British Mathematical Olympiad, Round 1, 2018 P1*

Proposer: *sjbs#9839 (434767660182405131)*

Problem ID: 98

Date: 2021-01-13

Tony divides 365 by each of  $1, 2, 3, \dots, 365$  in turn, writing down a list of the 365 remainders. Then Wang divides 366 by each of  $1, 2, 3, \dots, 366$  in turn writing down a list of the 366 remainders. What is the difference between the two summed lists?

*Solution.*

Remainder for remainder, the numbers in Wang's list will always be one greater than those in Tony's list - with exception to the cases where the remainder is 0, in which case, for a number  $n$  dividing 366, Tony will have a remainder of  $n - 1$ . Since  $366 = 2 \cdot 3 \cdot 61$ , there are 8 remainders in Wang's list that are 0. Now we wish to find the  $n - 1$  remainders. Well, we have  $n \in \{1, 2, 3, 6, 61, 122, 183\}$  so we must have the remainders for Tony, when Wang's remainder is 0, being  $\{0, 1, 2, 5, 60, 121, 182\}$ . To summarise these results, we have:

$$T + (0 + 1 + 2 + 5 + 60 + 121 + 182) = W + 365 - 8 + 1$$

$$\therefore \boxed{13} = W - T$$

□

### §6.2.4 Equilateral Decagon

Source: OTIS excerpts P118

Proposer: Angry Any#4319 (580933385090891797)

Problem ID:

Date:

Let  $ABCDEZYXWV$  be an equilateral decagon with interior angles  $\angle A = \angle V = \angle E = \angle Z = \angle C = 90^\circ$ ,  $\angle W = \angle Y = 135^\circ$ ,  $\angle B = \angle D = 225^\circ$ , and  $\angle X = 270^\circ$ . Find the sum of all  $1 < n < 100$  such that  $ABCDEZYXWV$  can be partitioned into  $n$  congruent polygons.

*Solution.*

Cut along horizontally such that the vertical distance for each line is constant. This shows that all  $n$  is possible. Thus, the answer is  $\frac{99 \cdot 100}{2} - 1 = 4949$ . □

**§6.2.5 Triangles on a Cubic**

---

*Source: 2017 AMC 12B, Q23*

*Proposer: Angry Any#4319 (580933385090891797)*

*Problem ID:*

*Date:*

---

The graph of  $y = f(x)$ , where  $f(x)$  is a polynomial of degree 3, contains points  $A(2, 4)$ ,  $B(3, 9)$ , and  $C(4, 16)$ . Lines  $AB$ ,  $AC$ , and  $BC$  intersect the graph again at points  $D$ ,  $E$ , and  $F$ , respectively, and the sum of the  $x$ -coordinates of  $D$ ,  $E$ , and  $F$  is 24. What is  $f(0)$ ?

*Solution.*

Art of Problem Solving write-ups



**§6.2.6 Jumping Frogs**

---

*Source: International Mathematics Competition for University Students, 2018 P8*

*Proposer: sjbs#9839 (434767660182405131)*

*Problem ID:*

*Date:*

---

Let  $\Omega = \{(x, y, z) \in \mathbb{Z}^3 : y + 1 \geq x \geq y \geq z \geq 0\}$ . A frog moves along the points of  $\Omega$  by jumps of length 1. Determine the number of paths the frog can take to reach  $(12, 12, 12)$  starting from  $(0, 0, 0)$  in exactly 36 jumps.

*Solution.*

(Art of Problem Solving write ups)

Answer:

□

### §6.2.7 Minimising Ratios

Source: China Team Selection Test, 2008 Quiz 1 P1

Proposer: sjbs#9839 (434767660182405131)

Problem ID:

Date:

Suppose a point  $P$  is a random point chosen inside a triangle  $XYZ$ . Extend the segment  $XP$  until it intersects with the circumcircle of  $PYZ$ , call the point of intersection which isn't  $P$ ,  $X_1$ . Similarly, for points  $Y$  and  $Z$  construct points  $Y_1$  and  $Z_1$ . What is the smallest value of

$$\left(1 + 2\frac{XP}{PX_1}\right) \left(1 + 2\frac{YP}{PY_1}\right) \left(1 + 2\frac{ZP}{PZ_1}\right)$$

Solution.

Art of Problem Solving write ups

Answer: 8

□



## §7 PSC's Adventure (Season 7)

### §7.1 Week 1

#### §7.1.1 The Jungle Polygon

---

*Source: Folklore*

*Proposer: sjbs#9839 (434767660182405131)*

*Problem ID: 103*

*Date: 2021-01-18*

---

The Problem Selection Committee ventures into a jungle where they find a regular polygon drawn on the ground. Brainysmurfs takes out his infinite precision protractor and measures one of the interior angles to be 179 degrees. How many sides does the polygon have?

*Solution.*

The formula for the sum of the interior angles is given by  $180(n - 2)$ , and thus we have  $180(n - 2) = 179n \iff n = \boxed{360}$   $\square$

### §7.1.2 Uphill and downhill

Source: Original Problem

Proposer: TaesPadhihary#8557 (665057968194060291)

Problem ID: 104

Date: 2021-01-19

After Brainy discovered a map hidden within the regular polygon, he is able to navigate the PSC through the Jungle. It takes Brainy and the team 4 hours to navigate out of the jungle, moving 2 kilometres per hour walking up inclined surfaces, 6 kilometres an hour when walking down, and 3 kilometres an hour on flat surfaces.

Upon reaching the edge of the jungle Brainy realises that they accidentally left .19 back where they started. Given that it took Brainy and the PSC 6 hours to return to where they started, to rescue .19, what is the total distance Brainy travelled (in kilometres)?

*Solution.* [Solution by TaesPadhihary#8557 (665057968194060291)]

Let the two “other” points be  $C$  and  $D$ , going from left to right (so that the labels when reading from left to right are  $A, C, D, B$ ). We let  $AC = x, CD = y, DB = z$ . We have  $\frac{x}{2} + \frac{y}{3} + \frac{z}{6} = 4$  and  $\frac{x}{6} + \frac{y}{3} + \frac{z}{2} = 6$ . Multiplying both equations by 6 gives  $3x + 2y + z = 24$  and  $x + 2y + 3z = 36$ . Adding them gives  $4x + 4y + 4z = 60$ , so our desired value, is  $\boxed{30}$ .  $\square$

*Solution.* [Solution by brainysmurfs#2860 (281300961312374785)]

Note that the harmonic mean of 2 and 6 kph is 3kph - in particular, since in total there's the same amount of uphill and downhill, we get that the average speed is 3 kph and thus the PSC travelled  $\boxed{30}$  km.  $\square$

**§7.1.3 Just a Beauty!**

Source: *CMC Mock ARML 2020 i7*

Proposer: *TaesPadhiary#8557 (665057968194060291)*

Problem ID: 105

Date: 2021-01-20

The PSC leave the jungle and find themselves on a long winding road, where they encounter a toll gate. The toll keeper, Joe, asks for an amount of smackaroo's equal to

$$31 \left( 1 + \frac{30}{2} \left( 1 + \frac{29}{3} \left( 1 + \frac{28}{4} \left( \cdots \left( 1 + \frac{17}{15} \right) \cdots \right) \right) \right) \right).$$

How many smackaroo's should the PSC pay Joe?

*(A four-function calculator may be used)*

*Solution.*

**Official Solutions Booklet p.9**

Answer: 9223372036854775806



### §7.1.4 Probability of Intervals

Source: *One Hundred Problems - 4th Edition Q93*

Proposer: *HoboSas#3200 (310725130097786880)*

Problem ID: 106

Date: 2021-01-21

After paying Joe what is due, the PSC team find themselves at the gates of a destitute underground research facility. Upon exploring the ransacked and decaying tomb they find themselves in a laboratory experimenting with teleportation. The PSC try to get it working:

.19, Brainy, and Yuchan enter the teleporter where they are each positioned randomly along a 1m strip.

For the teleporter to work, .19 and Brainy must be less than half a meter apart, and .19 and Yuchan must also be less than half a meter apart.


If the probability that the teleporter works is  $\frac{m}{n}$ , find  $100m + n$

*Solution.* [Solution by HoboSas#3200 (310725130097786880)]

The problem statement can be reduced to this:

Let  $a, b$  and  $c$  be real numbers randomly chosen from the interval  $[0, 1]$ , if the probability of  $|a - b| < \frac{1}{2}$  and  $|b - c| < \frac{1}{2}$  can be expressed as  $\frac{m}{n}$ , find  $100m + n$ .

I will pursue a geometrical approach using a 3D space with coordinates  $(a, b, c)$ , such that for every triplet  $(a, b, c)$  there exist one and only point enclosed in the cube whose vertices are  $(0, 0, 0)$   $(1, 0, 0)$   $(1, 1, 0)$   $(0, 1, 1)$  and so on. The probability we are seeking can be seen as ratio between volumes, in particular the volume of the intersection of the two solids  $|a - b| < \frac{1}{2}$  and  $|b - c| < \frac{1}{2}$ , over the volume of the cube with side 1. Let's focus on finding what the solid  $|a - b| < \frac{1}{2}$  looks like: if we consider the plane  $c=0$ , with some trivial analytic algebra we can draw the following polygon: Introducing the 3rd dimension,  $|a - b| < \frac{1}{2}$  turns out to be an



hexagonal prism (the base is shown above) whose height is 1 (along the  $c$ -axis). Similarly  $|b - c| < \frac{1}{2}$  is the same solid, but rotated 90 degrees around the line parallel to the  $b$ -axis and going through the point  $(\frac{1}{2}, \frac{1}{2}, 0)$ . Let's now compute the intersection between those solids, which turns out to be the sum of few smaller solids, in particular: 2 cubes with side  $\frac{1}{2}$ , 2 square pyramids with base length and height  $\frac{1}{2}$  and 4 triangular prism, having height  $\frac{1}{2}$  and an isosceles right triangle with side  $\frac{1}{2}$  as base. Finally the probability is

$$p = \frac{2 \cdot \frac{1}{8} + 2 \cdot \frac{1}{24} + 4 \cdot \frac{1}{8}}{1} = \frac{7}{12}$$

which leads to 712 as final answer. □

### §7.1.5 Dividing Functions

Source: Original Problem

Proposer: ChristopherPi#8528 (696497464621924394)

Problem ID: 106

Date: 2021-01-22

Find the smallest integer  $n > 1$  for which there exist positive integers  $a_1, a_2, \dots, a_n$  such that

$$(a_1)^2 + \dots + (a_n)^2 \mid [(a_1 + \dots + a_n)^2 - 1]$$

*Solution.* [Solution by ChristopherPi#8528 (696497464621924394)]

Note that squares are congruent to themselves (mod 2), so

$$(a_1)^2 + \dots + (a_n)^2 = a_1 + \dots + a_n = (a_1 + \dots + a_n)^2 \pmod{2}$$

Since  $(a_1)^2 + \dots + (a_n)^2$  divides  $(a_1 + \dots + a_n)^2 - 1$  which has a different parity, we must have the former odd and the latter even, so  $(a_1 + \dots + a_n)$  is odd. As all odd squares are 1 mod 8,  $(a_1 + \dots + a_n)^2 - 1$  must be divisible by 8, and since  $(a_1)^2 + \dots + (a_n)^2$  is odd, we must have

$$\frac{(a_1 + \dots + a_n)^2 - 1}{(a_1)^2 + \dots + (a_n)^2} \geq 8$$

Using Cauchy-Schwartz on the sequences  $(a_1, \dots, a_n)$  and  $(1, 1, \dots, 1)$  gives

$$n((a_1)^2 + \dots + (a_n)^2) \geq (a_1 + \dots + a_n)^2 > ((a_1 + \dots + a_n)^2 - 1)$$

, which is a direct contradiction if  $n$  is less than 9. Therefore we must have  $n \geq 9$ . It is easy to see  $n = 9$  works, with an example construction being  $(1, 1, 1, 1, 1, 1, 1, 2, 2)$  which works as  $7 * (1^2) + 2 * (2^2) = 15$  and  $(7 * 1 + 2 * 2)^2 - 1 = 120$ , with  $120/15 = 8$  being an integer.

9 □

**§7.1.6 Paper Monster**

---

Source: *Online Math Open* 2013 P29

Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 107

Date: 2021-01-23

---

After the TST the PSC meets Chrispi.

Chrispi has 255 sheets of paper, each labeled with a unique nonempty subset of 1, 2, 3, 4, 5, 6, 7, 8. Each minute, he chooses one sheet of paper uniformly at random out of the sheets of paper not yet eaten.

Then, he eats that sheet of paper, and all remaining sheets of paper that are labeled with a subset of that sheet of paper (for example, if he chooses the sheet of paper labeled with 1, 2, he eats that sheet of paper as well as the sheets of paper with 1 and 2).

The expected value of the number of minutes that Chrispi eats a sheet of paper before all sheets of paper are gone can be expressed in the form  $\frac{m}{n}$ , where  $m$  and  $n$  are relatively prime positive integers. Find  $100m + n$ . difficulty

*Solution.*

Art of Problem Solving write-ups

Answer:

□

### §7.1.7 Tan's Optimisation

Source: Cambridge Handout

Proposer: tanoshii#3160 (300065144333926400)

Problem ID: 108

Date: 2021-01-24

After Brainy and Yuchan placed joint first at the IMO, both perfect scoring, they decide to take the day off before going on a journey to replace .19 and add more people to the problem solving committee. As part of the recruitment process, they ask the following question to the applicants:

Given that the maximum value of

$$\frac{x_1x_2 + x_2x_3 + \cdots + x_{20}x_{21} + x_{21}x_1}{x_1^2 + \cdots + x_{21}^2}$$

is  $M$  for  $x_1 + \cdots + x_{21} = 0$ , find  $\lfloor 1000M \rfloor$

What answer should the applicants put down?

(A scientific calculator may be used)

*Solution.* [Solution by tanoshii#3160 (300065144333926400)]

We will prove it for a given  $n$ , here  $n = 21$ .

Let  $\omega$  be the principle  $n$ th root of unity.

Let

$$y_k = \frac{1}{\sqrt{n}} \sum_{i=1}^n x_i \omega^k.$$

Note that

$$|y_1|^2 + |y_2|^2 + \cdots + |y_n|^2 = |x_1|^2 + |x_2|^2 + \cdots + |x_n|^2$$

and

$$x_1x_2 + \cdots + x_nx_1 = \frac{1}{n} \sum_{k=1}^n (\omega^k + \omega^{-k}) y_k^2$$

Since  $x_1 + \cdots + x_n = 0$ , we get  $y_n = 0$ , so the sum is maximised when  $y_{n-1}$  or  $y_1$  is maximal since  $k = 1, n-1$  are the values where  $\omega^k + \omega^{-k}$  is the biggest and that gives  $2 \cos(\frac{2\pi}{n})$ .  $\square$



## §7.2 Week 2

### §7.2.1 A Digital Product equal to Half the Number

Source: Folklore

Proposer: *sjbs#9839 (434767660182405131)*

Problem ID: 110

Date: 2021-01-25

AiYa claims that he has found the smallest two-digit positive integer with the property that when divided by two, it is equal to the product of its digits.

Assuming AiYa is correct, what number has he found?

*(If needed, a four-function calculator may be used)*

*Solution.*

Let the integer have digits  $x$  and  $y$ . We require:

$$\begin{aligned} xy &= \frac{10x + y}{2} \\ 2xy - 10x - y &= 0 \\ 2xy - 10x - y + 5 &= 5 \\ (2x - 1)(y - 5) &= 5 \end{aligned}$$

Since 5 is prime, it has factors of either 1 or 5. Hence we have either  $y - 5 = 5$  or  $y - 5 = 1$ . We don't need to consider negatives as it's clear  $x$  would not be positive (and non-zero). We see that  $y = 6$ , as in the other case,  $y > 9$ . This means that  $2x - 1 = 5$ , giving  $x = 3$ . So the answer is 36 □

**§7.2.2 Unique Odd Numbers**

---

Source: Folklore

Proposer: Kiesh#0917 (544960202101751838)

Problem ID: 112

Date: 2021-01-26

---

Call a number unique if each of its digits are unique (no two are the same). How many odd integers in the interval  $[3 \cdot 10^4, 8 \cdot 10^4]$  are unique?

(A four-function calculator may be used)

*Solution.*

There are 5 ways to choose the first digit. If the first digit is odd, then we must consider the final digit too. In the even cases, there are 2 numbers to choose from for the first digit, then last digit we have 5 odd digits to choose from. Then for the other 3 digits, we have  $7!$  ways to pick digits, so there are  $2 \cdot 5 \cdot \frac{8!}{5!}$  ways when the first digit is even. When it is odd, however, there is one less odd digit to choose from for the last digit. There are 3 ways to choose the first digit and 4 ways to choose the last digit. As before there are  $\frac{8!}{5!}$  ways to choose the middle three digits. This gives us a total of  $3 \cdot 4 \cdot \frac{8!}{5!}$  when the first digit is odd.

Thus in total, there are  $2 \cdot 5 \cdot \frac{8!}{5!} + 3 \cdot 4 \cdot \frac{8!}{5!} = \boxed{7392}$

□

### §7.2.3 Maximising $a^4 + b^4$

Source: Folklore

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 113

Date: 2021-01-27

Returning home from Matteddy's demonstration, the PSC are trapped in a vault by NJOY, trying to join the PSC by coercion and brute force. On the wall are two haikus. The first reads:

The sum of the squares  
of  $a$  and  $b$  minus their  
product fears 15

They take this to mean that the given expression of  $a$  and  $b$  is at most 15, before reading the second haiku:

The unfearing sum  
of their fourth powers is on  
the door of freedom

which they take to mean that the door labeled with the maximal value of  $a^4 + b^4$  is their exit.

After some thought, they scan the walls for they want, climb up to it and walk out, to the rising moon. They've escaped.

What door did they leave through?

*Solution.*

Notice that  $a^4 + b^4 = (a^2 + b^2)^2 - 2(ab)^2$ . Thus,  $a^4 + b^4 \leq (2 + ab)^2 - 2(ab)^2$  - this is simply a quadratic in  $ab$ .

Therefore, we have, by *the method of maximising a quadratic of your choosing*

$$a^4 + b^4 \leq 450 - (ab - 2)^2$$

So our answer is 450.

□

### §7.2.4 Minimising Perimeter

Source: China Mathematical Competition (Extra Test), 2003 P2

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 114

Date: 2021-01-28

Find the **minimum** perimeter of a triangle having integer sides  $a > b > c > 0$ , such that

$$\frac{3^a}{10^4} - \left\lfloor \frac{3^a}{10^4} \right\rfloor = \frac{3^b}{10^4} - \left\lfloor \frac{3^b}{10^4} \right\rfloor = \frac{3^c}{10^4} - \left\lfloor \frac{3^c}{10^4} \right\rfloor,$$

where  $\lfloor x \rfloor$  denotes the greatest integer less than or equal to  $x$ , e.g.  $\lfloor \pi \rfloor = 3$ , and so on.

*Solution.*

Note that this condition implies  $l \equiv m \equiv n \pmod{\text{ord}_{10000}(3)}$ . Let us calculate  $\text{ord}_{10000}(3)$ .

By LTE,  $\nu_2(3^n - 1) = \nu_2(3 - 1) + \nu_2(n) + \nu_2(3 + 1) - 1 = \nu_2(n) + 2$ , and so if  $3^n \equiv 1 \pmod{2^4}$  we must have  $\nu_2(n) \geq 2$ , implying that  $n \geq 4$  and so  $\text{ord}_{2^4}(3) = 4$ .

By LTE again, note that  $\nu_5(81^n - 1) = \nu_5(81 - 1) + \nu_5(n) \geq 4$ . Hence we must have  $\nu_5(n) \geq 3$  which implies  $n \geq 125$ . Since  $81 = 3^4$  we get that  $\text{ord}_{5^4}(3) = 500$ .

Now  $\text{ord}_{10^4}(3) = \text{lcm}(\text{ord}_{2^4}(3), \text{ord}_{5^4}(3)) = 500$ .

Hence for our minimum we must have  $l = n + 1000, m = n + 500$ . Since they are the sides of a triangle  $l < m + n \implies n + 1000 < 2n + 500 \implies n > 500$ . So  $n = 501$  and so the smallest value of  $l + m + n$  is  $1501 + 1001 + 501 = \boxed{3003}$ .  $\square$

### §7.2.5 Binary Blocks

Source: Harvard-MIT Math Tournament, 2015 C5  
 Proposer: Angry Any#4319 (580933385090891797)  
 Problem ID: 116  
 Date: 2021-01-29

One night while sleeping, Brainy has a vision that the entirety of the new PSC has been found, and that they must now return to the jungle to find their mysterious advisor who will lead them out of this dimension.

The next day, Brainy picks up everybody, but is stopped at Yuchan's house by MODSbot, who plans to trap the PSC in this dimension.

To distract it, Brainy orders MODSbot to write out every integer from 1 to 256 in binary, with a space between each. It takes MODSbot 1 minute to write each unbroken block of 1s and a negligible amount of time to write 0s and spaces.

How long in minutes does Brainy have to collect the rest of the PSC and escape the city?

*Solution.*

Define  $g(0) = 0$ . Call a digit of a number represented in binary "good" if it is 1 and the preceding digit is 0. Then  $g(0) + g(1) + g(2) + \dots + g(255)$  is  $256E(X_8)$  where  $E(X_8)$  is the expected value of the number of "good" digits given that the number is less than  $2^8$ .

By linearity of expectation, the expected number of good digits is  $E(X_8) = E(G_1) + E(G_2) + \dots + E(G_8)$  where  $G_i$  is defined as

$$\begin{cases} 1 & \text{if the } i\text{th digit from the back is good} \\ 0 & \text{otherwise} \end{cases}$$

For example,  $G_3$  for 7 would be 1 but  $G_2$  for 7 would be 0 since there is already a 1 before that. Then we can find  $E(G_i) = \frac{1}{4}$  for all  $i = 1, 2, 3, 4, 5, 6, 7$  since it's just  $\frac{1}{2}$  chance that the  $i$ th digit is 1 multiplied by  $\frac{1}{2}$  chance that the preceding digit is 0. However,  $E(G_8) = \frac{1}{2}$  since the preceding digit is guaranteed to be 0, from which we can find  $E(X_8) = \frac{1}{2} + 7 \cdot \frac{1}{4} = \frac{9}{4}$

Thus  $g(0) + g(1) + g(2) + \dots + g(255) = 256 \cdot \frac{9}{4}$  and  $g(1) + g(2) + \dots + g(255) + g(256) = \boxed{577}$ .  $\square$

*Solution.* [Solution by RishiNandha Vanchi#3379 (562608039224410112)]

Let's say the bot's done printing until  $n - 1$  digits. Now let us append a digit.

We see that the bot will have to print all the numbers it has printed again so that one set of it can be appended with 0 and the other with 1. This takes exactly twice the amount of time taken for  $(n - 1)$  digits.

Whenever the ending digit was a 0 and the appended digit a 1. The bot takes a minute more to print it. It can be seen from some combinatorics that no. of numbers of  $(n - 1)$  digit numbers ending with 0 is exactly  $\binom{2}{1}^{n-2} \cdot 1$  and one resulting number appended with 1 for each of those.

Thus the recurrence is:

$$t_n = 2t_{n-1} + 2^{n-2}$$

1 to 256 would be all 8 digit strings of 1 or 0 along with the number 100000000. Thus the time Bot takes is:

$$t_8 + 1$$

It can be seen that:

$$t_n = 2^k t_{n-k} + k \cdot 2^{n-2} ; t_1 = 1$$

$$t_8 + 1 = \boxed{577}$$

□

### §7.2.6 Funkey Triangles

Source: Original Problem

Proposer: Matteddy#0482 (329956567132930048)

Problem ID: 117

Date: 2021-01-30

Having successfully gathered the PSC and escaped the scheming MODSbot, Brainy manages to lead the group to the jungle. Not far into the jungle, they meet Tan, who tells them that he is their mysterious advisor. He also says that he needs to take them to the last remaining population of wild triangles.

Some of the triangles are a deep shade of red; Tan explains that a triangle  $\triangle ABC$  is red if given its incenter  $I$ , points  $B_1$  and  $C_1$  the intersections of  $BI$  and  $CI$  with  $AC$  and  $AB$  and  $P, Q$  the intersections of line  $PQ$  with  $(ABC)$ ,  $\angle PIQ$  attains the minimal possible value over all acute triangles. Each red triangle has upon its face the value of  $\cos(\angle BAC)$ .

If the product of all numbers written on any triangle in the population for which  $AB = AC$  can be written in the form  $a - b\sqrt{c}$ , where  $a, b, c$  are positive integers with  $c$  squarefree, find  $10000a + 100b + c$ .

*Solution.* [Solution by Matteddy#0482 (329956567132930048)]

Let  $I_b$  and  $I_c$  be the excenters opposite  $B$  and  $C$  respectively. Since  $IAI_cB$  is cyclic, we have  $C_1A \cdot C_1B = C_1I \cdot C_1I_c$ , which implies  $C_1$  lies on the radical axis of  $(ABC)$  and  $(I_bII_c)$ , and so does  $B_1$ . This means that  $B_1C_1$  is the radical axis of those circles, which implies that  $P$  and  $Q$  are their intersection points. Since  $(ABC)$  is the Feuerbach circle of  $\triangle I_bII_c$ , the radius of  $(I_bII_c)$  is twice that of  $(ABC)$  for all triangles, which means that the minimal value of  $\angle PIQ$  happens when  $PQ$  is maximised, so when it is a diameter, and in this case it's easy to see  $\angle PIQ = 150^\circ$ . Now consider the case when  $\triangle ABC$  is funky and isosceles, and let  $O$  be the circumcenter,  $M$  be the midpoint of arc  $BC$  in  $(ABC)$  and suppose the radius of  $(ABC)$  is 1. Then  $2R\sin(\angle MAB) = MB = MI = R \pm OI = R(1 \pm \tan(15^\circ))$ , which gives the possible values of  $\cos(\angle BAC)$  to be  $\sqrt{3} - 1$  and  $3\sqrt{3} - 5$ , and their product is  $14 - 8\sqrt{3}$ . This therefore gives us an answer of

336

□

**§7.2.7 A Lotta Touples!**

---

Source: China Team Selection Test4 2017, P3

Proposer: Matteddy#0482 (329956567132930048)

Problem ID: 118

Date: 2021-01-31

---

The numbers of ordered touples  $(x_1, \dots, x_{100})$  which satisfy:

$$x_1, \dots, x_{100} \in \{1, 2, \dots, 2017\};$$

b)  $2017 \mid x_1 + \dots + x_{100};$

c)  $2017 \mid x_1^2 + \dots + x_{100}^2$

can be expressed as a product of its prime factors,  $p_1^\alpha p_2^\beta \dots$ , find  $p_1\alpha + p_2\beta + \dots$

*Solution.*

Art of Problem Solving write-ups

Answer: 197666

□



## §8 A New Chapter (Season 8)

### §8.1 Week 1

#### §8.1.1 Smallest Non-Factor

---

*Source: Senior Mathematical Challenge 1999 Q9*

*Proposer: sjbs#9839 (434767660182405131)*

*Problem ID: 119*

*Date: 2021-02-01*

*Difficulty: Beginner*

---

Let  $N = 50!$ . What is the smallest positive integer which does not divide  $N$ ?

*Solution.*

The question can be rephrased as 'what is the smallest prime number greater than 50'. To which the answer is 53 □

## §8.1.2 Product of Radii

Source: Senior Mathematical Challenge 2003 Q24

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 120


Date: 2021-02-02

Difficulty: Beginner

Let  $AOB$  be an isosceles right-angled triangle drawn in a quadrant of a circle of radius unit 1. The largest possible circle drawn in the minor segment cut by the line  $AB$  has radius  $r$ . The radius of the inscribed circle of the triangle  $AOB$  is  $R$ . Given that the value of  $Rr$  can be written in the form  $\frac{a-b\sqrt{c}}{d}$ , where  $a, b, c, d$  are positive integers and  $c$  is square-free.

What is the value of  $a^2 + b^2 + c^2 + d^2$ ?

Solution.



Observe that  $\epsilon + 2R + 2r = 1$  and  $\epsilon + 2R = \frac{1}{\sqrt{2}}$ , therefore we have  $r = \frac{\sqrt{2}-1}{2\sqrt{2}}$ . Then by the sine rule,  $\frac{\sin(45)}{R} = \frac{\sin(90)}{R+\epsilon} \Rightarrow R = \frac{1}{\sqrt{2}+2}$ , this gives  $Rr = \frac{3-2\sqrt{2}}{4}$ , hence our answer is  $3^2 + 2^2 + 2^2 + 4^2 = \boxed{33}$   $\square$

### §8.1.3 Two-to-One Functions

Source: Folklore

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 121

Date: 2021-02-03

Difficulty: Easy

Define a function  $f : \{1, 2, \dots, 12\} \rightarrow \{1, 2, \dots, 6\}$  such that for every  $y \in \{1, 2, \dots, 6\}$  there exists exactly two elements  $x_1, x_2 \in \{1, 2, \dots, 12\}$  such that  $f(x_1) = f(x_2) = y$ . How many such functions are there which map  $\{1, 2, \dots, 12\}$  to  $\{1, 2, \dots, 6\}$  with the described property?

(A four-function calculator may be used)

*Solution.*

This well known property can be derived simply by considering how we can choose two elements in  $\{1, 2, \dots, 12\}$  and enumerating that. Let the number of such functions be  $F$ , then we have:

$$\begin{aligned} F &= \prod_{i=1}^{12} \binom{2i}{2} \\ &= \frac{12!}{2^6} \\ &= \boxed{7484400} \text{ such functions} \end{aligned}$$

□

**§8.1.4 Modular Powers**

---

Source: New Zealand Mathematical Olympiad Round 1, 2019 Q4 (Adapted)

Proposer: brainysmurfs#2860 (281300961312374785)

Problem ID: 122

Date: 2021-02-04

Difficulty: Medium

---

Find the remainder when  $122^{2020} - 102^{2020} - 21^{2020}$  is divided by 2020.

*Solution.*

Let  $X = 122^{2020} - 102^{2020} - 21^{2020}$ .

Note that  $2020 = 20 \times 101$ . In particular, considering the expression mod 20 we get  $X = 2^{2020} - 2^{2020} - 1^{2020} \equiv -1 \pmod{20}$ , and considering it mod 101 we get  $X = 21^{2020} - 1^{2020} - 21^{2020} \equiv -1 \pmod{101}$ .

In particular, by Chinese Remainder Theorem we get  $X \equiv -1 \pmod{2020}$  which means that the remainder on division by 2020 of  $X$  is 2019.

*Note: The choice of 2020 in the exponent is not special.*

□

## §8.1.5 Australian Nim

Source: AMO 2020 P2

Proposer: ChristopherPi#8528 (696497464621924394)

Problem ID: 123

Date: 2021-02-05

Difficulty: Medium

sjbs and Brainy are playing a game. First, they'll use a random number generator to generate three random positive integers. Then, three piles of stones will magically appear before them; the number of stones in each pile are the numbers generated previously. Then they'll take turns, with sjbs going first. On a player's turn, they can pick one pile and split it into either two or three nonempty piles, throwing the rest of the stones in the other two piles away. The game ends when a player can't make a move. Suppose both players play perfectly like the smart people they are - then if the probability that sjbs wins is  $m/n$ , with  $\gcd(m, n) = 1$  and  $m, n$  positive integers, find  $m + n$ .

*Solution.*

Call a pile *perilous* if it contains a number of stones equal to  $3k + 1$  where  $k$  is a nonnegative integer and *safe* otherwise. I claim that a player can force a win, if at their turn they have at least one safe pile in front of them, and lose otherwise. Note that from a safe pile, the turn player can always give the other player either 2 or 3 perilous piles: simply consider splitting a pile of stones equal to  $3k + 2$  into two piles, one of 1 stone and 1 of  $3k + 1$  stones, which are both perilous, or a pile equal to  $3k$  stones into three piles, two of 1 stone and 1 of  $3k - 2 = 3(k - 1) + 1$  stones which are all perilous. Also note that from a perilous pile, it is impossible to leave only perilous piles behind, since two or three perilous piles imply that the original split pile was safe.

So if a player has at least one safe pile in front of them on their turn, they may make two or three perilous piles; then their opponent must return them at least one safe pile and they can repeat this; noting that the total number of stones in game is strictly decreasing so the game must end and that the losing state (all piles with 1 stone) is all perilous piles shows that they will win. And clearly if a player only has perilous piles before them then they must give their opponent at least one safe pile and thus their opponent wins. Thus sjbs wins if at least one of the three piles generated at the start is safe, and brainy wins if they are all perilous. The probability of brainy's win is thus  $\frac{1}{3} \cdot \frac{1}{3} \cdot \frac{1}{3} = \frac{1}{27}$  so the probability sjbs wins is  $1 - \frac{1}{27} = \frac{26}{27}$  and so the answer is  $26 \cdot 100 + 27 = \boxed{2627}$ .  $\square$

## §8.1.6 Another Geo Config

Source: Italian Team Competition Final 2020

Proposer: Matteddy#0482 (329956567132930048)


Problem ID: 124

Date: 2021-02-06

Difficulty: Challenging

For a triangle  $ABC$ , let  $P$  be on  $AB$ , and  $X$  be the circumcenter of  $APC$ ,  $Y$  be the circumcenter of  $BPC$ , and  $Z$  be the intersection of  $AX$  and  $BY$ . Given that  $AB = 91$ ,  $BC = 104$ , and  $CA = 65$ , what is the length of  $CZ$ ?

*Solution.*



By simple angle chasing  $\triangle CXA$  and  $\triangle CYB$  are similar, we have  $\angle ZAC = \angle XAC = \angle YBC = \angle ZBC$ , so  $Z$  is on  $\odot(ABC)$ . Standard calculations on  $\triangle ACX$  and  $\triangle ABC$  give  $\sin(\angle ZAC) = \frac{3\sqrt{3}}{14}$  and  $R_{\odot(ABC)} = \frac{91\sqrt{3}}{3}$ , so  $CZ = 2R_{\odot(ABC)}\sin(\angle ZAC) = 39$   $\square$

## §8.1.7 Product of Root Differences

Source: Original

Proposer: tanoshii#3160 (300065144333926400)

Problem ID: 125

Date: 2021-02-07

Difficulty: Challenging

Let  $\alpha_1, \alpha_2, \dots, \alpha_{2021}$  be the roots of  $x^{2021} + 20x^2 + 21$ . Find

$$\prod_{1 \leq i < j \leq 2021} (\alpha_i - \alpha_j)$$

*Solution.* [Write up by epicxtroll#6007 (300008472978653184)]

We solve for general  $f(x) = x^n + ax^2 + b$ , where  $n \equiv 1 \pmod{4}$ . Let  $P$  be our product, we have

$$P = \prod_{i < j} (\alpha_i - \alpha_j)^2 = \prod_{i \neq j} (\alpha_i - \alpha_j)$$

since we flipped the signs of  $\sum_{k=1}^{n-1} k \equiv 0 \pmod{2}$  factors. Fix  $i$ ; our product becomes

$$P = \prod_{i=1}^n \prod_{j \neq i} (\alpha_i - \alpha_j) = \prod_{i=1}^n \frac{\prod_{j=1}^n (\alpha_i - \alpha_j)}{\alpha_i - \alpha_i} = \prod_{i=1}^n \frac{f(\alpha_i)}{\alpha_i - \alpha_i}$$

after writing  $x^n + ax^2 + b = \prod_{i=1}^n (x - \alpha_i)$ . Although division by zero is undefined, since polynomials are continuous functions we can use L'Hopital's Rule to write the product as a limit, then use calculus to get

$$P = \prod_{i=1}^n \lim_{x \rightarrow \alpha_i} \frac{f(x)}{x - \alpha_i} = \prod_{i=1}^n n\alpha_i^{n-1} + 2a\alpha_i = -bn^n \prod_{i=1}^n \left( \alpha_i^{n-2} + \frac{2a}{n} \right).$$

Let  $\omega$  be a primitive  $(n-2)^{\text{nd}}$  root of unity and  $r = \left(\frac{2a}{n}\right)^{\frac{1}{n-2}}$ . The roots of  $x^{n-2} + r^{n-2}$  are  $-r\omega^k$  where  $0 \leq k < n-2$ , so we can rewrite our product as

$$P = -bn^n \prod_{i=1}^n \prod_{j=0}^{n-2} (\alpha_i + r\omega^j) = -bn^n \prod_{j=0}^{n-2} \prod_{i=1}^n (\alpha_i + r\omega^j) = bn^n \prod_{j=0}^{n-2} \prod_{i=1}^n (-r\omega^j - \alpha_i) = bn^n \prod_{j=0}^{n-2} f(-r\omega^j).$$

The rest is just computation.

$$\begin{aligned} P &= bn^n \prod_{j=0}^{n-2} (-r^n \omega^{jn} + ar^2 \omega^{2j} + b) = bn^n \prod_{j=0}^{n-2} [r^2 \omega^j (-r^{n-2} + a) + b] \\ &= bn^2 \prod_{j=0}^{n-2} [r^2 \omega^j a(n-2) + bn] = bn^2 \left[ r^{2(n-2)} a^{n-2} (n-2)^{n-2} + bn^{n-2} \right] \\ &= 4a^n b(n-2)^{n-2} + bn^n \\ &\equiv \boxed{821} \pmod{10000} \end{aligned}$$

□

*Solution.* [Write up by AiYa#2278 (675537018868072458)]

Recall the fundamental theorem of symmetric polynomials: *every symmetric polynomial of  $n$  variables can be written as a function of the  $n$ -variable elementary symmetric polynomials.* Since  $P = \prod_{i \neq j} (\alpha_i - \alpha_j)$  is symmetric in  $n$  variables and all elementary symmetric polynomials in  $n$  variables are zero except for  $e_{n-2}(\alpha) = -a$  and  $e_n(\alpha) = -b$  (where  $e_k(\alpha)$  is the  $k$ -th elementary symmetric sum),  $P$  can be written as a sum of  $(-a)^x(-b)^y$ .  $P$  has degree  $n(n-1)$ ,  $-a$  has degree  $n-2$ , and  $-b$  has degree  $n$  so we solve  $(n-2)x + ny = n(n-1)$  to find our suitable powers of  $-a$  and  $-b$ . Solving, we get  $(x, y) = (n, 1); (0, n-1)$  so  $P = Qa^nb + Rb^{n-1}$ . It remains to choose easy  $a, b$  to find  $P, Q$ ; an obvious choice is  $(a, b) = (0, 1)$  and we have

$$P = \prod_{i=1}^n n\alpha_i^{n-1} = n^n \iff R = n^n.$$

To find  $Q$ , we plug in  $(a, b) = (1, 1)$  and since  $\alpha^n + \alpha^2 + 1 = 0 \iff \alpha^{n-2} = -1 - \frac{1}{\alpha^2}$  we have

$$P = - \prod_{i=1}^n (n\alpha_i^{n-2} + 2) = - \prod_{i=1}^n \left[ (2-n) - \frac{n}{\alpha_i^2} \right] = - \sum_{k=0}^n (2-n)^{n-k} (-n)^k e_k \left( \frac{1}{\alpha^2} \right).$$

We first calculate  $e_k \left( \frac{1}{\alpha} \right)$ ; from substituting  $x \rightarrow \frac{1}{x}$  into  $f(x)$  we get  $e_2 \left( \frac{1}{\alpha} \right) = 1, e_n \left( \frac{1}{\alpha} \right) = -1, e_k = 0$  otherwise except for  $e_0 = 1$  as convention. We solve for  $e_1 \left( \frac{1}{\alpha^2} \right)$  as follows:

$$e_1^2 \left( \frac{1}{\alpha} \right) = e_1 \left( \frac{1}{\alpha^2} \right) + 2e_2 \left( \frac{1}{\alpha} \right) \iff e_1 \left( \frac{1}{\alpha^2} \right) = -2$$

and similarly

$$e_2^2 \left( \frac{1}{\alpha} \right) = e_2 \left( \frac{1}{\alpha^2} \right) + 2e_4 \left( \frac{1}{\alpha} \right) + 2e_1 \left( \frac{1}{\alpha^2} \right) e_2 \left( \frac{1}{\alpha} \right) = 1.$$

For larger values of  $k$ , note that

$$e_k^2 \left( \frac{1}{\alpha} \right) = e_k \left( \frac{1}{\alpha^2} \right) + 2 \sum_{j=0}^k e_j \left( \frac{1}{\alpha^2} \right) e_{2(k-j)} \left( \frac{1}{\alpha} \right)$$

so all of them evaluate to zero except for  $e_n \left( \frac{1}{\alpha^2} \right) = 1$ . The rest is computation.

$$\begin{aligned} P &= (n-2)^n - 2n(n-2)^{n-1} + n^2(n-2)^{n-2} + n^n \\ &= (n-2)^{n-2} (n^2 - 4n + 4 - 2n^2 + 4n + n^2) + n^n \\ &= 4(n-2)^{n-2} + n^n \iff P = 4(n-2)^{n-2} \\ P &= 4a^nb(n-2)^{n-2} + b^{n-1}n^n \\ &\equiv \boxed{821} \pmod{10000} \end{aligned}$$

□



**§8.2 Week 2****§8.2.1 Counting Intersections**

---

Source: Gray Kangaroo 2003 Q24

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 126

Date: 2021-02-08

Difficulty: Beginner

---

Rui draws 10 points on a large piece of paper, making sure that no three points are in a straight line. He then draws a segment joining each pair of points. If Orlo draws a straight line across Rui's diagram, without going through any of Rui's original points, what is the greatest possible number of lines that can be crossed?

*Solution.*

I claim the answer is  $\frac{420^2}{4} = 44100$ .

For a line drawn across Rui's construction splitting the points so that there are  $n$  on one side and  $420 - n$  on the other, clearly there will be  $n(420 - n)$  intersections. I.e.  $\frac{420^2}{4} - \left(n - \frac{420}{2}\right)^2$ . Thus the number of intersections is maximised when  $n = 210$ .

□

### §8.2.2 The Race

Source: UK Senior Kangaroo, 2013 Q20  
 Proposer: sjbs#9839 (434767660182405131)  
 Problem ID: 127  
 Date: 2021-02-09  
 Difficulty: Beginner

Zella and Samantha stand at either end of a straight track. They then run at a constant (not different) speeds to the other end of the track, turn and run back to their original end at the same speed they ran before. On their first leg, they pass each other 20m from one end of the track. When they are both on their return leg, they pass each other for a second time 10m from the other end of the track. How many meters long is the track?

*Solution.*

Let the distance of the track be  $d$ , and the speeds of the two runners be  $u$  and  $v$  respectively. And let  $t_i$  be the time for which the runners are at a particular position. Then on the first leg we must have  $20 = ut_1$  and  $d - 20 = vt_1$ , While on the second leg,  $d + 10 = ut_2$  and  $2d - 10 = vt_2$ . Therefore we have

$$\frac{ut_1}{vt_2} = \frac{ut_2}{vt_2} \Rightarrow \frac{20}{d - 20} = \frac{d + 10}{2d - 10} \quad (1)$$

$$d(d - 50) = 0 \quad (2)$$

Thus we must have  $d = \boxed{50}$

□

### §8.2.3 Remainder of Sums of Squares

Source: *British Mathematical Olympiad, Round 1, 1998 P2*

Proposer: *brainysmurfs#2860 (281300961312374785)*

Problem ID: 128

Date: 2021-02-10

Let  $a_1 = 19$ ,  $a_2 = 98$ . For  $n \geq 1$ , define  $a_{n+2}$  to be the remainder of  $a_n + a_{n+1}$  when it is divided by 100. What is the remainder when

$$a_1^2 + a_2^2 + \cdots + a_{1998}^2$$

is divided by 8?

*Solution.*

Consider the remainders of  $a_i \bmod 4$ : the first six are 3, 2, 1, 3, 0, 3 and it repeats every six. Since  $6 \mid 1998$  it suffices to find the remainder when  $a_1^2 + a_2^2 + \cdots + a_6^2$  is divided by 8; since odd squares are 1 mod 8,  $2^2$  is 4, and  $0^2$  is 0 we have  $4 \cdot 1^2 + 2^2 + 0^2 \equiv \boxed{0} \pmod{8}$ .  $\square$

## §8.2.4 Another 255 Subsets

Source: 2017 HMMT General #8

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 129

Date: 2021-02-11

Difficulty: Medium

Reimu has a collection of the 255 nonempty subsets of  $\{1, 2, 3, 4, 5, 6, 7, 8\}$ . Each minute, she takes two subsets chosen uniformly at random from the collection, and replaces them with either their union or their intersection, with each being equally likely. (The collection can contain repeated sets.) After 254 minutes, she is left with one set. The expected size of this subset can be expressed in the form  $\frac{m}{n}$ , where  $m$  and  $n$  are relatively prime positive integers. Find  $100m + n$ .

*Solution.* [Official Solutions File](#)

Answer: 102655

A formulaic proof of the invariance of the expected size of all remaining subsets is as follows. Let  $S_1, S_2, \dots, S_n$  be the sets in question, where a merge consists of taking two sets and replacing them with either their union or their intersection. If sets  $S_i, S_j$  are merged then the resulting set would be either  $S_i \cap S_j$  or  $S_i \cup S_j$ . Summing over all  $\binom{n}{2}$  pairs of  $i, j$ , our expected average size of sets after merging is

$$\frac{\binom{n}{2} 2S - \sum_{i \neq j} (|S_i \cap S_j| + |S_i \cup S_j|)}{2\binom{n}{2}} = \frac{n(n-1)S - (n-1)S}{n(n-1)} = \frac{S}{n}$$

where  $S = \sum_{i=1}^n |S_i|$ .

□

## §8.2.5 4-5-6 Triangle

Source: AMOC December 2020 Camp Exam 2 (A/G) P4

Proposer: ChristopherPi#8528 (696497464621924394)


Problem ID: 130

Date: 2021-02-12

Difficulty: Hard

Let  $ABC$  be a triangle with side lengths  $AB = 4$ ,  $BC = 5$ ,  $CA = 6$ . Suppose  $P$  is a point inside  $ABC$  such that the circumcenters  $B'$  and  $A'$  of triangles  $ACP$  and  $BCP$  respectively lie outside  $ABC$ , with  $A$ ,  $P$ ,  $A'$  collinear and  $B$ ,  $P$ ,  $B'$  collinear. The line through  $P$  parallel to  $AB$  meets the circumcircles of  $ACP$  and  $BCP$  at  $E$  and  $D$  respectively, where  $D$ ,  $E$ ,  $P$  are distinct. Suppose  $DE^2 - AP = (a - \sqrt{b})/c$  where  $a$ ,  $b$ ,  $c$  are integers and  $b$  is squarefree. Find  $a + 69b + 420c$ .

Solution.



We claim  $P$  is the incenter of  $\triangle ABC$ . By arc lengths,  $\angle PA'B = 2\angle BCP$  and since  $A'P = A'B$  we have  $\angle A'PB = 90 - \angle BCP \iff \angle APB = 90 + \angle BCP$ . However, doing this to the circle centered at  $B'$  gives  $\angle APB = 90 + \angle ACP \iff \angle ACP = \angle BCP = \frac{\angle C}{2}$  and  $\angle APB = 90 + \frac{\angle C}{2}$ . The locus of all points such that  $\angle APB = 90 + \frac{\angle C}{2}$  is  $(AIB)$ , which the  $C$ -angle bisector intersects twice, once at  $I$  and once outside of  $\triangle ABC$ ; since  $P$  is specified to be inside  $\triangle ABC$  we conclude  $P \equiv I$ .

Since  $\angle AEP = \angle ACP = \frac{\angle C}{2}$  and  $\angle CAP = \angle APE = \frac{\angle A}{2}$  we have  $\triangle EAP \cong \triangle CPA$  by AAS; similarly  $\triangle CPB \cong \triangle DBP$  so  $DE = EP + DP = AC + AB = 11$ . Also, the tangents to the incircle from  $A$  have length  $s - a = \frac{5}{2}$  and  $[ABC] = rs \iff r = \frac{\sqrt{7}}{2}$  so  $AP = 2\sqrt{2}$ . 1210202  $\square$

## §8.2.6 “What’s a signed integer?”

Source: Original Problem

Proposer: Constan#6792

Problem ID: 131

Date: 2021-02-13

Difficulty: Challenging

For all pairs  $(n, q)$  where  $n$  is a positive integer,  $q$  is a rational non-integer, and  $n^q - q$  is an integer, find  $\lfloor n + q \rfloor = x$ . The sum of all possible values of  $x$  is the answer. (For example, if we have  $(1, 1)$  and  $(2, 2)$ , then the answer is  $2 + 4 = 6$ )

*Solution.* [Solution by Constan#6792]

Let’s see that if  $n^q$  is rational and  $q$  is positive,  $n^q$  is integer. To do this, suppose it is rational and not integer where  $(a, b) = (c, d) = 1$ .  $n^{\frac{a}{b}} = \frac{c}{d}$   $d^b \cdot n^a = c^b$  Now we consider the exponent of a certain prime in the factorization of  $d, c, n$  which are  $e_d, e_c, e_n$  respectively.  $b \cdot e_d + a \cdot e_n = b \cdot e_c$   $a \cdot e_n \equiv 0 \pmod{b}$   $e_n \equiv 0 \pmod{b}$  So  $n$  is a  $b$ -th power.  $d^b \cdot k^{ab} = c^b$   $d \cdot k^a = c$   $k^a = \frac{c}{d}$  Absurd, because a perfect power must be a whole number.

Now if  $q$  is positive there are no solutions because if  $n^q$  and  $n^q - q$  are integers,  $q$  would be.

Then  $n = -\frac{a}{b}$  with  $a$  and  $b$  co-prime positive integers and  $\frac{1}{n^{\frac{a}{b}}} + \frac{a}{b}$  is an integer.

Now if  $\frac{n_1}{d_1} + \frac{n_2}{d_2} = \frac{n_1 d_2 + n_2 d_1}{d_1 d_2}$  is an integer where  $(n_1, d_1) = (n_2, d_2) = 1$  we have to:  $d_1 | n_1 d_2 \rightarrow d_1 | d_2$ .  $d_2 | n_2 d_1 \rightarrow d_2 | d_1$ . So  $d_1 = d_2$ .

That taking this to what we have tells us that  $n^{\frac{a}{b}} = b$  y  $n^a = b^b$  Where with a very similar reasoning as before,  $b$  is a  $a$ -th power.  $b = k^a$   $n^a = k^{a k^a}$   $n = k^{k^a}$

$$\frac{1}{(k^{k^a})^{\frac{a}{k^a}}} + \frac{a}{k^a} = \frac{a+1}{k^a}$$

Where:  $k^a \leq a + 1$  If  $k \geq 3$   $3^a \leq k^a \leq a + 1$  has no solutions. Then  $k \leq 2$  where if  $k = 1$ ,  $\frac{a}{b}$  would be an integer, so  $k = 2$ .  $2^a \leq a + 1$  Where the only solution is  $a = 1$  that gives us the triple:  $(k^{k^a}, k^a, a) = (4, 2, 1)$ .

Thus  $(n, q) = (4, -\frac{1}{2})$  the only solution. So the answer is 4.  $\square$

**§8.2.7 Diagonals in a Convex 1001-gon**

---

*Source: Sharygin 2019 Finals Grade 8 P8*

*Proposer: TaesPadhihary#8557 (665057968194060291)*

*Problem ID: 132*

*Date: 2021-02-14*

*Difficulty: Challenging*

---

What is the least positive integer  $k$  such that, in every convex 1001-gon, the sum of any  $k$  diagonals is greater than or equal to the sum of the remaining diagonals?

*Solution.*

[Official Solution, page 6 P8](#)

Answer:



## §10 CCCC After Math (Season 10)

### §10.1 Week 1

#### §10.1.1 EpicXtroll

---

*Source: Original*

*Proposer: epicxtroll#6007 (300008472978653184)*

*Problem ID: 133*

*Date: 2021-02-15*

*Difficulty: Beginner*

---

How many possibilities are there for the total score among all participants at the end of a standard QoTD season? (Assume that no rounding occurs. Recall that a standard QoTD season has 14 questions.)

*Solution.*

The answer is 15: if  $0 \leq N \leq 14$  is the number of questions solved (by anyone), then the total score is  $1000N$ . □



## §10.1.2 Similar Triangles

Source: Original

Proposer: Matteddy#0482 (329956567132930048)

Problem ID: 134


Date: 2021-02-16

Difficulty: Beginner

Let  $ABC$  be a triangle with  $AB = AC$  and  $BC = 10$ . Construct  $D$  and  $E$  outside  $ABC$  closer to  $B$  and  $C$  respectively such that  $\triangle DAB \sim \triangle ABC \sim \triangle EAC$ .

If  $DE = 45$ , then what is  $AB$ ?

Solution.



Note that  $\angle DAB = \angle ACB = \angle ABC$  and hence  $AD \parallel BC$ . Similarly  $AE \parallel BC$  and so  $A$  is the midpoint of  $DE$ , and  $A, D, E$  are collinear.

So  $AD = 22.5$  and hence  $AB = BC * \sqrt{\frac{AD}{BC}}$  by similarity, which is  $\boxed{15}$ . □

### §10.1.3 Factorial Sums and Divisors

Source: Folklore

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 135

Date: 2021-02-17

Difficulty: Easy

What is the largest prime number which divides

$$0! + 1! \times 1 + 2! \times 2 + 3! \times 3 + \cdots + 159! \times 159 + 160! \times 160$$

*Solution.*

We proceed by using the identity  $n! + n! \times n = (n+1)!$  and iterating:

$$0! + 1! \times 1 + \cdots \tag{3}$$

$$2! + 2! \times 2 + \cdots \tag{4}$$

$$3! + 3! \times 3 + \cdots \tag{5}$$

$$\cdots \tag{6}$$

$$161! \tag{7}$$

As neither 161 nor 159 are prime, the largest prime divisor of the sum is 157 □

*Solution.*

Note that  $(n+1)! - 1$  counts the non-identity permutations of  $(1, 2, \dots, n+1)$ . Consider a permutation where  $(1, 2, \dots, n-k)$  are fixed points but  $n-k+1$  is not a fixed point. So  $n-k+1$  can go in the  $n-k+2^{\text{nd}}$  to  $n+1^{\text{st}}$  spots, which is a total of  $k$  possibilities; the remainder of the  $k$  terms can be permuted willy-nilly. Thus, there  $k \cdot k!$  such possibilities, and summing over  $k$  from 1 to  $n$  (since  $k = n+1$  gives the identity permutation) yields the identity. 157 □

## §10.1.4 Brainy's Passcode

Source: Italian team competition

Proposer: HoboSas#3200 (310725130097786880)

Problem ID: 136

Date: 2021-02-18

Difficulty: Medium

Brainy forgot the numerical unlock code of his Nokia once again... This time he remembers that it is 69-digits long and the leading digit is greater or equal than the sum of all the remaining digits. Find the total number of possible codes.

*A computational aid may be used to calculate the final answer.*

*Solution.*

Let  $n$  be the number of digits of the code, in our case  $n = 69$  and let  $k$  be the leading digit, clearly  $k \in \{1, 2, \dots, 9\}$ . Let  $i$  be the sum of the remaining digits, where  $i \in \{1, 2, \dots, k\}$ . Using the stars and bars method (also called sticks and stones or Tonys and Wangs), the number of possible codes (with  $k$  and  $i$  fixed) is

$$\binom{(n-1) - 1 + i}{(n-1) - 1}$$

Cycling this for all possible values of  $k$  and  $i$  we get the following expression

$$\sum_{k=1}^9 \sum_{i=0}^k \binom{n-2+i}{n-2}$$

Using the Hockey-stick identity 2 times in a row we get

$$\begin{aligned} \sum_{k=1}^9 \sum_{i=0}^k \binom{n-2+i}{n-2} &= \sum_{k=1}^9 \binom{k+n-1}{k} \\ &= \binom{n+9}{9} - 1 \end{aligned}$$

Plugging in  $n = 69$  we get

$$\binom{78}{9} - 1 = \boxed{182364632449}.$$

□

*Solution.* [Write up by AiYa#2278 (675537018868072458)]

Let the  $k^{\text{th}}$  digit from the left be  $d_k$ . Our constraints become  $9 \leq d_1 \leq \sum_{k=2}^{69} d_k$ ; let  $y = 9 - d_1$  and  $x = d_1 - \sum_{k=2}^{69} d_k$ . Our constraints become

$$9 = d_1 + y = x + y + \sum_{k=2}^{69} d_k$$

so it suffices to count the ways to distribute 9 objects among 70 people. Letting 69 "sticks" divide the 9 "stones" into 70 parts, we see that this is  $\binom{78}{9}$  then subtract 1 because a string of 69 zeros is invalid.

$$\boxed{182364632449}$$

□

## §10.1.5 Production Loop

Source: Original/Folklore

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 137

Date: 2021-02-19

Difficulty: Hard (CN4)

Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be a function defined as follows:

$$f(n) = 2n + 1 - 2^{\lfloor \log_2 n \rfloor + 1}$$

and let  $f^a(n) = f(f^{a-1}(n))$ . Let  $t(n)$  be the smallest positive integer such that there exists a positive integer  $N$  such that  $f^{t(n)}(n) = f^{t(n)+N}(n)$ . Determine the remainder when  $\sum_{n=2^{2020}}^{2^{2021}} f^{t(n)}(n)$  is divided by 1009.

(A scientific calculator can be used)

*Solution.*

Consider what  $f$  does to  $n$  in binary:  $2n + 1$  concatenates a 1 to the end of  $n$ , while  $2^{\lfloor \log_2 n \rfloor + 1}$  subtracts the concatenation of 0 to the end of  $n$  from  $2n + 1$ . This can be seen as cycling the leading 1 of  $n$  to the end of  $n$ ; for instance  $f(6 = 101_2) = 11_2 = 3$ . After applying  $f$  enough times, we're left with the number of ones present in  $n$ . We now count the number of integers  $2^{2020} \leq n < 2^{2021}$  which have  $k$  ones present in their binary representation. All these numbers are 2021 digits long, so we want to distribute  $k - 1$  ones among 2020 digits; there are  $\binom{2020}{k-1}$  ways to do this and each such  $n$  contributes  $2^k - 1$  to the sum. Remembering that  $f^{t(n)}(n) = 1$  when  $n = 2^{2021}$ , it remains to sum

$$\begin{aligned} 1 + \sum_{k=1}^{2021} \binom{2020}{k-1} (2^k - 1) &= 1 + 2 \sum_{k=1}^{2021} \binom{2020}{k-1} 2^{k-1} - \sum_{k=1}^{2021} \binom{2020}{k-1} \\ &= 1 + 2 \cdot 3^{2020} - 2^{2020} \\ &\equiv 1 + 2 \cdot 81 - 16 \pmod{1009} \\ &= \boxed{147} \end{aligned}$$

where we have used  $(1 + 2)^n = \sum_{k=0}^n \binom{n}{k} 2^k = 3^n$  to help simplify. □

## §10.1.6 Weird polynomial

Source: NYCMT, 2020 P9 of 10

Proposer: TaesPadhiary#8557 (665057968194060291)

Problem ID: 138

Date: 2021-02-20

Let  $p = 2053$  be a prime. For positive integers  $x$  let

$$f(x) = x^{400} + x^{326} + x^{200} + x^{126} + 2$$

and let  $g(x)$  denote the unique integer  $0 \leq y \leq p-1$  such that  $y^{1759} - x$  is divisible by  $p$ . Compute the remainder when

$$\sum_{i=1}^{2052} g(f(i))$$

is divided by 2053.

*Solution.*

To find  $y$ , we want to raise  $y^{1759} \equiv x \pmod{p}$  to a power  $k$  such that  $1759k \equiv 1 \pmod{p-1}$ ; solving we get  $k \equiv 7 \pmod{p-1}$  and so  $y \equiv x^7 \pmod{p}$ . It remains to sum  $f(i)^7 \pmod{p}$ . Since  $g, g^2, \dots, g^{p-1}$  where  $g$  is a generator is a permutation of the nonzero residues mod  $p$  we find that

$$\sum_{i=1}^{p-1} i^k = \sum_{i=1}^{p-1} g^{ik} = g^k \left( \frac{g^{(p-1)k} - 1}{g^k - 1} \right) \equiv 0 \pmod{p}$$

unless  $p-1|k$  in which case each of the  $p-1$  terms is 1 for a sum of  $-1 \pmod{p}$ . Since the maximum coefficient in  $f(x)^7$  is  $7 \cdot 400 = 2800$  it suffices to find the coefficient of  $x^{2052}$  and the constant term. Using multinomial expansion, this is equivalent to finding nonnegative integers  $(a, b, c, d, e)$  such that  $400a + 326b + 200c + 126d + 0e = 2052 \iff 200a + 163b + 100c + 63d = 1026$  and  $a + b + c + d + e = 7$ . Notice  $a > 1$  since  $200 \cdot 1 + 163 \cdot 6 < 1026$  and  $a < 5$  since  $200 \cdot 5 = 1000$  and no combination of 160, 100, 63 will make that sum to 1026. Also, reducing mod 9 yields  $2a + b + c \equiv 0 \pmod{9} \iff a + 7 \equiv d + e \pmod{9}$ . When  $a = 2$ , we are forced  $d, e = 0$  so  $163b + 100c = 626 \iff (a, b, c, d, e) = (2, 2, 3, 0, 0)$ . When  $a = 3$ , we are forced  $d = 0, 1$  so either  $163b + 100c = 426, 363 \iff (a, b, c, d, e) = (3, 2, 1, 0, 1), (3, 1, 2, 1, 0)$ . When  $a = 4$ , we have  $163b + 100c + 63d = 226 \iff (a, b, c, d, e) = (4, 1, 0, 1, 1), (4, 0, 1, 2, 0)$ . Remembering the constant term is  $2^7$ , it remains to sum

$$-\binom{7}{2, 2, 3, 0, 0} - 2\binom{7}{3, 2, 1, 0, 1} - \binom{7}{3, 1, 2, 1, 0} - 2\binom{7}{4, 1, 0, 1, 1} - \binom{7}{4, 0, 1, 2, 0} - 2^7 \equiv \boxed{1983} \pmod{2053}.$$

□

## §10.1.7 Weird triangle

Source: Original

Proposer: tanoshii#3160 (300065144333926400)

Problem ID: 139

Date: 2021-02-21

Difficulty: Challenging (G7)

Let  $ABC$  be a  $13 - 14 - 15$  triangle, with  $AC = 14$ . Points  $X, Y$  satisfy

$$CX - BX = BX - AX = 1 = AY - BY = BY - CY.$$

The value of  $XY$  can be written in the form  $\frac{a\sqrt{b}}{c}$ , where  $b$  is not divisible by the square of any prime and  $a$  and  $c$  are relatively prime positive integers. Find  $10000a + 100b + c$ .

Solution.

Let  $I$  be the incenter of  $\triangle ABC$ . Construct three circles, each centered at one of  $A, B, C$ . Note that these three circles touch each other at the three intouch points of  $\triangle ABC$ . Thus their radii are 6, 7 and 8. WLOG assume  $AB = 13$ . Let  $W$  be the center of the unique circle externally tangent to our three circles at  $A, B, C$ , and  $Z$  the center of the unique circle internally tangent to our three circles at  $A, B, C$ . By definition, since the radius of the circle at  $C$  is 8, the radius of the circle at  $B$  is 7 and the radius of the circle at  $A$  is 6,  $W$  and  $Z$  must be precisely  $X$  and  $Y$ . Now consider an inversion with respect to the incircle. Let the intersection of the line  $AI$  and the circle at  $A$  further from  $I$  be  $D$ , and the one closer to  $I$  be  $E$ . Then if the radius of the circle at  $A$  is  $R$ , and the inradius is  $r$ ,  $DI \cdot EI = (AI - R)(AI + R) = AI^2 - R^2 = R^2 + r^2 - R^2 = r^2$ , so by definition  $D$  and  $E$  swap under our inversion. Thus the circle at  $A$  must map to itself under our inversion, and so do the circles at  $B$  and  $C$ . Then since inversion preserves tangency, the circle at  $X$  externally tangent to our three circles must swap with the circle at  $Y$  internally tangent to our three circles.

Now suppose a line through  $I$  meets the circle at  $X$  at  $M, N$  and the circle at  $Y$  at  $P, Q$ . Then since the image of  $M$  must lie on the ray  $MI$ , and also lie on the circle at  $Y$  since  $M$  lies on the circle at  $X$ , WLOG  $M$  and  $P$  swap, and then so do  $N$  and  $Q$ . Thus  $IM \cdot IP = r^2 = IN \cdot IQ$ , and by power of a point  $IM \cdot IN = R_x^2 - IX^2$  and  $IN \cdot IQ = R_y^2 - IY^2$  where  $R_x$  and  $R_y$  are the radii of the circles at  $X$  and  $Y$ . Since  $IM \cdot IP$  is independent of  $M$  and  $P$  we find that  $I$  is the insimilicenter of the circles at  $X$  and  $Y$  and therefore lies on the line segment  $XY$ . Thus if we set  $IX = cR_x$  and  $IY = cR_y$  then  $r^4 = R_x^2 R_y^2 (1 - c^2)^2$ ; by direct calculation (the inradius formula for  $r$  and Descartes' Kissing Circles Theorem for  $R_x$  and  $R_y$ ) we obtain four values for  $c$ . However, two of these values are negative and a third is greater than 1 (we know that clearly  $IX$  and  $IY$  are smaller than  $R_x$  and  $R_y$  by simply drawing the diagram in this case), so the only acceptable value of  $c$  is  $c = \frac{\sqrt{37}}{42}$ . Then by direct calculation we obtain that  $XY = IX + IY = cR_x + cR_y = \frac{672\sqrt{37}}{1727}$  so the answer is 6725427 as required.  $\square$

*Solution.*

An elementary, computationally feasible analytic solution is still possible using Cartesian coordinates. From the given condition, if we let  $BX = d$  then  $AX = d - 1$ ,  $CX = d + 1$ . Consider circles  $\omega_A, \omega_B, \omega_C$  centered at  $A, B, C$  with radii  $d - 1, d, d + 1$  respectively. Let  $A = (-5, 0), B = (0, 12), C = (9, 0)$ . Then the radical axis of  $\omega_A$  and  $\omega_C$  is given by

$$(x + 5)^2 + y^2 - (d - 1)^2 = (x - 9)^2 + y^2 - (d + 1)^2 \iff 7x = 14 - d$$

and the radical axis of  $\omega_A$  and  $\omega_B$  is given by

$$(x + 5)^2 + y^2 - (d - 1)^2 = x^2 + (y - 12)^2 - d^2 \iff 5x + 12y = 60 - d$$

so the radical center of  $\omega_A, \omega_B, \omega_C$  satisfies

$$(x, y) = \left( \frac{14 - d}{7}, \frac{350 - 2d}{84} \right).$$

This point must be on  $\omega_B$ , so plugging in and simplifying we have

$$\left( \frac{14 - d}{7} \right)^2 + \left( \frac{350 - 2d}{84} - 12 \right)^2 = d^2 \iff \frac{1727}{1764}d^2 + \frac{25}{126}d - \frac{2353}{36} = 0.$$

Solving this quadratic would be hell, but we don't need to. Let the two roots be  $r, s$  with  $r$  positive and  $s$  negative. Consider a point  $P$  with  $PA = |s - 1|, PB = |s|, PC = |s + 1|$ ; since  $s < -1$  we have  $PA = -s + 1, PB = -s, PC = -s - 1$ . This satisfies  $AP - BP = BP - CP = 1$ , so  $P \equiv Y$ ! Thus, it remains to find

$$XY = \sqrt{\frac{(r - s)^2}{7^2} + \frac{(r - s)^2}{42^2}} = \frac{r - s}{7} \sqrt{\frac{37}{36}}$$

and now the miraculous

$$(r - s)^2 = (r + s)^2 - 4rs = \frac{1764^2}{1727^2} \left( \frac{25^2}{126^2} + \frac{2353}{9} \cdot \frac{1727}{1764} \right) = 256 \cdot \frac{1764^2}{1727^2} \iff r - s = 16 \cdot \frac{1764}{1727}$$

$$\text{so } XY = \frac{672\sqrt{37}}{1727} \iff \boxed{6725427}.$$

□

**§10.2 Week 2****§10.2.1 Real Square Roots**

---

*Source: Purple Comet! Math Meet, High School, 2020 P4*

*Proposer: TaesPadhiary#8557 (665057968194060291)*

*Problem ID: 140*

*Date: 2021-02-22*

*Difficulty: Beginner*

---

Let  $\mathcal{S}$  be the set of integers  $n$  such that

$$\sqrt{\frac{(2021 - k)^2}{2021 - k^2}}$$

is a real number.

Compute the sum of all elements in  $\mathcal{S}$ .

*Solution.*

The numerator is always real, so we seek  $k$  such that  $2021 - k^2 > 0$ ; however if  $k$  satisfies this inequality so does  $-k$ . Then observe  $k = 2021$  makes the numerator 0, so it is real too. 2021. □



### §10.2.2 "What's a positive integer?"

Source: Original Problem

Proposer: TaesPadhihary#8557 (665057968194060291)

Problem ID: 141

Date: 2021-02-23

Difficulty: Beginner

Let  $n$  be a positive integer. Find the number of positive divisors of  $69420^{69}$  which are of the form  $4n + 1$ .

*Solution.* [Solution by TaesPadhihary#8557 (665057968194060291)]

Note that  $69420^{69} = 2^{138} \cdot 3^{69} \cdot 5^{69} \cdot 13^{69} \cdot 89^{69}$ . From these, only 3 is a prime which is equivalent to 3 (mod 4) and others, i.e. 5, 13, 89 are all 1 (mod 4). Therefore, the number of ways to choose a divisor of the form  $4n + 1$  is  $35 \times 70 \times 70 \times 70^5$ . But, since  $n$  is positive integer, so 1 can not be a chosen divisor. Hence, the desired answer is  $35 \cdot 70^3 - 1 = \boxed{12004999}$ .  $\square$

---

<sup>5</sup>35 ways being for the even powers of 3 and 70 otherwise

## §10.2.3 Converging Tangent Circles

Source: Original Problem


Proposer: HoboSas#3200 (310725130097786880)

Problem ID: 142

Date: 2021-02-24

Difficulty: Easy

In the following diagram every circle has two internally tangent circles also tangent to each other. The radius of the biggest circle is  $\frac{1}{\sqrt{\pi}}$ . If the area of the black region can be expressed as  $\frac{a}{b}$ , compute  $100a + b$ .



Solution. [Solution by HoboSas#3200 (310725130097786880)]

Let  $r_n$  and  $A_n$  be the radius and the area of the  $n$ th circle (in descending order). From construction we have  $r_{n+1} = \frac{r_n}{2}$  which leads to  $r_n = \frac{r_0}{2^n}$ , where  $r_0 = \frac{1}{\sqrt{\pi}}$ . Looking at the desired area  $A_\zeta$  as difference of black and white circles areas, we get the following:

$$A_\zeta = \sum_{n=0}^{\infty} (-2)^n A_n \quad (8)$$

$$= \sum_{n=0}^{\infty} (-2)^n (r_n)^2 \pi \quad (9)$$

$$= (r_0)^2 \pi \sum_{n=0}^{\infty} \left(-\frac{1}{2}\right)^n \quad (10)$$

$$= \left(\frac{1}{\sqrt{\pi}}\right)^2 \pi \frac{1}{1 - (-\frac{1}{2})} \quad (11)$$

$$= \frac{2}{3} \quad (12)$$

Which means 203 is the solution. □

Solution. [Solution by flame#6784 (185229437787176962)]

Let  $B$  be the total black area; the total white area is then  $1 - B$ . Looking at the two biggest white circles, we see that each of them is similar to the big circle by a factor of  $\frac{1}{2}$  with colors inverted; thus the total black area inside each of the two white circles is  $\frac{1-B}{4}$ . Since the black area outside the two big white circles is  $\frac{1}{2}$  we have

$$B = \frac{1}{2} + \frac{1-B}{2} \iff B = \frac{2}{3} \iff \boxed{203}.$$

□

**§10.2.4 Maze Escape**

---

*Source: 2018 HMMTNov Guts 18*

*Proposer: AiYa#2278 (675537018868072458)*

*Problem ID: 143*

*Date: 2021-02-25*

*Difficulty: Medium (C3)*

---

MODSbot has trapped Brainy at the southwest entrance of a two-by-fifteen grid of cells. To escape, Brainy has to enter into the grid from the southwest cell and exit through the northeast cell. However, MODSbot can fill any subset of the 20 cells with cement, effectively blocking off those cells. Out of all  $2^{30}$  possible grid configurations, in how many of them can Brainy escape?

*Solution.*

Follow the [official solution](#) to get the recursion for  $a_n$  and  $b_n$ . Iterating 15 indices of  $a$  and  $b$  is inefficient; for efficient computation we let  $c_n = a_n + b_n$  and note that  $c_{n+1} = 2c_n + c_{n-1}$ ; either directly recursing or finding a closed form yields  $c_{15} = 665857$ . By inspection, we have  $a_n = b_n - 1$  so  $a_n = \boxed{332928}$ .  $\square$

**§10.2.5 Primes divide powers of 5**

---

Source: China National Olympiad 2009, P2

Proposer: Matteddy#0482 (329956567132930048)

Problem ID: 144

Date: 2021-02-26

Difficulty: Hard

---

Suppose  $\mathcal{S}$  is the set of all pairs  $(p, q), p \leq q$  such that

$$pq \mid 5^p + 5^q.$$

Find

$$\sum_{(p,q) \in \mathcal{S}} (1000p + q).$$

*Solution.*

**AoPS Solution**

Answer: 14326

□

## §10.2.6 tRiVial

Source: Original Problem

Proposer: QuantumSigma#9996

Problem ID: 145

Date: 2021-02-27

Difficulty: Challenging

Consider an integer-valued function  $f(x, y)$  defined on the lattice points of the  $x - y$ -plane. The value of  $f$  at each point is average its 4 neighbours. The difference in  $f$  is less than 70000 for any pair of lattice points a distance of exactly 2021 apart. If  $f(a, b + c) - f(a, b) = 1780$  for some integers  $(a, b, c)$ , find the sum of all possible  $|c|$ .

*Solution.* [Solution by QuantumSigma#9996] 3738. Let  $F$  be the set of functions  $g : \mathbb{Z}^2 \rightarrow \mathbb{Z}$  fulfilling the 'average of neighbours' criteria.

1. For any  $g \in F$ ,  $g$  is either constant or unbounded (Klarijōn#0829, modified). Proof: Assume there exist 2 adjacent points  $(A, B)$  such that  $g(A) < g(B)$ . Then there exists point  $C$  adjacent to  $B$  such that  $g(B) < g(C)$ . Repeating ad infinitum, there exists an infinite strictly monotonic chain of points. Thus  $g$  is unbounded.
2. Define  $P$  and  $Q$ :

$$Pf(x, y) = f(x + 2021, y) - f(x, y)$$

$$Qf(x, y) = f(x, y + 2021) - f(x, y)$$

Then for any  $g \in F$ ,  $P\{g\}, Q\{g\} \in F$  (trivial)

3. For any  $f \in F$  such that  $P\{f\}, Q\{f\}$  are bounded,  $f$  is linear. Proof: By (1) and (2),  $Pf, Qf$  are constant. Let  $h = 2021f - Pf x - Qfy$ , then  $h \in F$ . Since  $h$  is doubly periodic (trivial), it is bounded, and thus constant. Then  $f = \frac{Pf x + Qfy + h}{2021}$ .

It follows that  $|c|$  is a factor of 1780. The problem is then trivial. □

**§10.2.7 Lamps in a Circle**

Source: *Olympic Revenge*, 2013 P5

Proposer: ChristopherPi#8528 (696497464621924394)

Problem ID: 146

Date: 2021-02-28

Difficulty: Challenging

Suppose  $n$  lamps are arranged in a circle, numbered from 1 to  $n$  clockwise, with  $0 \leq L \leq n$  lamps turned on. A testing procedure consists of performing the following operations:

for each lamp turned on, we take its number  $i$ ; the  $i$  lamps following it clockwise receive a signal

- After a signal has been sent for each lamp turned on, every lamp which received an odd number of signals will have its state changed; those receiving an even number of signals will not.

Let  $T$  be the set of all  $2^n$  possible initial configurations. We define a function  $t$  from  $T$  to  $T$ , where if  $X$  represents an initial configuration of lamps,  $t(X)$  is the configuration obtained after applying the testing procedure to  $X$ . Suppose  $S$  is the sum of all integers  $n < 2^{2015}$  such that the function  $t$  is bijective. Find the smallest positive integer  $c$  such that  $2^c$  is greater than the sum of every element of  $S$ .

*Solution.* (Art of Problem Solving write ups)

□

# §11 err¶¶: ¶¶ ¶¶¶ ¶¶¶¶¶¶¶¶ ¶¶¶¶ ¶¶ ¶¶¶ ¶¶¶¶ ¶¶¶¶¶. (Season 11)

## §11.1 Week 1

### §11.1.1 Integral AM-GM

Source: Original

Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)

Problem ID: 147

Date: 2021-03-01

Difficulty: Beginner

$p, q$  are positive integers such that  $pq = 2021^{2021}$ . What is the minimal number of digits in  $p + q$ ?

*Solution.*

By AM-GM the minimal value of  $p + q$  occurs when  $p, q$  are close; an obvious close pair is  $(p, q) = (43^{1010}, 47^{1010})$  and the minimal possible value of  $p + q$  (over positive reals) occurs at  $p = q = 2021^{1010.5}$ . The number of digits in a number  $n$  is  $\lfloor \log_{10}(n) \rfloor + 1$ ; noticing this we see that both our obvious guess and the absolute minimum have 3341 digits. □

**§11.1.2 d20 but not D20***Source: Original**Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)**Problem ID: 148**Date: 2021-03-02**Difficulty: Beginner*

21 people, including IcosahedralDice, each roll an icosahedral die, which have faces numbered 1 through 20 and each face is equally likely to land face-up. Given that IcosahedralDice rolled the (strictly) largest number, find the expected value of his roll.

*Solution.*

Let IcosahedralDice roll  $r$ . The probability that a person rolls between 1 and  $r - 1$  inclusive is  $\frac{r-1}{20}$  so the probability that all 20 other people rolled less than  $r$  is  $\left(\frac{r-1}{20}\right)^{20}$ ; thus the probability IcosahedralDice's roll is strictly greater than everyone else's is  $\sum_{r=1}^{20} \left(\frac{r-1}{20}\right)^{20}$ . The expected value is then

$$E = \frac{\sum_{r=1}^{20} r \cdot \left(\frac{r-1}{20}\right)^{20}}{\sum_{r=1}^{20} \left(\frac{r-1}{20}\right)^{20}} \iff \lfloor 100000E \rfloor = \boxed{1952901}.$$

□



**§11.1.3 Is it better to walk or run in the rain?***Source: Original**Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)**Problem ID: 149**Date: 2021-03-03**Difficulty: Easy*

Rain is falling at an angle of 2.021 (radians) measured anticlockwise from the positive  $x$ -axis, with velocity 2021. A unit sphere rolls from the origin to (2021, 0) with constant speed  $v$ .

Given that  $v$  minimizes the amount of rain that falls on the ball, find  $\lfloor 10^2 v \rfloor$ , where  $\lfloor x \rfloor$  is the greatest integer less than or equal to  $x$ .

*Note that if you run too fast, you end up running into the rain.*

*Solution.*

Change the frame of reference such that the rain is still relative to you; this can be done by moving the entire plane with velocity  $(2021 \cos \theta, 2021 \sin \theta)$  where  $\theta = 2.021$ , the angle at which rain falls. This means that your destination point will be on the line with slope  $\tan \theta$  passing through (2021, 0) and your new velocity vector is  $(v + 2021 \cos \theta, 2021 \sin \theta)$ . To minimize the amount of rain that falls on you, you should run through as little rain as possible, which means your path should be perpendicular to the destination line; this is achieved when your velocity vector has direction  $\tan(\theta - \frac{\pi}{2})$ . Solving, we get

$$\tan\left(\theta - \frac{\pi}{2}\right) = \frac{2021 \sin \theta}{v + 2021 \cos \theta} \iff v = 4644.3870 \iff \boxed{464438}.$$

□

**§11.1.4 Don't over think this***Source: Original**Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)**Problem ID: 150**Date: 2021-03-04**Difficulty: Medium*

Let  $N$  be the number of ways to put the numbers  $2, 4, 8, \dots, 2^{420}$  into a 20-by-21 grid such that the sums of rows are increasing from top to bottom and the sums of columns are increasing from left to right. How many digits does  $N$  have?

*Solution.*

We solve for a general  $m$ -by- $n$  grid. Note that permuting rows leaves the sums of columns intact, and vice versa. For each of the  $(mn)!$  ways to fill the grid, we must divide by  $m!$  to correct for the ordering of the rows and  $n!$  for the columns; as we established before these events are indepent so our answer is  $\frac{(mn)!}{m!n!}$ ; in the case of  $(m, n) = (20, 21)$  our answer is  $\lfloor \log_{10} \left( \frac{420!}{20!21!} \right) \rfloor + 1 = \boxed{883}$ .  $\square$

**§11.1.5 Genfun is Not Fun***Source: Original**Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)**Problem ID: 151**Date: 2021-03-05**Difficulty: Hard*

Suppose  $a_0 = 0$ ,  $a_1 = 1$  and for  $n \geq 1$ ,

$$a_n = n(a_{n-1} + 3a_{n+1}).$$

If  $S = \sum_{n \geq 0} a_n$  then find the 2021<sup>st</sup> to 2025<sup>th</sup> digits of  $S$ . (For instance, the third to fifth digits of  $\pi$  is 415.)

*Solution.*

We let  $f(x) = \sum_{n \geq 0} a_n x^n$  be a power series; our desired sum  $S$  is just  $f(1)$ . To get something of the form  $na_n x^n$  we can differentiate to get  $f'(x) = \sum_{n \geq 0} (n+1)a_{n+1}x^n$  where  $a_k = 0$  if  $k \leq 0$  for convenience. Now, if we can construct a function  $F(x)$  using a linear combination of  $f(x)$  and  $f'(x)$  such that the coefficient of  $x^n$  in  $F(x)$  is  $n(a_{n-1} + 3a_{n+1})$  then we know  $F(x) = f(x)$ . We can shift indices by multiplying or dividing by powers of  $x$  accordingly; we use this to try and turn  $3f'(x) = \sum_{n \geq 0} 3(n+1)a_{n+1}x^n$  into  $\sum_{n \geq 0} 3na_{n+1}x^n$  so we need to subtract some shifted form of  $f(x)$ . Indeed,  $\frac{f(x)}{x} = \sum_{n \geq 0} a_{n+1}x^n$  so  $f'(x) - \frac{f(x)}{x} = \sum_{n \geq 0} na_{n+1}x^n$ ; similarly we have  $x^2 f'(x) + x f(x) = \sum_{n \geq 0} na_{n-1}x^n$ . So, we have

$$F(x) = 3f'(x) - \frac{3f(x)}{x} + x^2 f'(x) + x f(x) = \sum_{n \geq 0} (a_{n-1} + 3a_{n+1}) x^n = \sum_{n \geq 0} a_n x^n = f(x)$$

which is a separable differential equation; chucking this into WolframAlpha we get

$$f(x) = \frac{Cx}{x+3} \exp\left(\frac{\arctan\left(\frac{x}{\sqrt{3}}\right)}{\sqrt{3}}\right)$$

and since  $\sum_{n \geq 0} a_{n+1}x^n = \frac{f(x)}{x}$  is 0 when  $x = 0$  we have  $C = 3$  and

$$f(1) = \frac{3}{4} \exp\left(\frac{\pi\sqrt{3}}{18}\right)$$

and using WolframAlpha our required 5-digit number is 64744. □

## §11.1.6 Harmonic Sum

Source: Original

Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)

Problem ID: 152

Date: 2021-03-06

Difficulty: Challenging

Let  $S$  be the set of positive integers which when written in base  $2021^{2021}$  have no identical consecutive digits. Find the first five digits of

$$\sum_{n \in S} \frac{1}{n}.$$

Solution.

Call an integer *distinctive* if it is in  $S$ . Let  $S(d, n)$  be the set of distinctive positive integers which when written in base  $b = 2021^{2021}$  are  $n$  digits long and have leading digit  $d$  and let  $T(d, n)$  be the sum of the reciprocals of the integers in  $S$ . We will attempt to bound  $T(d, n)$  as follows, since if the leading digit of an integer is  $d$  then the second digit cannot be  $d$ :

$$T(d, n) = \sum_{k \in S(d, n)} \frac{1}{k} = \sum_{k=1, k \neq d}^{b-1} T(d, n-1) = \sum_{k=0, k \neq d}^{b-1} \sum_{h \in S(k, n-1)} \frac{1}{h}$$

and note that  $h$  takes the form of  $d \cdot b^{n-1} + k \cdot b^{n-2} + O(b^{n-3})$ . An upper bound would be to let  $O(b^{n-3}) = 0$  and count  $T(d, n-1)$ ; since there are  $(b-1)^{n-2}$   $n-1$ -digit distinctive numbers with a fixed leading digit this gives the bound

$$T(d, n) < \sum_{k=0}^{b-1} \frac{(b-1)^{n-2}}{d \cdot b^{n-1} + k \cdot b^{n-2}} = \sum_{k=0}^{b-1} b^{2-n} \frac{(b-1)^{n-2}}{db + k} = \frac{(b-1)^{n-2}}{b^{n-2}} (H_{db+b-1} - H_{db});$$

letting  $T(n)$  be the sum of  $T(d, n)$  over all  $d$  we have

$$T(n) < \frac{(b-1)^{n-2}}{b^{n-2}} (H_{b^2} - H_b)$$

or

$$\sum_{n \in S} \frac{1}{n} < \sum_{n \geq 1} \frac{(b-1)^{n-2}}{b^{n-2}} (H_{b^2} - H_b) \approx \sum_{n \geq 1} \left( \frac{b-1}{b} \right)^{n-2} \ln b = \frac{b^2 \ln b}{b-1}$$

whose first five digits are 54518.

By AM-HM inequality, if two distinct positive integers  $a, b$  have average  $m$  then  $\frac{1}{a} + \frac{1}{b} > \frac{2}{m}$ ; noting that  $n-1$ -digit distinctive numbers that start with  $d$  come in pairs where the last  $n-2$  digits of each pair add up to  $b^{n-1} - 1$  (try subtracting any four-digit base-10 distinctive integer from 9999 and see what I mean) we can bound

$$T(d, n) > \sum_{k=0}^{b-1} \frac{2(b-1)^{n-2}}{2d \cdot b^{n-1} + (2k+1)b^{n-1} - 1} - \frac{2(b-1)^{n-2}}{2d \cdot b^{n-1} + d \cdot b^{n-2}}$$

and since  $b$  is so large we take ignore the  $-1$  term and get

$$T(d, n) > \sum_{k=0}^{b-1} \frac{2(b-1)^{n-2}}{b^{n-1}} \cdot \frac{1}{2bd + 2k + 1} - \frac{2(b-1)^{n-2}}{b^{n-2}} \cdot \frac{1}{2bd + 1}.$$

Now summing over  $d$ , the first harmonic sum turns into

$$\frac{1}{2b+1} + \frac{1}{2b+3} + \cdots + \frac{1}{2b^2+1} = \log(2b^2+1) - \log(2b) - \frac{1}{2} [\log(b^2) - \log(b+1)]$$

and the second harmonic sum is

$$\frac{1}{2b+1} + \frac{1}{4b+1} + \cdots + \frac{1}{2b^2-b+1} < \frac{1}{2b} \log(b-1) = O\left(\frac{1}{b}\right)$$

which we can safely ignore (feel free to plug it in if you want; it's just extra terms). This is a bound sharp enough to also give 54518 as its first five digits, and we are done.  $\square$

### §11.1.7 Lonely Tetrahedron

*Source: Original*

*Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)*

*Problem ID: 153*

*Date: 2021-03-07*

*Difficulty: Challenging*

For a point  $P$  on the surface of a regular tetrahedron with side length 1, define the *isolation* of  $P$  to be the smallest  $d$  such that an ant crawling on the surface of the tetrahedron, starting at  $P$ , can reach any other point on the tetrahedron by traveling at most  $d$  units. Find the total length of the locus of points whose isolation is  $\frac{21}{20}$ .

*Solution.*

□

## §11.2 Week 2

## §11.2.1 Don't over think this part 2

---

Source: Original

Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)

Problem ID: 154

Date: 2021-03-08

Difficulty: Beginner

---

Find the number of real solutions to  $\{x^{21}\} = \{x^{20}\}$ , where  $20 \leq x \leq 21$ .

*Solution.*

If the fractional parts of  $x^{20}$  and  $x^{21}$  are equal, then  $x^{21} - x^{20}$  must be an integer. Since  $p(x) = x^{21} - x^{20}$  is continuous and strictly increasing when  $20 \leq x \leq 21$ , it remains to find the number of integers between  $p(21)$  and  $p(20)$  inclusive, which is  $p(21) - p(20) + 1 = \boxed{21^{21} - 21^{20} - 20^{21} + 20^{20} + 1}$ .  $\square$

## §11.2.2 Lattice Points on a Circle

Source: Original

Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)

Problem ID: 155


Date: 2021-03-09

Difficulty: Beginner

A circle with area  $\frac{20}{21}$  is placed randomly in the plane. What is the probability that the circle will contain a lattice point? (Formally, if  $p_M$  is the probability that a circle with center chosen randomly in  $[-M, M]^2$  contains a lattice point, find  $\lim_{M \rightarrow \infty} p_M$ .)

*Solution.*

By symmetry, it suffices to consider the probability when the circle's center is chosen randomly inside a unit square. Draw four quartercircular arcs, one at each vertex; such a circle's center must be contained in the union of those arcs in order for it to contain a lattice point (otherwise a lattice point would more more than  $r$  away from the circle's center). Shown below is the diagram.



Since the arcs overlap, we split the area into four isosceles triangles and four arcs. The base angle of the triangle is  $\arccos\left(\frac{1}{2r}\right)$  and the height of the triangle is  $\sqrt{r^2 - \frac{1}{4}}$  so the angle of the arc is  $\frac{\pi}{2} - 2\arccos\left(\frac{1}{2r}\right)$ . Multiplying by four and plugging in  $r = \sqrt{\frac{20}{21\pi}}$  we get

$$P = 2\sqrt{\frac{20}{21\pi} - \frac{1}{4}} + \frac{20}{21} \left( 1 - \frac{4\arccos\left(\frac{\sqrt{21\pi}}{2\sqrt{20}}\right)}{\pi} \right).$$

□



**§11.2.3 WA is Required***Source: Original**Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)**Problem ID: 156**Date: 2021-03-10**Difficulty: Easy*

Find the number of digits in the smallest positive integer  $x$  such that there exists a positive integer  $y$  such that

$$2020x^{2021} = 2021y^{2120}.$$

*Solution.*

Since  $2020 = 2^2 \cdot 5 \cdot 101$  and  $2021 = 43 \cdot 47$  we split  $x, y$  into their prime factors; for  $x$  to be minimal it should only have the necessary prime factors 2, 5, 43, 47, 101. Let  $x = 2^{x_1} \cdot 5^{x_2} \cdot 43^{x_3} \cdot 47^{x_4} \cdot 101^{x_5}$  and  $y = 2^{y_1} \cdot 5^{y_2} \cdot 43^{y_3} \cdot 47^{y_4} \cdot 101^{y_5}$ ; we now can solve for the exponent of each prime independently. Looking at the first equation we get  $2 + 20^{21}x_1 = 21^{20}y_1$ , or the equivalent  $2 + 20^{21}x_1 \equiv 0 \pmod{21^{20}}$ . Let  $I(a)$  be the modular inverse of  $a \pmod{21^{20}}$ ; then we have  $x_1 \equiv -2I(20^{21}) \pmod{21^{20}}$ . Solving similar equations for  $x_2$  through  $x_5$  we get  $x_2 \equiv -I(20^{21}) \pmod{21^{20}}$ ,  $x_3 = x_4 = I(20^{21}) \pmod{21^{20}}$ . Putting this together, the number of digits is

$$\begin{aligned} & \lfloor [21^{20} - 2I(20^{21})] \log_{10}(2) + [21^{20} - I(20^{21})] \log_{10}(505) + I(20^{21}) \log_{10}(2021) \rfloor + 1 \\ &= \boxed{835858264863808864064926340}. \end{aligned}$$

□

**§11.2.4 Hard***Source: Original**Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)**Problem ID: 157**Date: 2021-03-11**Difficulty: Hard*

Let  $n$  be a power of 10 greater than  $10^{100}$ . A regular  $n$ -gon is inscribed in a unit circle, and a subset  $S$  is chosen uniformly at random from the  $2^n$  subsets of the polygon's vertices. If  $A$  is the expected area of the convex hull of  $S$ , find  $\pi - A$ .

*Solution.*

By linearity of expectation, finding  $\pi - A$  is the same as finding the expected area outside the convex hull inside the circle. So, we find the expected area of a circular segment (the little cap thingy formed when you subtract the triangle off from the circular arc). Label the points 1 through  $n$  counterclockwise. Consider a circular segment that reaches from 1 to  $k$ . The probability of getting such a circular segment from your point selection is  $2^{-k}$ , since the endpoints of the segment (vertices 1 and  $k$ ) need to be included while the  $k - 2$  middle segments must be excluded. The area of such a circular segment is  $\frac{(k-1)\pi}{n} - \frac{1}{2} \sin\left(\frac{2\pi(k-1)}{n}\right)$  (yes, this covers segments with an obtuse angle as well - why?). Summing, we see that  $k$  can range from 2 to  $n$ ; however we need to correct for the edge cases where we select one or zero points, which contributes an expected area of  $\frac{(n+1)\pi}{2^n}$  (the whole circle). Since there are  $n$  of each of the circular segments, our sum is

$$n \sum_{k=2}^n 2^{-k} \left[ \frac{(k-1)\pi}{n} - \frac{1}{2} \sin\left(\frac{2\pi(k-1)}{n}\right) \right] + \frac{(n+1)\pi}{2^n} = \frac{n}{2} \sum_{k=1}^{n-1} 2^{-k} \left[ \frac{k\pi}{n} - \frac{1}{2} \sin\left(\frac{2k\pi}{n}\right) \right] + \frac{(n+1)\pi}{2^n}$$

and since  $n$  is huge we can approximate  $n - 1$  with  $\infty$  and ignore the edge case (since  $2^n$  grows way faster than  $n$ ). The first sum of  $k2^{-k}$  is an arithmo-geometric series and evaluates to 2; the second sum is the imaginary part of the sum of complex numbers

$$\sum_{k=1}^{\infty} \left(\frac{z}{2}\right)^k = \frac{z}{2} \cdot \frac{1}{1 - \frac{z}{2}} = \frac{z}{2 - z}$$

where  $z = \cos(a) + i \sin(a)$  and  $a = \frac{2\pi k}{n}$ . Simplifying further, we have

$$\frac{z}{2 - z} = \frac{\cos(a) + i \sin(a)}{2 - \cos(a) - i \sin(a)} = \frac{2 \cos(a) + 2i \sin(a)}{5 - 4 \cos(a)}$$

and scaling by a factor of 2 we have

$$\sum_{k=1}^{\infty} 2^{-k} \frac{1}{2} \sin\left(\frac{2k\pi}{n}\right) = \frac{\sin(a)}{5 - 4 \cos(a)}$$

so putting it all together the expected value of  $\pi - A$  is

$$\pi - \frac{n \sin\left(\frac{2\pi}{n}\right)}{10 - 8 \cos\left(\frac{2\pi}{n}\right)}.$$

Since  $n$  is huge, we use the first two terms of the Taylor series to approximate our trigonometric functions;

doing this we get  $\boxed{\frac{26\pi^3}{3n^2}}$ . □

*Solution.* [Write-up by Boldizsár#3106 (694967174921715853)]

$n$  is huge. Without loss of generality, one can assume that  $S$  has at least 3 elements. We try to compute the expected area of one slice, i. e. the part of a circular slice outside the polygon. Because of linearity of expectation,  $n$  times this is  $\pi - A$ . Assume that, on the left of this circular sector, the  $k$ -th vertex is the first one in  $S$  and on the right, the  $l$ -th one. So the minimum case is when  $k = l = 0$  when two neighboring vertices are both in  $S$ . The probability of one choice of  $k, l$  happening is  $\frac{1}{2^{k+1}2^{l+1}}$ . The whole circular sector our slice lies in consists of  $k + l + 1$  slices, so it has the area  $\frac{\pi(k+l+1)}{n}$ , the corresponding triangle has area  $\frac{\sin(\frac{2\pi(k+l+1)}{n})}{2}$ . As  $n$  is huge, we can sum to infinity without problems and we are interested in

$$n \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{\frac{\pi(k+l+1)}{n} - \frac{\sin(\frac{2\pi(k+l+1)}{n})}{2}}{2^{k+1}2^{l+1}(k+l+1)}.$$

Now, again,  $n$  is huge so we can use the "fundamental theorem of engineering refined" without any concerns, i. e. that  $\sin(x) = x - \frac{x^3}{6}$ , so we have:

$$\begin{aligned} \frac{n}{24} \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{\left(\frac{2\pi(k+l+1)}{n}\right)^3}{2^{k+l+1}(k+l+1)} &= \frac{\pi^3}{3n^2} \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{(k+l+1)^2}{2^{k+l+1}} = \\ &= \frac{\pi^3}{3n^2} \sum_{m=0}^{\infty} \sum_{k=0}^{m-1} \frac{m^2}{2^m} = \frac{\pi^3}{3n^2} \sum_{m=0}^{\infty} \frac{m^3}{2^m}. \end{aligned}$$

So we ask WolframAlpha and we are surprised that there comes a nice answer, namely 26, a random integer. Where the 26 comes from is left to the reader. Finally, we are left with  $\frac{26\pi^3}{3n^2}$ . As we all know, the PSC consists of nice people, so they set  $n$  and hence  $n^2$  to a power of 10, so we now understand why they asked for the 2021st to 2030th significant figures, namely because this is just the 2021th to 2030th digits of  $\frac{26\pi^3}{3}$ , which is, according to WolframAlpha, 8481609952.  $\square$

**§11.2.5 hard***Source: Original**Proposer: tanoshii#3160 (300065144333926400) and iparis#5018 (269634843468496906)**Problem ID: 158**Date: 2021-03-12**Difficulty: Hard*

Let  $N = 20^{21}$ . You play a game, starting with \$1. At each of  $N$  steps, you can bet any nonnegative amount of money,  $m$ , up to how much you currently have, and a computer flips a coin. If heads, you double your bet; if tails you lose it.

However, before you start, you hack the computer, and you know that it will generate exactly  $\frac{N}{2} + K$  heads.

Suppose that you can guarantee yourself at least \$2021. If  $K$  is minimal, find the leading five digits of  $K$ .

*Solution.*

Let  $g(h, t)$  denote the largest amount of money you can guarantee yourself if you start out with \$1. We claim that if we start out with \$ $s$ , then the largest amount of money you can guarantee yourself is  $sg(h, t)$ . Clearly,  $sg(h, t)$  is attainable, since we can use the same betting procedures when we did for \$1 and scale the bets proportionally by  $s$ . And if we can guarantee more than  $sg(h, t)$  then dividing each of the bets by  $s$  would mean that we can guarantee more than  $g(h, t)$  if we started out with \$1, contradiction. (Intuitively, the amount of money you gain or lose in a bet is homogeneous with the amount of money you already have, so scalings are valid.)

On our first step, suppose our optimal bet is \$ $b$ . If heads, the largest amount of money we can guarantee ourselves is  $(1 + b)g(h - 1, t)$ ; if tails it is  $(1 - b)g(h, t - 1)$  and so

$$g(h, t) = (1 + b)g(h - 1, t) = (1 - b)g(h, t - 1).$$

Noting that the sum of heads and tails decreased by 1, we seek  $g(h, t)$  in terms of  $g(h - 1, t)$  and  $g(h, t - 1)$ . Indeed,

$$(1 + b) + (1 - b) = \frac{g(h, t)}{g(h - 1, t)} + \frac{g(h, t)}{g(h, t - 1)} \iff \frac{2}{g(h, t)} = \frac{1}{g(h - 1, t)} + \frac{1}{g(h, t - 1)}$$

so  $g(h, t)$  is the harmonic mean of  $g(h - 1, t)$  and  $g(h, t - 1)$ . With the values  $g(h, 0) = 2^h$  (go all-in every time) and  $g(0, t) = 1$  (bet nothing every time) we get

$$g(h, t) = \frac{2^{h+t}}{\sum_{k=0}^t \binom{h+t}{k}} \iff \sum_{k=0}^t 2^{-N} \binom{N}{k} \leq \frac{1}{2021}$$

after substituting relevant values. The last sum is a binomial distribution with  $n = N$  and  $p = q = \frac{1}{2}$ ; that is, think of  $n$  independent events that can each take one of two outcomes: pass or fail, where each event passes with probability  $p$  and fails with probability  $q = 1 - p$ . Letting  $X$  be the random variable that denotes the number of events that passed, we see that our expression is equal to  $\Pr(X \leq t) \leq \frac{1}{2021}$ ; since  $n$  is large we can approximate this binomial distribution by the normal distribution; in particular the quantity  $\frac{X}{n}$  is normally distributed with mean  $\mu = np = \frac{n}{2}$  and variance  $\sigma^2 = \frac{pq}{n}$ . So, we want to find a  $z$  such that

$$\Pr\left(\frac{X/n - \mu}{\sigma} \leq z\right) = \frac{1}{2021} \iff \frac{t}{n} = \sigma z + \mu \iff K = \lfloor z \rfloor n \sigma$$

and plugging  $\frac{1}{2021}$  as our  $p$ -value into a  $p$ -value calculator we get that  $K = \boxed{3.2946\sqrt{0.25 \cdot 20^{21}}}$ . □

## §12 Return to Normalcy (Season 12)

### §12.1 Week 1

#### §12.1.1 Quadrilateral Perimeter

---

*Source: Folklore*

*Proposer: flame#6784 (185229437787176962)*

*Problem ID: 161*

*Date: 2021-03-29*

*Difficulty: Beginner*

---

Let  $ABCD$  be a convex quadrilateral satisfying

$$AB = 1, BC = 6, CD = 18, AC \perp BD.$$

Find its perimeter.

*Solution.*

Let  $E$  be the intersection of the diagonals of  $ABCD$ . Then  $AB^2 + CD^2 = AE^2 + EB^2 + CE^2 + ED^2$  and  $BC^2 + DA^2 = BE^2 + EC^2 + DE^2 + EA^2$  by the Pythagorean Theorem. This implies that  $AB^2 + CD^2 = BC^2 + DA^2$ , or  $1^2 + 18^2 = 6^2 + x^2$ . Solving yields  $x = 17$ , so the answer is  $1 + 18 + 6 + 17 = \boxed{42}$ .  $\square$

**§12.1.2 Involution Convolution**

---

Source: Australian Senior Mathematical Challenge, 2019 Q30

Proposer: brainysmurfs#2860 (281300961312374785)

Problem ID: 162

Date: 2021-03-30

Difficulty: Easy

---

A function  $f$ , defined on the set of positive integers, has  $f(1) = 2$  and  $f(2) = 3$ . Also  $f(f(f(n))) = n + 2$  if  $n$  is even and  $f(f(f(n))) = n + 4$  if  $n$  is odd. What is  $f(777)$ ?

*Solution.*

Observe that  $777 = 4 \times 194 + 1$ . Thus

$$\begin{aligned} f(777) &= f(f^{194 \times 3}(1)) \\ &= f^{194 \times 3}(f(1)) \\ &= f^{194 \times 3}(f(2)) \\ &= 2 + 194 \times 2 \\ &= \boxed{390} \end{aligned}$$

□

## §12.1.3 Coordinates without Bash

Source: Original Problem

Proposer: Denial#4301 (118831126239248397)

Problem ID: 163

Date: 2021-03-31

Difficulty: Medium

A circle of radius 1 is centred at  $(1, 1)$ . Points  $A$  and  $B$  have co-ordinates  $(a, 0)$  and  $(0, b)$ , with  $a, b > 2$ . If  $AB$  is tangent to the circle, what is the sum of all possible integer values of  $a$  and  $b$ ?

*Solution.* [Solution by Denial#4301 (118831126239248397)]

Let  $O, C, D$  and  $E$  have coordinates  $(1, 1), (0, 0), (0, 1)$  and  $(1, 0)$  respectively, and let the point of tangency of  $AB$  with the circle be  $F$ .

Note since  $O$  and  $D$  have the same  $y$  co-ordinate  $OD$  is parallel to the  $x$ -axis and hence perpendicular to the  $y$ -axis. Since  $D$  is also on the  $y$ -axis,  $BC$  is tangent to the circle at  $D$ . Similarly,  $O$  and  $E$  have the same  $x$  co-ordinate, so  $OE$  is parallel to the  $y$ -axis and hence perpendicular to the  $x$ -axis. Since  $E$  is also on the  $x$ -axis,  $AC$  is tangent to the circle at  $E$ .

Since the tangents to a circle from a point have equal length,  $BF = BD$  and  $FA = AE$ . Summing,  $BF + FA = BD + AE$ , so  $BA = (BC - CD) + (AC - CE) = (b - 1) + (a - 1) = a + b - 2$ , giving  $AB^2 = (a + b - 2)^2$ . However, since  $\angle ACB = 90^\circ$ , by Pythagoras' theorem,  $AB^2 = a^2 + b^2$ , and so we have

$$a^2 + b^2 = (a + b - 2)^2$$

$$a^2 + b^2 = a^2 + b^2 + 4 + 2ab - 4a - 4b$$

$$0 = 2ab - 4a - 4b + 4$$

$$ab - 2a - 2b + 2 = 0$$

$$ab - 2a - 2b + 4 = 2$$

$$(a - 2)(b - 2) = 2$$

However,  $a, b > 2$ , so if  $a$  and  $b$  are integers, then  $a - 2$  and  $b - 2$  are positive integers and hence must both be positive factors of 2. Since 2 is prime, its only factorisation is as  $2 = 2 \times 1$ , so either  $a - 2 = 2$  and  $b - 2 = 1$  or  $a - 2 = 1$  and  $b - 2 = 2$ . Hence the possible integer values of  $a$  and  $b$  are  $(a, b) = (3, 4)$  or  $(4, 3)$ . Giving a final answer of 14 □



**§12.1.4 April Fool's**

---

*Source: brain*

*Proposer: brainysmurfs#2860 (281300961312374785)*

*Problem ID: 164*

*Date: 2021-04-01*

*Difficulty: Random*

---

How many official solves will this Sunday's problem get?

*Solution.*

Answer: 21 Five people managed to guess 21.



## §12.1.5 FE

Source: 2011 Kosovo TST #5

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 165

Date: 2021-04-02

Difficulty: Hard

There exist functions  $f : \mathbb{R} \setminus \{1, -1\} \rightarrow \mathbb{R}$  such that the following holds:

$$f\left(\frac{x-3}{1+x}\right) + f\left(\frac{x+3}{1-x}\right) = x$$

Then the average value of

$$\sum_{k=2}^{100} \frac{f(k)}{k}$$

can be expressed in the form  $\frac{m}{n}$ , where  $m, n$  are relatively prime integers and  $n$  is positive. Find  $100m + n$ .

(For example, if the solutions to that FE were  $f(x) \equiv 0$  and  $f(x) \equiv -x^2$  then the average value of that sum would be  $-\frac{5049}{2}$  and your answer would be  $2 - 504900 = -504898$ .)

(A four-function calculator may be used.)

*Solution.*

The unique function can be found using [AoPS solutions](#); then note that  $\frac{f(x)}{x}$  can be decomposed into partial fractions and so the sum telescopes. Answer: -13243675 □

**§12.1.6** So this is where American problem quality went

---

Source: 2012 AIME II #15

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 166

Date: 2021-04-03

Difficulty: Challenging

---

Triangle  $PQR$  is inscribed in circle  $\Gamma$  with  $PQ = 5$ ,  $QR = 7$ , and  $RP = 3$ . The bisector of angle  $P$  meets side  $\overline{QR}$  at  $X$  and circle  $\Gamma$  at a second point  $Y$ . Let  $\gamma$  be the circle with diameter  $\overline{XY}$ . Circles  $\Gamma$  and  $\gamma$  meet at  $Y$  and a second point  $Z$ . Then  $PZ^2 = \frac{m}{n}$ , where  $m$  and  $n$  are relatively prime positive integers. Find  $100m + n$ .

*Solution.*

**AoPS Solutions** Answer: 90019

□

## §12.1.7 640 Generators of Pain

Source: 2015 PUMAC Number Theory A7

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 167

Date: 2021-04-04

Difficulty: Challenging

For positive integers  $n$ , let  $s(n)$  be the sum of the  $n$ -th powers of the primitive roots mod 1601. Find the number of positive integers  $n \leq 2021$  such that  $s(n)$  is divisible by 1601.

*Solution.*

First, some lemmas about orders and generators (skip if you think people should know this already):

**Lemma 1.** The order of  $g^n$ , where  $g$  is a generator mod prime  $p$ , is  $d = \frac{p-1}{(n, p-1)}$  where  $(a, b)$  is the greatest common divisor function.

**Proof.** Since  $g^{nd} \equiv 1 \pmod{p}$  we have  $p-1 \mid nd$  so  $\frac{p-1}{(n, p-1)} \mid d$ . Since  $d$  is defined to be minimal we have  $d = \frac{p-1}{(n, p-1)}$ .

**Lemma 2.** The number of residues mod  $p$  with order  $d$  is  $\varphi(d)$ , where  $\varphi$  is Euler's totient function.

**Proof.** We want to find the number of positive integers  $k$  such that  $d$  is the least positive integer for which  $\left(g^{k \frac{p-1}{d}}\right)^d \equiv 1 \pmod{p}$ . For that to be satisfied,  $k$  and  $d$  cannot share any common factors; otherwise  $\left(g^{k \frac{p-1}{d}}\right)^{\frac{d}{(k, d)}} \equiv 1 \pmod{p}$ . Also, since  $d$  is the order only  $k < d$  generate unique residues. The number of positive integers  $k$  less than  $d$  such that  $(d, k) = 1$  is  $\varphi(d)$ .

Let  $p = 1601$  since 1601 is prime. The order of  $g^n$  mod  $p$  is  $d = \frac{p-1}{(n, p-1)}$ ; also each of the  $\varphi(p-1)$  terms in  $s(n)$  can take only one of  $\varphi\left(\frac{p-1}{(n, p-1)}\right)$  residues with order  $d$  mod  $p$ . By symmetry, each residue of order  $d$  is represented the same amount of times in  $s(n)$ . Now let  $r(d)$  be the sum of the residues with order  $d$  mod  $p$ , so

$$s(n) = \frac{\varphi(p-1)}{\varphi\left(\frac{p-1}{(n, p-1)}\right)} r(d).$$

Notice that the ratio of totient functions is less than  $p$ , so if  $p \mid s(n)$  then  $p \mid r(d)$ .

Consider a residue  $a$  with order  $d$ ; then  $p \mid a^d - 1 = (a-1)(a^{d-1} + a^{d-2} + \dots + 1)$  and since  $a < p$  we have  $a^{d-1} + a^{d-2} + \dots + 1 \equiv 0 \pmod{p}$ . Remembering  $\sum_{d \mid n} \varphi(d) = n$  we conjecture that that sum is  $\sum_{e \mid d} r(e) \equiv 0 \pmod{p}$ ; indeed we can treat  $a$  like the "generator" of the residue class of order  $d$  mod  $p$  and follow the steps of Lemma 2.

**Claim.**  $r(d) \equiv 0 \pmod{p}$  if and only if  $d$  is not squarefree.

As with any divisor sum, we try products of primes:  $r(1) = 1$ ,  $r(p) = -1$ ,  $r(pq) + r(p) + r(q) + 1 \equiv 0 \iff r(pq) = 1$  and induct on  $k$  to get  $r(p_1 p_2 \dots p_k) = (-1)^k$ . Notice  $r(p^2) + r(p) + r(1) \equiv 0 \iff r(p^2) \equiv 0$ ; a similar induction argument yields  $r(d) \equiv 0$  if  $d$  is not squarefree.

We are homeward-bound: for  $d = \frac{p-1}{(n, p-1)} = \frac{1600}{(n, 1600)}$  to be squarefree we must have  $2^5 \cdot 5 = 160 \mid n$ . We have  $2021 - \lfloor \frac{2021}{160} \rfloor = \boxed{2009}$  such  $n$ . □

*Solution.* [Solution by Boldizsár#3106 (694967174921715853)]

$n$ -th powers of roots is something that polynomials can handle very well. The definition of order,  $g^{p-1} \equiv 1 \pmod{p} \iff g^{p-1} - 1 \equiv 0 \pmod{p}$ , motivates us to look at the polynomial  $x^{p-1} - 1$ . Which integers  $g$  satisfy  $g^{p-1} - 1 \equiv 0 \pmod{p}$  but not  $g^d - 1 \equiv 0 \pmod{p}$  for all  $d < p$ ? We claim that the polynomial in question is  $\Phi_{p-1}(x)$ , the  $p-1$ -th cyclotomic polynomial. (If you don't know what a cyclotomic polynomial is, google.) While it is easy to compute  $\Phi_p(x)$ , computing  $\Phi_{p-1}(x) = \Phi_{1600}(x)$  will be harder. Using  $\Phi_n(x) = \prod_{d|n} \Phi_d(x)$  we can get

$$x^{p^k} - 1 = (x^{p^{k-1}} - 1) \Phi_{p^k}(x) = \left[ (x^{p^{k-1}})^p - 1 \right] \iff \Phi_{p^k}(x) = \sum_{h=0}^{p-1} x^{hp^{k-1}}.$$

We now turn our attention to  $\Phi_{p^k n}(x)$  for  $(p, n) = 1$ . Note that  $\Phi_n(x^{p^k})$  has roots of unity  $\exp\left(\frac{\tau m}{p^k n}\right)$  where  $(m, p^k n) = 1, p, \dots, p^k$ . However, we want only  $(m, p^k n) = 1$ ; note then, that  $\Phi_n(x^{p^{k-1}})$  has roots of unity  $\exp\left(\frac{\tau m}{p^{k-1} n}\right) = \exp\left(\frac{\tau pm}{p^k n}\right)$  where  $(m, p^{k-1} n) = 1, p, \dots, p^{k-1}$  so  $(pm, p^k n) = p, p^2, \dots, p^k$  and so

$$\Phi_{p^k n}(x) = \frac{\Phi_n(x^{p^k})}{\Phi_n(x^{p^{k-1}})}.$$

Thus, we have  $\Phi_{25}(x) = 1 + x^5 + x^{10} + x^{15} + x^{20} = \frac{x^{25}-1}{x^5-1}$  and so

$$\Phi_{1600}(x) = \Phi_{2^6 \cdot 25}(x) = \frac{\Phi_{25}(x^{64})}{\Phi_{25}(x^{32})} = \frac{x^{800} + 1}{x^{160} + 1} = 1 - x^{160} + x^{320} - x^{480} + x^{640}$$

where we have omitted two difference of squares factorizations since  $2^6 = 2^5 \cdot 2$  and so  $x^{2^6} = (x^{2^5})^2$ .

*Newton's Sums.* Let  $r_1, r_2, \dots, r_n$  be the roots of polynomial  $a_n x^n + a_{n-1} x^{n-1} + \dots + a_0 x^0$ , and let  $s_k = \sum_{i=1}^n r_i^k$ . Then  $\sum_{i=0}^n a_{n-i} s_{k-i} = 0$  if  $k \geq n$  and  $(-1)^k k a_{n-k} + \sum_{i=0}^{k-1} (-1)^i a_{n-i} s_{k-i} = 0$  if  $k < n$ .

Using Newton's sums, it becomes clear that the only nonzero terms of  $s(n)$  happen when  $160|n$ ; our answer is thus 2009. □

*Solution.* [Solution by Hopeless#9949 (399043449443057704)]

Let  $g$  be a known generator mod  $p$ . Note that if  $(k, p-1) = 1$  then  $g^k$  is also a generator mod  $p$ . We have  $p-1 = 1600 = 2^6 \cdot 5^2$ ; the integers not coprime to  $p-1$  are either multiples of 2, multiples of 5, or multiples of 10. Now observe that

$$\sum_{a=1}^{1600} g^a$$

sums over all powers of  $g$ ; however since we only want coprime powers we need to subtract off powers that are multiples of 2 or 5. Using inclusion-exclusion, this is

$$s(n) = \sum_{a=1}^{1600} g^{an} - \sum_{b=1}^{800} g^{bn} - \sum_{c=1}^{320} g^{cn} + \sum_{d=1}^{160} g^{dn}$$

which can be rewritten and re-summed as

$$\frac{g^{1601n} - 1}{g^n - 1} - \frac{g^{1602n} - 1}{g^{2n} - 1} - \frac{g^{1605n} - 1}{g^{5n} - 1} + \frac{g^{1610n} - 1}{g^{10n} - 1} \equiv 0 \pmod{p}$$

with a small catch: if any of  $g^n, g^{2n}, g^{5n}, g^{10n}$  is 1 the geometric series sum representation fails, and it is easy to see that  $s(n)$  will not be 0 mod  $p$  in that case. Thus, the cases where  $s(n) \not\equiv 0 \pmod{p}$  is when  $160|n$ ; our answer is then 2009. □

## §12.2 Week 2

### §12.2.1 Even Numbers with Restrictions

---

*Source: AMC10, 2011 Q13*

*Proposer: sjbs#9839 (434767660182405131)*

*Problem ID: 168*

*Date: 2021-04-05*

---

How many even integers are there between 200 and 700 whose digits are all different and come from the set  $\{1, 2, 5, 7, 8, 9\}$

*Solution.*

**AoPS Solutions** Answer:

□

**§12.2.2 Enjoyable**

---

Source: Canadian Open Mathematics Challenge 2018

Proposer: TaesPadhiary#8557 (665057968194060291)

Problem ID: 169

Date: 2021-04-06

Difficulty: Beginner

---

Five *distinct* integers  $e, n, j, o, y$  are such that

$$e \cdot n \cdot j \cdot o \cdot y = 775.$$

Find  $e + n + j + o + y$ .

*Solution.*

Note that  $775 = 5^2 \cdot 31$ . In particular, this means that the only way to factor into 5 distinct integers is to have  $\{e, n, j, o, y\} = \{1, -1, 5, -5, 31\}$ , giving us a sum of  $1 - 1 + 5 - 5 + 31 = \boxed{31}$ .  $\square$



## §12.2.3 Isosceles Tetrahedron

Source: IMTS Year 1991 R4 Problem 5 (modified)

Proposer: flame#6784 (185229437787176962)

Problem ID: 170

Date: 2021-04-07

Difficulty: Easy

Paper  $\triangle XYZ$  has  $XY = 69$ ,  $YZ = 91$ ,  $ZX = 100$ . Let  $A, B, C$  be the midpoints of  $\overline{XY}$ ,  $\overline{YZ}$ ,  $\overline{ZX}$  respectively. The triangle is folded about  $AB$ ,  $BC$ , and  $CA$  to form a tetrahedron  $\mathcal{T}$ . What is the volume of  $\mathcal{T}$ ?

*Solution.*

### AoPS Solution using Pythagorean Theorem

We will prove that in an isosceles tetrahedron (opposite edges have the same length) with edge lengths  $a, b, c$ , the volume of such a tetrahedron is

$$V = \sqrt{\frac{(a^2 + b^2 - c^2)(a^2 - b^2 + c^2)(-a^2 + b^2 + c^2)}{72}}.$$

First, note that every "acute" tetrahedron can be inscribed in a parallelepiped prism, with each edge being a diagonal of each face (see [here](#) for a picture). This is because there is exactly one pair of parallel planes that pass through each pair of opposite edges (since opposite edges are skew lines, so you can't just rotate the planes wrt the axis of one edge). Also note that since opposite edges are skew, this means that the opposite edges take up the two different diagonals of their inscribed parallelogram face. Since opposite edges are equal in an isosceles tetrahedron, this implies that the parallelogram has equal diagonals, implying that it's a rectangle, and the circumscribed parallelepiped is actually a rectangular prism. Letting  $x, y, z$  represent the edges of the prism, we have

$$x^2 + y^2 = a^2, y^2 + z^2 = b^2, z^2 + x^2 = c^2 \iff (x^2, y^2, z^2) = \left( \frac{-a^2 + b^2 + c^2}{2}, \frac{a^2 - b^2 + c^2}{2}, \frac{a^2 + b^2 - c^2}{2} \right)$$

and now we claim that the volume of the prism is three times the volume of the tetrahedron; indeed the space outside the inscribed tetrahedron but inside the prism is composed of three congruent right-angle-cornered tetrahedra, each with volume  $\frac{xyz}{6}$ . Thus

$$V = \frac{xyz}{3} = \sqrt{\frac{(a^2 + b^2 - c^2)(a^2 - b^2 + c^2)(-a^2 + b^2 + c^2)}{72}}.$$

Answer: 7605

□

**§12.2.4 Sums of powers of 2 but weird**

---

Source: 2018 PUMAC Algebra A5

Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 171

Date: 2021-04-08

Difficulty: Medium

---

For  $k \in \{0, 1, \dots, 9\}$ , let  $\varepsilon_k \in \{-1, 1\}$ . Find the minimum possible value of

$$\sum_{i=1}^9 \sum_{j=0}^{i-1} \varepsilon_i \varepsilon_j 2^{i+j}.$$

*Solution.*

**Solution PDF** Answer: 174762

□

## §12.2.5 "D4 I swear"

Source: 2012 Winter OMO 43

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 172

Date: 2021-04-09

Difficulty: Hard

Let  $p = 2017$ . An integer  $n$  is uniformly and randomly selected between 1 and  $(p^p - 1)!$  inclusive. The probability that  $n^n - 1$  is divisible by  $p$  can be written in the form  $\frac{m}{n}$ , where  $m$  and  $n$  are relatively prime positive integers. Find  $100m + n$ .

*Solution.*

Fix an order  $d$ ; we find the probability that  $d|n$  and  $n$  has order  $d \bmod p$ . Since the first probability involves  $n$ 's behavior mod  $d$  and the second  $n$ 's behavior mod  $p$  and  $\gcd(d, p-1) = 1$ , these events are independent. Since our interval is from 1 to  $(p^p - 1)!$ , each residue mod  $p$  and mod  $p-1$  (and thus mod  $d$ ) appears the same amount of times. Thus, our first event happens with probability  $\frac{1}{d}$  and our second with probability  $\frac{\varphi(d)}{p}$ , where we have used the fact that there are exactly  $\varphi(d)$  residues with order  $d \bmod p$ . It remains to sum this probability over all possible values of  $d$ , or,

$$P = \frac{1}{p} \sum_{d|p-1} \frac{\varphi(d)}{d} = \frac{1}{p} s(p-1),$$

where  $s(n) = \sum_{d|n} \frac{\varphi(d)}{d}$ . As with all divisor sums, we test with  $n = p, p^2, pq$ :  $s(p) = 1 + \frac{p-1}{p} = 2 - \frac{1}{p}$ ,  $s(p^2) = s(p) + \frac{p(p-1)}{p^2} = 3 - \frac{2}{p}$ ,  $s(pq) = 1 + \frac{p-1}{p} + \frac{q-1}{q} + \frac{(p-1)(q-1)}{pq} = \left(2 - \frac{1}{p}\right) \left(2 - \frac{1}{q}\right)$ ; we conjecture that  $s(n)$  is multiplicative. Indeed, letting  $n = p_1^{e_1} p_2^{e_2} \cdots p_k^{e_k}$ , two induction arguments, one on  $k$  and one on  $e$ , yields

$$s(n) = \prod_{i=1}^k e_i + 1 - \frac{e_i}{p_i}$$

$$\text{so } s(p-1) = s(2016) = \left(6 - \frac{5}{2}\right) \left(3 - \frac{2}{3}\right) \left(2 - \frac{1}{7}\right) = \frac{91}{6} \text{ so } P = \frac{91}{6p} \iff \boxed{21202}.$$

□

### §12.2.6 "change the point distribution up so weekends don't carry people"

Source: Unknown

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 173

Date: 2021-04-10

Difficulty: Challenging

2021 coplanar lines intersect at  $2021 \cdot 1010$  distinct points  $K_1, K_2, \dots, K_{2021 \cdot 1010}$ . Let  $P$  be a point in the plane not on any of the lines. We call an intersection point  $K$  *k-reachable* if segment  $PK$  intersects at most  $k$  lines, excluding the intersection point at  $K$ . If 1999420 points are *k-reachable*, find the largest possible value of  $k$ .

*Solution.*

First note that no two of these 2021 lines are parallel and no three of them are concurrent, otherwise the number of intersection points would be less than  $2021 \cdot 1010$ . After some playing around with small cases of  $n$ , we claim that the minimal number of *k-reachable* points among  $n$  such lines is  $\binom{k+2}{2}$ . The base case  $k = 0$  for arbitrary  $n$  is trivial. Assume the condition holds for  $k - 1$  and  $n - 1$ . Consider the line with minimal distance to  $P$ . Label its intersection points  $P_1, P_2, \dots, P_n$  where  $KP_1 < KP_2 < \dots < KP_n$ . Among these  $n$  points, at least  $k$  of them are *k-reachable* (consider the worst-case scenario where  $P$  is on the left of all the intersection points). Now remove this line; among the remaining points there are at least  $\binom{k+1}{2}$   $(k - 1)$ -reachable points. When we add the line back, those  $(k - 1)$ -reachable points become *k-reachable* points (any point  $Q$  in that set either had  $PQ$  cross that deleted line or not). It remains to bound  $\binom{k+2}{2} \approx (k + 2)^2/2$ . Noticing  $1999420 \approx 1999200 = 1400 \cdot 1428 \approx 1414^2$  and  $\sqrt{2} = 1.414$ , we have  $2 \cdot 1414^2 \approx 2000^2 \iff 1999420 > \binom{2000}{2}$ . Our answer is 1998.  $\square$

## §12.2.7 Notice me, sqing-senpai

Source: 2019 China Mathematical Olympiad P1, Modified

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 174

Date: 202-04-11

Difficulty: Challenging

Let  $(a_1, a_2, a_3, a_4, a_5, a_6)$  be reals no less than  $-2020$  that sum to  $2021$ . Let  $M$  and  $m$  be the maximum and minimum, respectively, of the function

$$f(a_1, a_2, a_3, a_4, a_5, a_6) = \prod_{k=1}^6 (a_k + a_{k+1}),$$

where indices are taken modulo 6, and let  $N$  and  $n$  be the number of ordered pairs which maximize and minimize  $f$ , respectively. Find  $\lfloor \ln(-Mm) \rfloor + N + n$ .

(A scientific calculator may be used.)

*Solution.*

We switch variables to  $(a, b, c, d, e, f)$  for sake of legibility. WLOG  $a$  is maximal; therefore  $a + b, f + a \geq 0$ . We solve for the minimum first, as that is less painful. Negative values are possible, which correspond to either one or three negative factors. If there is only one negative factor, then there are at least two positive variables less than  $2021$ ; flipping their signs and increasing the value of the greatest two variables accordingly produces a product greater in absolute value, so we want three negative factors. Let  $a + b, b + c, c + a < 0$  (the factor adjacency really does matter as we'll see in a bit). Applying AM-GM twice, once to the positive factors and once to the negative, we have

$$f \leq \left( \frac{-a - 2b - 2c - d}{3} \right)^3 \left( \frac{d + 2e + 2f + a}{3} \right)^3$$

and since  $d + 2e + 2f + a = 2 \cdot 2021 - (a + 2b + 2c + d)$  we see that the maximum occurs when  $a + 2b + 2c + d$  is minimized; this happens at  $a = b = c = d = -2020$  and  $e = f = \frac{10101}{2}$ . Note that the variables with coefficient 2 must be different variables; in other words, the three factors we chose to be negative matters. In essence, the two positive terms have to be next to each other, so  $n = 6$ .

Now for the maximum. By the same argument, having all positive factors does not yield a maximal configuration; therefore there are either two or four negative factors. Applying AM-GM in the same fashion (once to the positive factors and once to the negative factors) we get that the equality case is one of four and that the maximal configuration happens when all the variables in the negative terms are equal to  $-2020$ . Thus, four negative factors is optimal; this happens at  $a = b = c = d = e = -2020$  and  $f = 12121$  and  $N = 6$ .

Our answer is

$$\lfloor \ln(2^5 \cdot 2020^6 \cdot 6061^2 \cdot 10101^3) \rfloor + 6 + 6 = \boxed{113}.$$

□

## §13 Quality Uncontrol (Season 13)

### §13.1 Week 1

#### §13.1.1 Choi's Balls

---

*Source: Folklore*

*Proposer: sjbs#9839 (434767660182405131)*

*Problem ID: 175*

*Date: 2021-04-19*

*Difficulty: Beginner*

---

Choi has three types of counters, coloured red, blue and green which she must place on a numberline consisting of the numbers 1,2 and 3.

In how many ways can she arrange the counters such that no red counter is directly to the left of a blue counter?

*Solution.*

There are  $3^3$  ways of arranging the counters without restrictions and  $2 \cdot 3 = 6$  ways to pick the red counter being next to the blue counter, thus we have  $3^3 - 6 = \boxed{21}$  total ways.

□

**§13.1.2 Biscriminant**

---

*Source: Folklore*

*Proposer: sjbs#9839 (434767660182405131)*

*Problem ID: 176*

*Date: 2021-04-20*

*Difficulty: Beginner*

---

Given that for  $f(x) = x^2 + bx + 7$  and  $f(x) \neq -18$ , the possible values of  $b$  can be written in the form  $x < b < y$ . What is the value of  $x^2 + y^2$ ?

*Solution.*

We want to find the values of  $b$  such that  $x^2 + bx + 7 = -18$  has no real solutions. By taking the discriminant, we see that we require  $b^2 - 4 \cdot 25 < 0 \iff -10 < b < 10$ . Therefore our answer is 200.

□

**§13.1.3 IMOK Maclaurin2012 P6**

---

Source: 2012 IMOK Maclaurin Problem 6

Proposer: alchemyst\_ttk#7268 (531054379852365824)

Problem ID: 177

Date: 2021-04-21

Difficulty: Easy

---

Three different positive integers have the property that each of them divides the sum of the other two. Let  $S$  be a possible set of three such integers. What is the smallest possible integer in  $S$  such that the sum of the three integers in  $S$  is greater than 2021?

*Solution.*

Let the three integers be  $a, b, c$  where  $0 < a < b < c$ . This means that  $a + b < 2c$  and because  $c|a + b \rightarrow a + b = c$ . We also know that  $b|a + c \rightarrow b|2a + b \rightarrow b|2a$ .  $b > a$  meaning that  $b = 2a$  and this gives us the solutions  $(a, 2a, 3a)$ . This gives that the sum of the integers in  $S$  is  $6a$  and the smallest multiple of 6 greater than 2021 is 2022. This gives that  $a$  is  $\frac{2022}{6} = \boxed{337}$  which is the answer.  $\square$



§13.1.4 NICE Geometry

Source: 2021 NICE Problem 10


Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 178

Date: 2021-04-22

Difficulty: Medium

A circular sector of angle  $\theta < 180^\circ$  is inscribed inside another sector of the same angle, shown below. If  $\cos(\theta) = -\frac{m}{n}$ , where  $m, n$  are relatively prime positive integers, find  $100m + n$ .




*Solution.*

Let  $r < R$  be the two radii of the sectors. Notice that the two red triangles shown below are both right triangles (one's a tangent, the other's the symmetry axis of an isosceles triangle) and one of their acute angles is  $\frac{\theta}{2}$  in both triangles; since they share the same hypotenuse the two are congruent. Then by Pythagorean Theorem we have

$$r^2 + \left(\frac{r}{2}\right)^2 = R^2 = \frac{5r^2}{4} \iff \frac{r}{R} = \frac{2}{\sqrt{5}} = \sin\left(\frac{\theta}{2}\right)$$

so by double angle formula we have

$$\cos(\theta) = 1 - 2\sin^2\left(\frac{\theta}{2}\right) = -\frac{3}{5} \iff \boxed{305}.$$



□

### §13.1.5 Graph Theory in Disguise

Source: CMC Mock ARML 2019 P10

Proposer: TaesPadhihary#8557 (665057968194060291)

Problem ID: 179

Date: 2021-04-23

Difficulty: Hard

Tony and Wang play a game in which Tony chooses a list of 6 real numbers and Wang devises 6 questions, each of the form “What is the sum of the  $x^{\text{th}}$  and  $y^{\text{th}}$  number in your list?” where  $1 \leq x < y \leq 6$ . Without regard to order, compute the number of ways Wang can choose his questions so that he can determine Tony’s numbers.

*Solution.* [Solution by TaesPadhihary#8557 (665057968194060291)]

We can consider the the game as a graph with 6 vertices, representing Tony’s numbers, and 6 edges, representing Wang’s questions. Note that we can’t have a cycle with even length since then the corresponding system of equations wouldn’t be linearly independent. For example, the system of equations  $x + y = c_1$ ,  $y + z = c_2$ ,  $z + w = c_3$ ,  $w + x = c_4$  is not linearly independent, since the sum of the 1<sup>st</sup> and 3<sup>rd</sup> equations is equal to the sum of the 2<sup>nd</sup> and 4<sup>th</sup>; hence one equation is wasted, yet 6 independent equations are required to find 6 variables. Since the largest tree in a graph with 6 vertices consists of 5 edges, a cycle must exist. We distinguish three cases, depending on the length of the largest cycle:

1. We have a 5-cycle. There are 6 ways to choose which number is left out of the 5-cycle, 12 ways to order the elements of the 5-cycle, and 5 ways to connect the number left out to the cycle, for a total of 360 possibilities.
2. There are two 3-cycles. As the cycles are disjoint, there are  $\frac{\binom{6}{3}}{2} = 10$  such possibilities.
3. There is only one 3-cycle. We choose the cycle  $C$  in  $\binom{6}{3} = 20$  ways, and then choose the paths to connect the other three vertices  $(X, Y, Z)$  to  $C$ , where each of these points must have exactly 1 path. If  $X \rightarrow C$ ,  $Y \rightarrow C$  and  $Z \rightarrow C$ , there are  $3^3 = 27$  ways; if  $X \rightarrow Y \rightarrow C$ , and  $Z \rightarrow C$  (and permutations), there are  $6 \times 3^2 = 54$  ways; if  $X \rightarrow Z$ ,  $Y \rightarrow Z$  and  $Z \rightarrow C$  (and permutations), there are  $3 \cdot 3 = 9$  ways; finally, if  $X \rightarrow Y \rightarrow Z \rightarrow C$  (and permutations), there are  $6 \rightarrow 3 = 18$  ways, for a total of  $20 \rightarrow (27 + 54 + 9 + 18) = 2160$  possibilities. Thus, Wang has a total of  $360 + 10 + 2160 = \boxed{2530}$ .

□

### §13.1.6 What the H-E-double hockey sticks?

Source: old italian contest

Proposer: Matteddy#0482 (329956567132930048)

Problem ID: 180

Date: 2021-04-24

Difficulty: Challenging

There are 2021 people working in a 2021-floor building, one on each floor. On day zero, each worker writes a zero on their sheet of paper. On day  $k > 0$ , the worker on floor  $f > 1$  writes on their sheet of paper the sum of the numbers of the worker on floor  $j - 1$  had written in the days up to  $k - 1$ . The worker on floor 1, meanwhile, writes  $k^2$ . If  $N$  is the number the worker on floor 2021 writes on day 10000, find the number of digits in  $\nu_2(N) + \nu_3(N) + \nu_5(N)$ .

(A scientific calculator may be used.)

*Solution.*

We denote worker  $f$  as the worker who works on floor  $f$ . It's clear that on day  $k$ , worker 2 writes

$$1^2 + 2^2 + \dots + (k-1)^2 = \frac{k(k-1)(2k-1)}{6}$$

using the well-known sum of squares formula, but it's hard to see where it goes from here. Instead, write  $k^2$  as a sum of binomial coefficients. We know that  $2\binom{k}{2} = k(k-1) = k^2 - k$  and  $\binom{k}{1} = k$ , so we have

$$k^2 = 2\binom{k}{2} + \binom{k}{1} \iff 1^2 + 2^2 + \dots + (k-1)^2 = 2 \left[ \binom{2}{2} + \binom{3}{2} + \dots + \binom{k-1}{2} \right] + \left[ \binom{1}{1} + \binom{2}{1} + \dots + \binom{k-1}{1} \right]$$

which sums to  $2\binom{k}{3} + \binom{k}{2}$  after applying the hockey stick identity. Use the hockey stick identity again and again to show that on day  $k$ , worker  $f$  writes down the number

$$2\binom{k}{f+1} + \binom{k}{f} = \binom{k+1}{f+1} + \binom{k}{f+1} = \binom{k}{f+1} \cdot \frac{2k-f+1}{k-f}.$$

Substituting  $f = 2021, k = 10000$  (and doing either Legendre or Krummer's) yields 25. □

**§13.1.7 "What's an mgf?"**

---

Source: 2013 HMMT C9

Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 181

Date: 2021-04-25

Difficulty: Challenging

---

Given a permutation  $\pi$  of  $\{1, 2, \dots, 2021\}$ , let  $p(\pi)$  be the number of fixed points of  $\pi$ . If  $P$  is the set of all permutations  $\pi$ , then let

$$N = \sum_{\pi \in P} p(\pi)^4.$$

Find  $\nu_2(N) + \nu_3(N) + \nu_5(N)$ .

(A scientific calculator may be used.)

Solution.

Contest Solutions Answer:

□

## §13.2 Week 2

### §13.2.1 It's Okay :)

---

*Source: Adapted from a FTW problem.*

*Proposer: ==#5400 (658718130326994954)*

*Problem ID: 182*

*Date: 2021-04-26*

*Difficulty: Beginner*

---

A two-digit positive integer,  $n$ , is doubled and then added to 2 to obtain  $N$ . Then, the digits of  $n$  are reversed to form a new positive integer  $m$ . Given that  $N = m$ , find  $n$ .

*Solution.*

(Sorry, I don't know how to use LaTeX.)

Answer: 25

It is not too difficult to find the answer through educated guessing and checking, but below is a solution that does not require guessing.

Let  $x$  be the tens digit of  $n$  and let  $y$  be the ones digit of  $n$ .

Then the problem can be translated into:  $2(10x + y) + 2 = 10y + x$

Simplifying the equation, we get:  $19x = 8y - 2$

We can easily deduce that:  $19x = 8y - 2 = 38$

Therefore:  $x = 2$  and  $y = 5$

Finally:  $n = 10x + y = 25$

□

**§13.2.2 Points on a Line**

---

*Source: Online Mathematics Open Spring 2020 Problem 1*

*Proposer: TaesPadhiary#8557 (665057968194060291)*

*Problem ID: 183*

*Date: 2021-04-27*

*Difficulty: Beginner*

---

Let  $\ell$  be a line and let points  $A, B, C$  lie on  $\ell$  so that  $AB = 7$  and  $BC = 5$ . Let  $m$  be the line through  $A$  perpendicular to  $\ell$ . Let  $P$  lie on  $m$ . Compute the smallest possible value of  $PB + PC$ .

*Solution.*

Clearly, the minimum occurs when  $A = P$ . If  $C$  is between  $A$  and  $B$  then the distance  $PB + PC$  is lower than if  $C$  is not between  $A$  and  $B$ , giving us an answer of 9. □

### §13.2.3 Diamond Operator

---

Source: PUMaC 2017 A1

Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 184

Date: 2021-04-28

Difficulty: Easy

---

Let  $a \diamond b = ab - 4(a + b) + 20$ . Evaluate

$$1 \diamond (2 \diamond (3 \diamond (\cdots (2020 \diamond 2021) \cdots)))$$

Solution.

Official Solution Answer:



**§13.2.4 Weird Wilson**

---

*Source: Baltic Way [Unknown]*

*Proposer: epicxtroll#6007 (300008472978653184)*

*Problem ID: 185*

*Date: 2021-04-29*

*Difficulty: Medium*

---

Find the smallest prime factor of  $712! + 1$ .

*Solution.*

Unfortunately, 713 is not prime; neither is 715 nor 717. 719 is prime, so by Wilson's Theorem we know that  $718! \equiv 1 \pmod{719}$ . Now we rewrite  $718!$  as follows:

$$718! = 718 \cdot 717 \cdot 716 \cdot 715 \cdot 714 \cdot 713 \cdot 712! \equiv (-1)(-2)(-3)(-4)(-5)(-6) \cdot 712! \equiv 712! \equiv 1 \pmod{719}$$

so  $\boxed{719} \mid 712! + 1$ . □



**§13.2.5 Get the CRUX of this inequality!**

---

Source: *CRUX Mathematicorum*

Proposer: *TaesPadhary#8557 (665057968194060291)*

Problem ID: 186

Date: 2021-04-30

Difficulty: Hard

---

Six positive real numbers  $r_1, r_2, r_3, r_4, r_5, r_6$  are such that

$$3 \cdot \left( \sum_{i=1}^6 (-1)^i r_i \right) = \sum_{i=1}^6 r_i^2 = 18.$$

If  $M$  is the least possible constant such that  $r_1 r_2 r_3 r_4 r_5 r_6 \leq M$ , find the value of  $\lfloor 100M \rfloor$ .

*Solution.*

Source: **CRUX Mathematicorum (go to Page 44-45)** Answer: 100

□

## §13.2.6 Don't overthink this

---

Source: Original (?)

Proposer: ChristopherPi#8528 (696497464621924394)

Problem ID: 187

Date: 2021-05-01

Difficulty: Challenging

---

Points  $A, B, C, D$  are located at  $(-3, 0), (1, 0), (6, 0), (7, 0)$  in the coordinate plane. If the locus of points  $P$  such that the circumcircles of  $\triangle ABP$  and  $\triangle CDP$  are tangent has perimeter  $p$ , find  $\lfloor 100p \rfloor$ .

*Solution.*

Let  $X$  have coordinates  $(x, y)$ , and  $Y$  be the intersection of the radical axis of the two circumcircles and segment  $\overline{BC}$ . Let  $BY = b$ ; then  $CY = 5 - b$ . By Power of a Point, we have

$$XY^2 = BY \cdot AY = CY \cdot DY = b(4 + b) = (5 - b)(6 - b) \iff b = 2, XY^2 = 12$$

so  $X$  is always a distance of  $\sqrt{12}$  from the point  $Y = (3, 0)$ ; this is a circle with radius  $\sqrt{12}$  and so our answer is  $\boxed{2176}$ . □

### §13.2.7 The Party, The Politburo, and the PSC

Source: IZhO 2021/5

Proposer: Circumrectangular Hyperbola#8766 (446065841172250638)

Problem ID: 188

Date: 2021-05-02

Difficulty: Challenging

On a party with 102 guests, hosts Tony and Wang play a game (the hosts are not regarded as guests). There are 102 chairs arranged in a circle; initially, all guests hang around those chairs. The hosts take turns alternately. By a turn, a host orders any standing guest to sit on an unoccupied chair  $c$ . If some chair adjacent to  $c$  is already occupied, the same host orders one guest on such chair to stand up (if both chairs adjacent to  $c$  are occupied, the host chooses exactly one of them). All orders are carried out immediately. Tony makes the first move; her goal is to fulfill, after some move of hers, that at least  $k$  chairs are occupied. Determine the largest  $k$  for which Tony can reach the goal, regardless of Wang's play.

*Solution.*

We claim the answer is  $\boxed{35}$ . Note the number of chairs does not decrease, given that every empty chair must be next to a full chair, we have at least 34 chairs. In addition, when there are 33 chairs, every third chair is occupied.

If Wang forces Tony into such a configuration, say Tony placed a guest on chair  $x$  and Wang on  $y$ . However, if  $y$  is adjacent to  $x$ , Tony could have easily have chosen  $y$  instead and wins. Otherwise, she could have placed it on  $x \pm 1$ , as there were no guests there, and now not every third chair is occupied. Thus, Wang never reaches a position with 34 chairs such that no more guests can be placed for Tony, and Tony can happily place her last guest.

To show that Wang can win otherwise, we show inductively that Wang can hold a strategic position:

**Claim.** Wang can make sure that in the sets  $S_2, S_3, \dots, S_{34}$  where  $S_k = \{3k+1, 3k+2, 3k+3\}$ , none of  $3k+1$  and  $3k+3$  is occupied. Furthermore, there is at most two occupied chairs in  $S_1$ .

**Proof.** The proof is by induction.

*Base Case.* No moves are made, in which case the claim holds.

*Induction Hypothesis.* Assume that Wang holds such a position. We claim after 2 moves, Wang can hold this position again. *Induction Step.* If Alice's move holds the position, Wang can simply put a person somewhere in  $S_1$  (as chairs 4 and 102 are unoccupied, we can once again only have at most 2 people in  $S_1$ ). If it doesn't, she could not have operated on  $S_1$ , and thus placed a guest in some of  $S_2, \dots, S_{99}$ . However, if she operates on  $S_k$ , Wang places a person in the middle of  $S_k$  and removes the mischief caused by Alice. Thus Wang can hold his position.  $\square$

## §14 Mityushikha Bay (Season 14)

### §14.1 Week 1

#### §14.1.1 Rectangle Rotation

Source: Original

Proposer: RandomGuy#8751 (821212688633823235)

Problem ID: 189

Date: 2021-05-17

Difficulty: Beginner

A 35-by-42 rectangle is rotated around one of its vertices. If  $A$  is the area the rectangle sweeps out, find  $\lfloor 100A \rfloor$ .

(A scientific calculator may be used.)

*Solution.*

Noticably after a rectangle is rotated around one of its vertex, it will then form a circle.

The radius of the circle is a diagonal because the furthest point from one of its vertex is a diagonal and in a few rotation, it will cover an area of circle with the radius of the diagonal length.

Firstly, calculate the radius of the circle by calculating the rectangle diagonal

$$r = \sqrt{a^2 + b^2}$$

$$r = \sqrt{35^2 + 42^2}$$

$$r = \sqrt{2989}$$

so the area of the circle of  $2989\pi \iff \boxed{939022}$ .

*Remark.* 3.1415926 and  $\frac{355}{113}$  are the least "precise" approximations of  $\pi$  needed to get the correct answer.  $\square$

**§14.1.2 NJOY Troll (again)**

---

*Source: Original*

*Proposer: TaesPadhihary#8557 (665057968194060291)*

*Problem ID: 190*

*Date: 2021-05-18*

*Difficulty: Beginner*

---

If the minimum of  $(20 \cos x + 21)^2 + 2021$  is  $M$ , find  $\lfloor 100M \rfloor$ .

*Solution.*

The expression inside the square has a minimum of 1, achieved when  $\cos x = -1$ . Answer: 202200

□

## §14.1.3 You Just Got Vector'ed

Source: Sixth Term Examination Paper I, 2013 Q3

Proposer: sjbs#9839 (434767660182405131)

Problem ID: 191

Date: 2021-05-19

Difficulty: Easy

Given for any two points  $X$  and  $Y$ , with position vectors  $\mathbf{x}$  and  $\mathbf{y}$  respectively,  $X * Y$  is defined to be the point  $Z$  along the line  $\overrightarrow{XY}$  such that

$$\frac{XZ}{ZY} = \frac{\lambda}{1 - \lambda},$$

where  $\lambda$  is a fixed number.

The points  $P_1, P_2, \dots$  are defined by  $P_1 = X * Y$  and, for  $n \geq 2$ ,  $P_n = P_{n-1} * Y$ . Given that  $X$  and  $Y$  are distinct and that  $0 < \lambda < 1$ , the ratio in which  $P_n$  divides the line segment  $XY$  can be expressed in the form  $1 : A$ . If  $\lambda = 0.75$  and  $n = 69$ , find  $\lfloor 10^{12} A \rfloor$ .

(A scientific calculator may be used.)

*Solution.*

We have

$$P_1 = X * Y$$

$$P_2 = (X * Y) * Y$$

...

$$P_n = (X * Y) * Y) * Y) \dots * Y$$

Where  $P_n$  has  $n$   $Y$ 's. Substituting  $X * Y = \lambda \mathbf{x} + (1 - \lambda)\mathbf{y}$  and going through a number of iterations of  $*Y$ , we guess  $P_n = \lambda^n \mathbf{x} + (1 - \lambda^n)\mathbf{y}$ .

We now proceed by induction on  $n$ . Given we have already verified the case of  $n = 1$ , let  $n = k$ ,  $k \in \mathbb{N}$  and assume  $P_k = \lambda^k \mathbf{x} + (1 - \lambda^k)\mathbf{y}$ . Then for  $n = k + 1$ , we have  $P_{k+1} = P_k * Y$ , and so we have  $P_{k+1} = \lambda^{k+1} \mathbf{x} + \lambda(1 - \lambda^k)\mathbf{y} + (1 - \lambda)\mathbf{y}$ . This simplifies to  $P_{k+1} = \lambda^{k+1} \mathbf{x} + (1 - \lambda^{k+1})\mathbf{y}$ , as required. Therefore:

$$P_n = \begin{pmatrix} \lambda^n \\ 1 - \lambda^n \end{pmatrix}$$

Now it's simply a case of

$$XY : P_n = \begin{pmatrix} 1 \\ 1 \end{pmatrix} : \begin{pmatrix} \lambda^n \\ 1 - \lambda^n \end{pmatrix} \Rightarrow XY : P_n = 1 : \frac{\lambda^n}{1 - \lambda^n}$$

Hence when  $\lambda = 0.75$  and  $n = 69$ , we have  $\{10^{12} \cdot A\} = \boxed{2}394$

□

### §14.1.4 Answer Extraction

Source: Original

Proposer: Charge#3766(481250375786037258)

Problem ID: 192

Date: 2021-05-20

Difficulty: Medium

On a QoTD, the PSC has a probability  $\frac{m}{n}$  that they need to decide how to make people submit. Sjsb thinks that people should submit  $m + n$ . Brainy however, thinks that the people should submit  $100m + n$ . The rest of the PSC come up with an even better idea of people submitting  $420m + 69n$ .

If the PSC decides every single answer will be 420 in a season, the number of possible ways they can choose  $m$  and  $n$  and a way of formatting their answer (  $420m + 69n$ ,  $100m + n$ , or  $m + n$  ) in a 14 day season can be written as  $a^b$  where  $a$  and  $b$  are positive integers and  $b$  is maximized. Compute  $420a + 69b$ .

(The PSC can use a specific  $\frac{m}{n}$  more than once.)

*Solution.*

We can do some casework.

Case 1: The PSC choose to use the  $420m + 69n = 420$  system Clearly only  $m = 1, n = 0$  works, however this will not fit into the constraints.

Case 2: The PSC choose to use the  $100m + n = 420$  system Only  $(0, 420), (1, 320), (2, 220), (3, 120), (4, 20)$  work, and of these, only  $(1, 320)$  works.

Case 3: The PSC choose to use the  $m + n = 420$  system. For this to work we must have  $\gcd(m, n) = \gcd(m, 420 - m) = \gcd(m, 420) = 1$ . So to calculate the number of  $m$  possible, we do  $\varphi(420) = 420 \cdot \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{4}{5} \cdot \frac{6}{7} = 96$ . Since  $m$  must also be greater than  $n$  we divide by 2 to get 48 ways.

So our answer is  $49^{14} = 7^{28} \iff \boxed{4872}$ . □

## §14.1.5 Not a Fan

Source: Baltic Way 2020 Problem 18

Proposer: Circumrectangular Hyperbola#8766 (446065841172250638)

Problem ID: 193

Date: 2021-05-21

Difficulty: Hard

Let  $n \geq 1$  be a positive integer. We say that an integer  $k$  is a *fan* of  $n$  if  $0 \leq k \leq n-1$  and there exist integers  $x, y, z \in \mathbb{Z}$  such that

$$\begin{aligned}x^2 + y^2 + z^2 &\equiv 0 \pmod{n}; \\xyz &\equiv k \pmod{n}.\end{aligned}$$

Let  $f(n)$  be the number of fans of  $n$ . Determine  $f(2020)$ .

*Solution.*

We note that if  $4 \mid x^2 + y^2 + z^2$ , then clearly  $x \equiv y \equiv z \equiv 0 \pmod{2}$ , so  $xyz \equiv 0 \pmod{4}$ . Similarly, if  $5 \mid x^2 + y^2 + z^2$ , then as  $x^2 \in \{0, 1, 4\}$ , one of  $x, y, z$  is divisible by 5. Now, we claim we can find  $x, y, z$  such that  $x^2 + y^2 + z^2 \equiv 0 \pmod{101}$  and  $xyz \not\equiv 0 \pmod{101}$ . We can let  $\alpha = \sqrt{-1} \pmod{101}$  which has to exist as  $101 \equiv 1 \pmod{4}$ . Thus if  $101 \nmid z$ ,

$$(x - \alpha y)(x + \alpha y) = x^2 - (\alpha y)^2 = -z^2$$

We have  $101 - 1 = 100$  choices for  $x + \alpha y$ , and only two values of  $x + \alpha y$  force  $x + \alpha y = \pm(x - \alpha y)$ , implying that we have a solution such that  $0 \nmid x, y, z$  and

$$101 \mid x^2 + y^2 + z^2$$

Note that  $3 \nmid 101 - 1$ , so thus taking this solution and multiplying by an arbitrary number  $w$ , we see that  $(xw, yw, zw)$  works and thus spans all possible values of  $k \pmod{101}$ . Thus, there are 101 *fans* of 101, but only 1 of 20, so there are  $\boxed{101}$  of 2020.  $\square$



## §14.1.6 you're welcome, brainy

---

Source: 2019 Turkey Math Olympiad, P3

Proposer: Alp Eren#5330 (691708663462887464)

Problem ID: 194

Date: 2021-05-22

Difficulty: Challenging

---

There are 2021 crewmates in the universe, and some of these crewmates are members of different crews (crewmates can belong to multiple crews at once). Each crew has a security council consisting of 12 crewmates who are members of that particular crew. An *emergency meeting* (for a particular crew) can be realized only when each participant (of the meeting) is a member of that crew, and moreover, each of the 12 crewmates forming the security council (for that crew) are present among the participants. It is known that each subset of at least 12 crewmates can realize an emergency meeting for exactly one crew. If the total number of different crews with exactly 29 crewmates is  $N$ , find  $\nu_2(N) + \nu_3(N) + \nu_5(N)$ .

*Solution.*

AoPS Solutions

The answer is  $\binom{2003}{11} \iff \boxed{4}$ .

□

## §14.1.7 RDS-1

Source: 2007 Sharygin Correspondence Round 18

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 195

Date: 2021-05-23

Difficulty: Challenging

$O, G$  are points in the plane such that  $OG = 20.21$ . Let  $S$  be the set of vertices of all triangles with positive area that have circumcentre  $O$  and centre of mass  $G$ , and  $T$  the set of points in the plane that are not in  $S$ . If the area of  $T$  is  $U$ , find  $\lfloor 100U \rfloor$ .

(A scientific calculator may be used.)

*Solution.*


The official solution is in Russian. I don't understand Russian.

The main motivation here is the Euler Line and the points on it.

*Euler Line.*  $O, G, H$ , circumcentre, centre of mass, and orthocentre of a triangle, are collinear with  $2OG = GH$  or all three coincident. Consider the homothety that sends each vertex of  $\triangle ABC$  to the midpoint of its opposite side, thus sending  $\triangle ABC$  to its medial triangle; this homothety is then centered at  $G$  (why?). Then,  $H$  is mapped to the orthocentre of the medial triangle, but this point is the circumcentre of  $\triangle ABC$ .

*Lemma.* Let  $F$  be the midpoint of  $\overline{AB}$ . Then,  $CH = 2OF$ . This is closely related with the proof of the existence of the nine-point-circle. Consider the homothety centered at  $N_9$  that sends  $O$  to  $H$ ; this homothety has scale  $-1$ . Then,  $F$  is sent to the midpoint of  $\overline{CH}$  since the foot of the  $C$ -altitude is on the nine-point-circle, implying that  $F$  and the midpoint of  $\overline{CH}$  are diametrically opposite. We are done.

But note that  $OF < R$ , so we have the constraint  $CH = 2OF < 2R = 2OC$ . The locus for which  $\frac{CH}{OC} \leq 2$  is a union of Apollonian circles forming a disk with the boundary circle being the Apollonian circle where  $\frac{CH}{OC} = 2$ . Two diametrically opposite points on this circle are  $G$  and  $G'$ , where  $G'$  is on the Euler line and  $G'O = 60.63$ , making the disk have radius 40.42. To construct such  $\triangle ABC$ , let  $F$  be the point on  $\overrightarrow{CG}$  such that  $CF > CG$  and  $CG = 2GF$ . Then draw a line at  $F$  perpendicular to  $\overline{OF}$  and a circle centered at  $O$  with radius  $OC$ ; the points  $A, B$  are the intersections of circle and line.



We are missing one detail: what if  $A, B, C$  are all collinear? Then, since  $\overline{FO} \parallel \overline{CH}$  we have  $\angle CFO = \angle GCH$ , but since  $A, B, C$  collinear we have  $\angle CFO = 90$ , or,  $C$  is on the circle with diameter  $\overline{GH}$ , which in our case has radius 20.21 and does not intersect the aforementioned disk.

Our answer is  $100\pi (40.42^2 + 20.21^2) \iff \boxed{641582}$ .

*Remark.* RDS-1 was the first Soviet nuke.

□

**§14.2 Week 2****§14.2.1 All About that Base**

---

Source: Komal C1336

Proposer: flame#6784 (185229437787176962)

Problem ID: 196

Date: 2021-05-24

Difficulty: Beginner

---

Let  $a$ ,  $b$ , and  $c$  denote the digits 0, 1, and 2, in some order. If

$$aaa_3 \times bb_3 = acbab_3,$$

evaluate  $bcba_{10} \times bcbb_{10}$ .

(A 4-function calculator may be used.)

*Solution.*

If  $a$  or  $b$  are 0, then  $aaa_3 \times bb_3 = 0_3$ , which is clearly false. Hence  $c = 0$ . Looking at the units digits,  $a_3 \times b_3 = b_3$ , so  $a = 1$  and  $b = 2$ . (It can be checked that  $111_3 \times 22_3 = 10212_3$ .) We seek  $2021 \times 2022 = \boxed{4086462}$ .  $\square$

## §14.2.2 Number of QoTD Seasons

Source: Original

Proposer: cooked#8292 (573431880091828224)

Problem ID: 197

Date: 2021-05-25

Difficulty: Beginner

A QoTD season, lasting 14 days, has 3 alg, 3 combi, 3 geo and 3 NT problems, as well as 2 other problems in any of alg, combi, geo or NT. If  $N$  is the number of distinct possible QoTD topic sequences, find  $\nu_2(N) + \nu_3(N) + \nu_5(N)$ . (Each question belongs to exactly one subject, so no CG or anything.)

*Solution.*

Without loss of generality, we can consider two cases- the case that both problems are the same topic and the case that both problems are in different topics.

Consider the case the 2 problems are in the same topic. Without loss of generality, let us call the 4 topics A, B, C and D, and let us assume the 2 problems are in topic A, so that there are 5 A problems, 3 B problems, 3 C problems and 3 D problems. Now, since this is permutations of like objects, we can see the number of topic sequences using A, B, C and D is

$$\frac{14!}{5! \times 3! \times 3! \times 3!}$$

, which is 3363360. We can also see that there are  $4!$  ways to map Alg, Combi, Geo and NT to the topics A, B, C and D, so there are actually 3363360

×

$4!$  topic sequences, which is 80720640.

Now consider the case the two problems are in different topics. Without loss of generality, let us call the 4 topics A, B, C and D, and let us assume the two problems are in topic A and B respectively, so that there are 4 A problems, 4 B problems, 3 C problems and 3 D problems. Like before, we can see the number of topic sequences using A, B, C and D is

$$\frac{14!}{4! \times 4! \times 3! \times 3!}$$

is 4204200. Like before, there are  $4!$  ways to map Alg, Combi, Geo and NT to the topics A, B, C, and D, so there are actually 4204200

×

$4!$  topic sequences, which is 100900800.

Add to get

$$\frac{14! \cdot 4!}{3!^3 \cdot 4!} \left( \frac{1}{5} + \frac{1}{4} \right) = 2^6 \cdot 3^4 \cdot 5 \cdot 7^2 \cdot 11 \cdot 13 \iff \boxed{11}.$$

□

## §14.2.3 Square and Point

Source: Original

Proposer: RandomGuy#8751 (821212688633823235)

Problem ID: 198

Date: 2021-05-26

Difficulty: Beginner

There exist a square  $ABCD$  with its side length 5 and there also exist a random point  $P$  on the same plane such that  $AP = 1$ .

Let  $M$  be the longest distance possible between  $P$  and  $C$ . Let  $m$  be the shortest distance possible between  $P$  and  $C$ .

Find  $\lfloor 1000M + 1000m \rfloor$ .

(A scientific calculator may be used.)

*Solution.*

We notice that the distance  $CP$  could be maximum and minimum if  $C$ ,  $A$ , and  $P$  are collinear. This can be proved by drawing a circle around  $A$ . Then, we continue by finding  $M$  and  $m$

$$M = \sqrt{5^2 + 5^2} + 1 = 5\sqrt{2} + 1$$

$$m = \sqrt{5^2 + 5^2} - 1 = 5\sqrt{2} - 1$$

$$\text{Hence, } M + m = 5\sqrt{2} + 1 + 5\sqrt{2} - 1 = 10\sqrt{2} = \sqrt{200} \iff \boxed{14142}.$$

□

## §14.2.4 2012 HMMT A6

Source: Harvard-MIT Mathematics Tournament 2012 February A6

Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 199

Date: 2021-05-27

Difficulty: Medium

Let  $a_0 = -2$ ,  $b_0 = 1$ , and for  $n \geq 0$ , let

$$a_{n+1} = a_n + b_n + \sqrt{a_n^2 + b_n^2}$$

$$b_{n+1} = a_n + b_n - \sqrt{a_n^2 + b_n^2}$$

If  $a_{2020}$  can be represented in the form  $a^x \sqrt{b} - c^y$ , where  $a, b, c$  are integers such that  $a, c$  are minimized and  $c$  is square free. Find  $x + y$ .

*Solution.*

By adding the two equations and multiplying them respectively, we get

$$a_{n+1} + b_{n+1} = 2(a_n + b_n)$$

$$a_{n+1}b_{n+1} = (a_n + b_n)^2 - (a_n^2 + b_n^2) = 2a_nb_n$$

Thus, we get that

$$a_{2020} + b_{2020} = -2^{2020}$$

$$a_{2020}b_{2020} = -2^{2021}$$

Either by vieta's or substitution, we get  $a_{2020}, b_{2020}$  are the roots of the quadratic

$$x^2 + 2^{2020}x - 2^{2021}$$

where  $a_{2020}$  is the larger root since the positive sqrt was added. Solving the quadratic gives that

$$a_{2020} = 2^{1010} \sqrt{2^{2018} + 2} - 2^{2019}$$

Thus, the answer is 3029

□

§14.2.5 [under construction]



## §14.2.6 Projective Pain

Source: Unknown

Proposer: ChristopherPi#8528 (696497464621924394)


Problem ID: 201

Date: 2021-05-22

Difficulty: Challenging

$\triangle ABC$  has circumcircle  $\Gamma$  and incentre  $I$ , with sidelengths  $AB = 44, BC = 37, CA = 15$ . Let  $L$  be the midpoint of the arc  $\widehat{BC}$  containing  $A$ .  $\overleftrightarrow{LI}$  meets  $\Gamma$  again at  $D$ , and the tangent to  $\Gamma$  at  $A$  meets  $\overleftrightarrow{BC}$  at  $P$ .  $\overleftrightarrow{PD}$  meets  $\Gamma$  again at  $X$ . If  $Y$  is the point on  $\overline{BC}$  such that  $\angle BAX = \angle CAY$ , the ratio  $\frac{BY}{CY}$  can be expressed in the form  $\frac{m}{n}$ , where  $m, n$  are relatively prime positive integers. Find  $100m + n$ .

Solution.



We claim that for a general cevian  $\ell$  through  $A$ ,  $Y$  will be the reflection across the midpoint  $M$  of  $\overline{BC}$  of the intersection of the isogonal line of  $\ell$  and  $\overline{BC}$ . Consider the following transformation that sends points on  $\overline{BC}$  to points on  $\overline{BC}$ : take the projections of points wrt  $A$  onto  $\Gamma$ , invert  $\triangle ABC$  and those points wrt  $P$ , project points back onto  $\overline{BC}$  wrt  $A$ , then invert the points back onto the original  $\triangle ABC$ . Let  $\overline{AD}$  intersect  $\overline{BC}$  at  $D_1$ , and let  $D_2$  be the intersection of the isogonal line of  $\overline{AD}$  and  $\overline{BC}$ . Applying this transformation to  $\overline{AD}$  and  $\overline{AX}$ , we let  $D', X', B', C', D'_2$  be the images of  $D, X, B, C, D_2$  under inversion, and  $Y'$  the intersection of  $\overline{AD'}$  and  $\overline{B'C'}$ ; by angle chase we have  $\overline{AX'}$  isogonal to  $\overline{AD}$  and  $\overline{AD'}$  isogonal to  $\overline{AX}$ . By inversion distance formula we find

$$\frac{BY}{CY} = \frac{CD_2}{BD_2} = \frac{PB}{PC};$$

furthermore since  $\overline{PA}$  is a tangent we have  $\overline{A'C'} \parallel \overline{AB}, \overline{A'B'} \parallel \overline{AC}$  and  $\overline{AD'} \parallel \overline{AX}, \overline{AX'} \parallel \overline{AD}$  so by similar triangles and inversion distance formula we have

$$\frac{C'D'_2}{A'D'_2} = \frac{BD_1}{AD_1}, \frac{B'D'_2}{A'D'_2} = \frac{CD_1}{AD_1} \iff \frac{C'D'_2}{B'D'_2} = \frac{PB}{PC} = \frac{BD_1}{AD_1} = \frac{CD_2}{BD_2}$$

and we are done.

In the case of this problem, it is well-known (in EGMO; invert at  $A$ ) that  $D$  is the touchpoint of the  $A$ -mixtilinear incircle and  $\Gamma$ , and so the isogonal line passes through the  $A$ -excircle touchpoint, so  $Y$  is actually the incircle touchpoint on  $\overline{BC}$  so our answer is  $\frac{s-b}{s-c} = \frac{33}{4} \iff \boxed{3304}$ .  $\square$

## §14.2.7 AN602

Source: 2021 Bundeswettbewerb Mathematik R1P4, Modified

Proposer: Boldizsár#3106 (694967174921715853)

Problem ID: 202

Date: 2021-05-23

Difficulty: Challenging

Consider a pyramid with a regular  $n$ -gon as a base. Draw a line segment in either blue or red between any two points among the  $n$  vertices of the base and the apex, except for the edges of the base, which are not drawn at all. Find the smallest integer  $n$  with the following property: There exist 4 points among the aforementioned  $n + 1$  points such that all six segments connecting those four points are drawn in the same color.

*Solution.*

The solution is  $n = \boxed{33}$ . We first show that  $n = 32$  does not work, i. e. that we can draw the pyramid with a 32-gon as a base such that there is no complete quadrilateral in a single color. By  $R(4, 4) = 18$ , there exists a complete graph  $G$  with 17 vertices, the edges drawn in red and blue that contains no  $K_4$  (the complete graph with four vertices) drawn in only one color. Now, organise the 32 vertices of the base into pairs, each pair consisting of two neighboring vertices. We have 17 units now, 16 pairs of neighboring base vertices and one apex. Now, match every unit with one vertex of  $G$  and connect two units in the color in which the edge connecting the corresponding vertices of  $G$  is drawn. Connect two vertices of the pyramid in the color in which the units they are in are connected. As there are no edges drawn within units (because those are neighboring vertices of the base), our construction will indeed work. Now show that for  $n = 33$ , there indeed is a  $K_4$  as a subgraph drawn in a single color. So consider the pyramid with the regular 33-gon as a base. WLOG, 17 vertices are connected with the apex in red. But among those 17 vertices, there are 9 from which any two are connected. But  $R(4, 3) = 9$ , so those 9 vertices contain either a red  $K_3$  or a blue  $K_4$ , where in the first case the pyramid contains a red  $K_4$ , as desired.

This problem is almost impossible without the knowledge of Ramsey numbers, but it can be done; the theory required to discover isn't all that much and you can always search up  $R(3, 4)$  and  $R(4, 4)$ .

*2-colour Ramsey numbers.*  $R(b, w)$  is the smallest positive integer  $n$  such that any edge colouring in black and white of the complete graph on  $n$  vertices  $K_n$  will always contain either an induced subgraph  $K_b$  of all black edges or an induced subgraph  $K_w$  of all white edges. For example,  $R(3, 3) = 6$ . To see this, fix one vertex  $v$ ; by pigeonhole at least three of its edges will be of one color, WLOG black. Let the other vertices of those black edges be  $x, y, z$  respectively. If none of the edges formed by connecting any of  $x, y, z$  together are black, then  $x, y, z$  is  $K_3$ ; if any of them are then  $v$  and those two vertices forming the other black edge is  $K_3$ . To prove  $R(3, 3) \neq 5$  we can construct a counterexample: take  $K_5$  as a pentagon; the perimeter of the pentagon is coloured black while all the interior diagonals are coloured white. Convince yourself that this works.

*Ramsey's Theorem.*  $R(b, w) \leq R(b, w - 1) + R(b - 1, w)$ . We proceed by induction on  $b + w$ . First, convince yourself that  $R(b, 2) = b$ ; this is our base case. Now consider the complete graph on  $R(b, w - 1) + R(b - 1, w)$  vertices with edges coloured in black or white. Fix one vertex  $v$ , and let  $B$  be the set of all vertices whose edges with  $v$  are black, and  $W$  the set of all vertices whose edges with  $v$  are white. Since  $R(b, w - 1) + R(b - 1, w) = |B| + |W| + 1$  we have either  $|B| \geq R(b - 1, w)$  or  $|W| \geq R(b, w - 1)$ ; WLOG the first case is true. If we have  $K_w$  with edges all white we're done; otherwise we have  $K_{b-1}$  with edges all black; the add in vertex  $v$  and all the black edges connected to  $v$ , which are all from  $B$ ; thus we have  $K_b$  all black edges and we're done. If both  $R(b, w - 1)$  and  $R(b - 1, w)$  are even, the inequality strengthens to  $R(b, w) < R(b, w - 1) + R(b - 1, w)$ . Consider the complete graph on  $R(b, w - 1) + R(b - 1, w) - 1$  vertices with edges coloured in black or white. Order the vertices, and let  $b_i$  be the *black-degree* of vertex  $v_i$ ; that is, the number of black edges

emanating from  $v_i$ . By handshaking lemma there are an even number of vertices with odd degree; since  $R(b, w-1) + R(b-1, w) - 1$  is odd there must exist at least one vertex with even degree; WLOG it's  $v_1$ . Then  $|B| = b_i$ ,  $|W| = R(b, w-1) + R(b-1, w) - 2 - b_i$  are both even, so by parity it is impossible for both  $|B| < R(b-1, w) - 1$  and  $|W| < R(b, w-1)$  and we are done.

$R(3, 4)$  is 9. Since  $R(3, 3) = 6$  and  $R(2, 4) = 4$  are both even we conclude  $R(3, 4) < 10$ . We can colour a regularly octagonal  $K_8$  as follows: all perimeter edges are black, and all edges connecting diametrically opposite vertices (vertices 4 apart from another) are black; everything else is white. This way, the only way to get from vertex to vertex along black edges is by moving along by 1 or 4 vertices; no combination of three ones and fours will be zero mod 8, so there is no black  $K_3$ ; similarly, the only way to get from vertex to vertex along white edges is by moving along by 2 or 3 vertices, and the only 4-tuplet of twos and threes that is zero mod 8 are  $(2, 2, 2, 2)$  which clearly doesn't work since edges joining antipodal vertices are coloured black.

$R(4, 4)$  is 18. We know that  $R(4, 4) \leq 2R(3, 4) = 18$ . We can colour a regularly 17-gonal  $K_{17}$  as follows: all edges that connect vertices that are a power of two apart are black and everything else is white. The only combination of  $\{1, 2, 4, 8\}$  that is zero mod 17 is  $(8, 4, 4, 1)$  and that doesn't work because a pair of vertices are a distance of twelve apart, and that edge would be coloured white. Similarly, the only combination of  $\{3, 5, 6, 7\}$  that is zero mod 17 is  $(3, 3, 5, 6)$  and that doesn't work because a pair of vertices are a distance of eight apart, and that edge would be coloured black.

*Remark.* Mityushikha Bay was the testing site for AN602 (nicknamed the Tsar Bomba), along with many other Russian nukes.  $\square$

## §15 SALT-1 (Season 15)

### §15.1 Week 1

#### §15.1.1 "What are distinct integers?"

---

*Source: 2018 Fall OMO 1*

*Proposer: TaesPadhiary#8557 (665057968194060291)*

*Problem ID: 204*

*Date: 2021-06-07*

*Difficulty: Beginner*

---

Call two integers *primely* if 1 is the greatest integer dividing both of them. Find the minimum possible sum of 2021 nonnegative, primely integers.

*Solution.*

One zero and 2020 ones. Answer:

□

## §15.1.2 Troll Pigeonhole

---

Source: Original Problem

Proposer: Ada Virus#6266 (439054981966856202)

Problem ID: 205

Date: 2021-06-08

Difficulty: Beginner

---

In the following fraction every letter represents a different decimal digit.

$$\frac{B \cdot L \cdot U \cdot E \cdot B \cdot E \cdot R \cdot R \cdot Y}{I \cdot C \cdot E \cdot C \cdot R \cdot E \cdot A \cdot M}$$

If the sum of all possible real values this expression can take is  $S$ , find  $\lfloor 100S \rfloor$ .

*Solution.*

There are exactly 10 different letters (A, B, C, E, I, L, M, R, U, Y), so one of the letters must be 0. 0 divided by anything (except 0 which will make it undefined) is 0 so the answer is 0. □

## §15.1.3 WA Can't Solve This

Source: 2019 Pui Ching Contest

Proposer: QuantumSigma#9996 (374937087301910528)

Problem ID: 206

Date: 2021-06-09

Difficulty: Easy

$$\frac{69}{420} + \frac{69 \cdot 70}{420 \cdot 421} + \frac{69 \cdot 70 \cdot 71}{420 \cdot 421 \cdot 422} + \cdots = \frac{m}{n},$$

where  $m, n$  are relatively prime positive integers. Find  $100m + n$ .

Solution.

$$\begin{aligned} & \frac{69}{420} + \frac{69 \cdot 70}{420 \cdot 421} + \frac{69 \cdot 70 \cdot 71}{420 \cdot 421 \cdot 422} + \cdots \\ &= \frac{1}{350} \left[ \frac{69 \cdot (420 - 70)}{420} + \frac{69 \cdot 70 \cdot (421 - 71)}{420 \cdot 421} \right. \\ & \quad \left. + \frac{69 \cdot 70 \cdot 71 \cdot (422 - 72)}{420 \cdot 421 \cdot 422} + \cdots \right] \\ &= \frac{1}{350} \left[ \frac{69 \cdot 420}{420} - \frac{69 \cdot 70}{420} + \frac{69 \cdot 70 \cdot 421}{420 \cdot 421} - \frac{69 \cdot 70 \cdot 71}{420 \cdot 421} \right. \\ & \quad \left. + \frac{69 \cdot 70 \cdot 71 \cdot 422}{420 \cdot 421 \cdot 422} - \frac{69 \cdot 70 \cdot 71 \cdot 72}{420 \cdot 421 \cdot 422} + \cdots \right] \\ &= \frac{69}{350} \end{aligned} \tag{13}$$

Thus, the answer is 7250.

□

**§15.1.4 Prime Product**

---

Source: HMMT-Feb-2018-AN4

Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 207

Date: 2021-06-10

Difficulty: Medium

---

Distinct prime numbers  $p, q, r$  satisfy the equation

$$2pqr + 50pq = 7pqr + 55pr = 8pqr + 12qr = A$$

for some positive integer  $A$ . What is  $A$ ?

*Solution.*

Comparing the first two equations, we get

$$\begin{aligned} 50pq - 55pr &= 5pqr \iff \\ 10q - 11r &= qr \iff \\ (q + 11)(r - 10) &= -110 \end{aligned}$$

Since  $q, r$  are positive integers,  $(r - 10)$  must be negative and  $(11 + q)$  must be a divisor of 110 larger than 11. A quick check gives that  $q$  must be 11, which gives that  $r = 5$ . Plugging these values into the original equation gives  $p = 3$  and thus  $A = \boxed{1980}$ .

□

### §15.1.5 The Rare Non-Horrible States Problem

Source: David Patrick

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 208

Date: 2021-06-11

Difficulty: Hard

We play a game. The pot starts at \$0. On every turn, you flip a fair coin. If you flip heads, I add \$100 to the pot. If you flip tails, I take all of the money out of the pot, and you are assessed a “strike.” You can stop the game before any flip and collect the contents of the pot, but if you get 3 strikes, the game is over and you win nothing. Find, with proof, the expected value of your winnings if you follow an optimal strategy.

*Solution.*

We work backwards. If we have only one strike left, when is it advantageous to flip again? Suppose there are  $d$  dollars in the pot. Then, the expected pot value of our next flip is  $\frac{1}{2}(d + 100)$ , so we want to solve  $\frac{1}{2}(d + 100) > d \iff d < 100$ . However, the only value for which  $d < 100$  is  $d = 0$ , so, with only one strike remaining, the best strategy is to flip once and end the game, no matter what the result; our expected earnings is \$50.

What about with two strikes? Suppose there are  $d$  dollars in the pot. This time, the expected pot value is  $\frac{1}{2}(d + 100) + \frac{1}{2} \cdot 50$ , since if it's heads we add \$100 and if we lose we play as if we have one strike left, which optimal expected earnings is \$50 as covered before. Solving  $d + 100 + 50 > 2d$  we have  $d < 150$ . Now, we can split into cases: the result of our first coin, and the result of our second (if the first coin is heads). Our expected earnings are then  $\frac{1}{2} \cdot 50 + \frac{1}{2} \left( \frac{1}{2} \cdot 200 + \frac{1}{2} \cdot 50 \right) = 87.5$ .

For three strikes, it's exactly the same: solve  $\frac{1}{2}(d + 100) + \frac{1}{2} \cdot 87.5 > d \iff d < 187.5$ ; then our expected earnings are  $\frac{1}{2} \cdot 87.5 + \frac{1}{2} \left( \frac{1}{2} \cdot 200 + \frac{1}{2} \cdot 87.5 \right) = 115.625 \iff \boxed{11562}$ .  $\square$



## §15.1.6 Minimal Area, Minimal Difficulty for a Saturday

Source: Original (?)

Proposer: Matteddy#0482 (329956567132930048)


Problem ID: 209

Date: 2021-06-12

Difficulty: Hard

Right isosceles  $\triangle ABC$  has area 1200.  $X, Y, Z$  are points on sides  $\overline{AB}, \overline{BC}, \overline{CA}$  respectively such that  $\triangle XYZ$  is also right and isosceles. Find the minimal area of  $\triangle XYZ$ .

Solution.



Let  $\angle B$  be the right angle. If  $Z$  is the right angle of  $\triangle XYZ$ , then we leave it as an exercise to the reader to show that the minimal configuration happens when  $\triangle XYZ$  is the medial triangle of  $\triangle ABC$ . However, there is another case: the right angle is on one of the legs. Let  $Y$  be the right angle; then let  $\angle XYB = a$ . Through some angle chasing, we find  $\triangle AXZ \sim \triangle CZY$  with ratio  $\sqrt{2}$ . Letting  $s$  be the leg length of  $\triangle ABC$  and  $x$  the leg length of  $\triangle XYZ$  we have

$$BY = x \cos(a) \iff YC = s - x \cos(a) \iff AZ = \sqrt{2}(s - x \cos(a))$$

and

$$BX = x \sin(a) \iff XA = s - x \sin(a) \iff CZ = \frac{s - x \sin(a)}{\sqrt{2}}$$

and so

$$AZ + CZ = AC = s\sqrt{2} \iff s = x(2 \cos(a) + \sin(a)).$$

The maximum of  $2 \cos(a) + \sin(a)$  is  $\sqrt{5}$  (use Cauchy-Schwarz or properties of combined sinusoidal functions) so our minimum  $x$  is  $\frac{s}{\sqrt{5}}$  and our minimum area is  $\frac{[ABC]}{5} = \boxed{240}$ .  $\square$

**§15.1.7 Detente**

---

Source: 2016 CMIMC N10

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 210

Date: 2021-06-13

Difficulty: Challenging

---

Let

$$S = \sum_{n=1}^{\infty} \text{lcm}[n, 69]^{-2}.$$

Find  $\lfloor 10^9 S \rfloor$ .

(A scientific calculator may be used.)

Hint.  $\sum_{n=1}^{\infty} \text{lcm}[n, 1]^{-2} = \frac{\pi^2}{6}$ .

Solution.

**Solution** Answer:

□

**§15.2 Week 2****§15.2.1 Senior Math; What Challenge?**

---

*Source: 1998 SMC*

*Proposer: sjbs#9839 (434767660182405131)*

*Problem ID: 211*

*Date: 2021-06-14*

*Difficulty: Beginner*

---

A cube is inscribed in a sphere of diameter 1. What is the surface area of the cube?

*Solution.*

This means that the cube has space diagonal 1, so  $3s^2 = 1 \iff s^2 = \frac{1}{3} \iff SA = 6s^2$ . Answer: 4 □

**§15.2.2 Prisoner's Stupid Dilemma**

---

*Source: Original**Proposer: brainysmurfs#2860 (281300961312374785)**Problem ID: 212**Date: 2021-06-15**Difficulty: Beginner*

---

There are 100 prisoners numbered 1 to 100, and 100 boxes also labelled 1 to 100. The boxes are kept inside a room and the numbers 1 to 100 are randomly placed inside each box (one number in each box, and each number appears exactly once). In sequence, each prisoner enters the room and is allowed to look inside 50 boxes. If they find their number, they are released and allowed to communicate with the next prisoners. Otherwise, all the prisoners are executed. If the prisoners play optimally, the probability they are all released is  $p$ . Find  $\lfloor 1000p \rfloor$ .

*(A scientific calculator may be used.)*

*Solution.*

Suppose Brainy is the first prisoner. He clearly has no better strategy than random guessing, getting his number with probability  $\frac{1}{2}$ . If he does get his number, then he should remember the numbers in the remaining 49 boxes that he opened and tell that to the remaining 99 prisoners. Then, each prisoner will either know which box to open (because Brainy opened the box with their number in it) or they will know 50 boxes that don't contain their number (Brainy doesn't mention their number, so of the 50 boxes he opened none of them contain their number). But this means that they can all successfully escape! Our answer is 500.  $\square$

**§15.2.3 Coordinates without Bash**

---

*Source: Original Problem*

*Proposer: nexal#3868 (381673088216989742)*

*Problem ID: 213*

*Date: 2021-06-16*

*Difficulty: Easy*

---

There is a circle that goes through the points  $(2 + \sqrt{2}, 1 - \sqrt{7})$  and  $(5, 1)$  with radius 3. Let  $(a, b)$  be the circle's center. Given  $a + b$  is a positive integer, find  $a^b + b^a$ .

*Solution.*

Notice that  $\sqrt{2}^2 + \sqrt{7}^2 = 3^2$ , so we try and see that a circle centered at  $(2, 1)$  does go through  $(2 + \sqrt{2}, 1 - \sqrt{7})$ . Using the distance formula, we see it also goes through  $(5, 1)$ . The other and only possible circle has its center reflected across the midpoint of the two given points. Since one point has lattice coordinates and the other is irrational, the midpoint must have irrational coordinates. Therefore, this other circle's center has irrational coordinates. Since 2 and 7 are relatively prime, it is impossible for the sum of the this center's coordinates to be an integer. Therefore the circle centered at  $(2, 1)$  is the only possible circle, giving us the answer of  $2^1 + 1^2 = \boxed{3}$ .  $\square$

### §15.2.4 WA Can't Solve This, Either

Source: 2013 HMMT A7

Proposer: Angry Any#4319 (580933385090891797)

Problem ID: 214

Date: 2021-06-17

Difficulty: Medium

$$\sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \cdots \sum_{k_8=0}^{\infty} \frac{k_1 + k_2 + \cdots + k_8}{5^{k_1+k_2+\cdots+k_8}} = \frac{m}{n},$$

where  $m, n$  are relatively prime positive integers. Find the sum of positive divisors of  $mn$ .

(A scientific calculator may be used.)

Solution.

Solution Answer:

□

**§15.2.5 WA Can't Solve This, Either**

---

Source: USAMTS Y19R1P3

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 215

Date: 2021-06-18

Difficulty: Hard

---

Let  $S$  be the set of all ordered triples of positive integers  $(a, b, c)$  such that

$$\arctan \frac{1}{a} + \arctan \frac{1}{b} + \arctan \frac{1}{c} = \frac{\pi}{4},$$

where the range of  $\arctan(x)$  is  $[-\pi/2, \pi/2]$ . Find

$$\sum_{(a,b,c) \in S} abc.$$

*Solution.*

**Solution** Answer: 1293

□

## §15.2.6 cHriS geO

Source: 2020 Iran MO R3G4

Proposer: ChristopherPi#8528 (696497464621924394)

Problem ID: 216

Date: 2021-06-19

Difficulty: Challenging

$\triangle PQR$  has  $PQ = 6, QR = 7, RP = 8$ , incentre  $I$ , and circumcentre  $O$ . The external angle bisector of  $P$  meets  $\overleftrightarrow{QR}$  at  $S$ , and  $I_Q$  is the  $Q$ -excentre. The point  $T$  is chosen along  $\overleftrightarrow{IQ}$  such that  $QT = 2IQ$  and  $IT > IQ$ . Let  $F$  be the point on the circumcircle of  $\triangle DI_QT$  such that  $DF$  is a diameter. Then the perimeter of  $\triangle OIF$  can be expressed in the form  $\frac{a+b\sqrt{c}}{d}$ , where  $\gcd(a, b, d) = 1$ ,  $a, b$  are integers, and  $c, d$  are positive integers. Find  $1000a + 100b + 10c + d$ .

Solution.

**Solution** so  $O, I, F$  are collinear; using  $OI^2 = R^2 - 2Rr$  we get  $IF = \frac{16\sqrt{15}}{15} \iff \boxed{3530}$ . □



## §15.2.7 Treaty Violation

Source: chinese guy

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 217

Date: 2021-06-20

Difficulty: Challenging

Let  $f : \mathbb{R}^{2020} \rightarrow \mathbb{R}$  be the function

$$f(a_1, a_2, \dots, a_{2020}) = \frac{a_1^2 + a_2^2 + \dots + a_{2020}^2 + (a_1 + a_2 + \dots + a_{2020})^2}{\sqrt{a_1^4 + a_2^4 + \dots + a_{2020}^4 + (a_1 + a_2 + \dots + a_{2020})^4}}.$$

Then the maximum of  $f$  is  $M$ , and the minimum of  $f$  is  $m$ . Find  $\lfloor 10^9 M + 10^6 m \rfloor$ .

(A scientific calculator may be used.)

*Solution.*

In fact,  $f$  has the range

$$\left[ \sqrt{\frac{m(m-1)}{m^2-3m+3}}, \sqrt{\frac{m(m^2-1)}{m^2+3}} \right]$$

where  $m = n + 1$ , giving us an answer of  $\boxed{44956512067}$ . The condition is equivalent to  $a_1 + \dots + a_m = 0$ , where  $m$  is odd. To find the minimum, WLOG  $a_m$  be the largest term in absolute value (positive or negative) then use C-S on

$$a_1^2 + \dots + a_{m-1}^2 = (a_1 + \dots + a_{m-1})^2 - 2 \sum_{i < j} a_i a_j = a_m^2 - 2 \sum_{i < j} a_i a_j$$

to arrive at an answer. The end result is that we should maximize  $a_m^2$  while minimizing  $a_1^2 + \dots + a_{m-1}^2$ .

Unfortunately, the maximum is not so easy. Intuitively from QM-AM, the maximum should happen when the  $a$ 's are close together; in fact, the equality case happens when there are  $\lfloor m/2 \rfloor$  equal terms of one sign and  $\lceil m/2 \rceil$  equal terms of the other. The proof of this is absolute hell and involves 4 different inequalities (one that you have to make up yourself) and smoothing/mixing variables, so just trust that the fakesolve works.  $\square$

## §16 Quality Control (Season 16)

### §16.1 Week 1

#### §16.1.1 The Rare non-Troll NJOY Problem

---

*Source: Unknown*

*Proposer: TaesPadhiary#8557 (665057968194060291)*

*Problem ID: 218*

*Date: 2021-06-28*

*Difficulty: Beginner*

---

If  $n^2\%$  of  $n$  is 80, find  $n$ .

*Solution.*

We have

$$\frac{n^2}{100} \cdot n = 80 \iff n = \boxed{20}.$$

□

### §16.1.2 The Common Troll NJOY Problem

Source: 2017 Fall OMO #4

Proposer: TaesPadhiary#8557 (665057968194060291)

Problem ID: 219

Date: 2021-06-29

Difficulty: Beginner

Chris draws a line segment between each of the points


$$A(4, 4), B(-4, 4), C(-4, -4), D(4, -4), E(2, 0), F(0, 2), G(-2, 0), H(0, -2)$$

on a sheet of paper large enough so that none of the line segments touch the edge of the paper. He wants to separate all the regions obtained by the line segments and later on eat them.

What is the maximum number of regions of paper that Chris can eat?

*Solution.*

Note that the diagram is diagonally symmetric, so we need only to look at the intersections in the interior of  $\triangle AOB$ , where  $O$  is the origin. Count that there are 15 intersections, then multiply by 4 to get 60. However, the troll is that Chris is cutting out sections on a piece of paper, so the outside of square  $ABCD$  is also a region and our answer is 61.



□

## §16.1.3 Parallelogram PTSD

Source: Italian Team Competition

Proposer: HoboSas#3200 (310725130097786880)

Problem ID: 220


Date: 2021-06-30

Difficulty: Medium

Diagonals of parallelogram  $NJSM$  intersect at  $E$ , angle bisectors of  $\angle MNE$  and  $\angle EJS$  intersect at  $L$ . Given that  $MESL$  is a parallelogram and  $\overline{NM} = 35$ , compute  $\overline{NJ}^2$ .

*Solution.*

Note  $NS \parallel ML$  so  $\angle NLM = \angle LNE = \angle MNL$  which makes  $\triangle MNL$  isosceles, hence  $\overline{MN} = \overline{ML} = \overline{ES}$ . With a similar reasoning we get  $\triangle JSL$  isosceles, with  $\overline{JS} = \overline{SL} = \overline{EM}$ , but  $\overline{MN} = \overline{JS}$  so  $\overline{EM} = \overline{ES}$ , but recalling  $E$  is the midpoint of both diagonals,  $\overline{NS} = \overline{JM}$ , making  $NJSM$  a rectangle. Now  $\overline{MJ} = 2 \cdot \overline{MN}$ , which gives  $\overline{NJ} = \sqrt{3} \cdot \overline{NM}$ , so finally  $\overline{NJ}^2 = 3 \cdot \overline{NM}^2 = \boxed{3675}$ .  $\square$



**§16.1.4 WA Could Solve This and you Still Failed***Source: Original**Proposer: AiYa#2278 (675537018868072458)**Problem ID: 221**Date: 2021-07-01**Difficulty: Medium*

A real number  $a$  is selected uniformly at random between  $-50$  and  $50$  inclusive. The probability that  $20x - |21x - |x + \frac{20a}{21}|| = 42|x - 1|$  has real solutions can be expressed in the form  $\frac{m}{n}$ , where  $m$  and  $n$  are relatively prime positive integers. Find  $100m + n$ .

*Solution.*

Let the LHS be  $f(x)$ . The graph of  $f$  looks like a bunch of lines, and  $f$  is continuous (why?). The maximum absolute value of the slope of a line that's part of  $f$  can be is 42, which means that if  $f$  crosses the RHS then  $f(1) \geq 0$ , since  $x = 1$  is where the minimum of  $42|x - 1|$  is attained; if  $f(1) < 0$  then  $f$  will never cross  $42|x - 1|$ . This is because for  $f$  to cross  $42|x - 1|$  for  $x < 1$  when  $f(1) < 0$ ,  $f$  would have to somehow go from being above the V-shaped absolute value graph of  $42|x - 1|$  to below it at  $x = 1$ , impossible since no part of  $f$  has slope of absolute value greater than 42; analogously we see that  $f$  cannot cross  $42|x - 1|$  when  $x > 1$ . Solve to get  $-\frac{441}{10} \leq a \leq -\frac{21}{10}$  and  $0 \leq a \leq 42$ . Our probability is  $\frac{21}{25} \iff \boxed{2125}$ .  $\square$

### §16.1.5 Surprising System

Source: AoPS Mock AIME

Proposer: AiYa#2278 (675537018868072458)

Problem ID: 222

Date: 2021-07-02

Difficulty: Hard

$p, q, r$  are reals such that the following system holds:

$$\begin{aligned}(p+q)(q+r) &= -1 \\ (p-q)^2(1+(p+q)^2) &= 20 \\ (q-r)^2(1+(q+r)^2) &= 21\end{aligned}$$

Find  $(r-p)^2(1+(r+p)^2)$ .

*Solution.*

Consider the three points  $P = (p, p^2)$ ,  $Q = (q, q^2)$ ,  $R = (r, r^2)$ . Then  $\overrightarrow{PQ}$  has slope  $p+q$  and  $\overrightarrow{QR}$  has slope  $q+r$ , so  $\overrightarrow{PQ} \perp \overrightarrow{QR}$ . Then, using distance formula, we have  $PQ^2 = (p-q)^2 + (p^2 - q^2)^2 = 20$  and likewise for  $QR$ , so  $(r-p)^2(1+(r+p)^2)^2 = RP^2 = PQ^2 + QR^2$  from Pythagorean Theorem. Our answer is 41.  $\square$

## §16.1.6 Really Weird NT

Source: National High School Math League (China) 2004 Test 2 Question 3, altered greatly

Proposer: watanuki-taiga-2#1569 (769480637245554688)

Problem ID: 223

Date: 2021-07-03

Difficulty: Challenging

Let  $m$  and  $n \geq 4$  be positive integers. If any  $k$ -element subset of  $\{m, m+1, \dots, m+n-1\}$  has at least 3 mutually coprime elements, and the minimum value of  $k$  is 2021, calculate the sum of all possible values of  $n$ .

*Solution.*

Let  $f(n)$  defined for integers  $n \geq 4$  be the minimum value required such that any  $f(n)$ -element subset of  $\{m, m+1, \dots, m+n-1\}$  has at least 3 mutually coprime elements. We wish to solve for  $f(n) = 2021$ . First, it is trivial that the number of elements in  $\{2, 3, 4, \dots, n+1\}$  which are either a multiple of 2 or 3 is given by  $\lfloor \frac{n+1}{2} \rfloor + \lfloor \frac{n+1}{3} \rfloor - \lfloor \frac{n+1}{6} \rfloor$ , where  $\lfloor x \rfloor$  is the greatest integer less than or equal to  $x$ . Therefore  $f(n) \geq \lfloor \frac{n+1}{2} \rfloor + \lfloor \frac{n+1}{3} \rfloor - \lfloor \frac{n+1}{6} \rfloor + 1$ . We show that the RHS is indeed equal to  $f(n)$ .

We will show that  $f(4) = 4$ ,  $f(5) = 5$ ,  $f(6) = 5$ ,  $f(7) = 6$ ,  $f(8) = 7$ ,  $f(9) = 8$  and then proceed with induction.

Step 1. When  $n = 4$ , let the set be  $\{m, m+1, m+2, m+3\}$ . One of the triples  $(m, m+1, m+2)$  and  $(m+1, m+2, m+3)$  must all be mutually coprime. Therefore it is true for  $n = 4$ . It follows that this is also true for  $n = 5$ .

When  $n = 6$ , let the set be  $\{m, m+1, m+2, m+3, m+4, m+5\}$ . Let us choose three numbers from  $x_1, x_2, x_3, x_4, x_5$ . If three of them are odd, then they are mutually coprime. Else there are three even numbers, let them be  $x_1, x_2$  and  $x_3$ . Since  $|x_i - x_j| \in \{2, 4\}$  ( $1 \leq i < j \leq 3$ ), therefore at most one is divisible by 3, at most one is divisible by 5, so there must exist one that is neither divisible by 3 nor 5, WLOG let it be  $x_3$ . Then  $x_3, x_4, x_5$  are coprime. It follows that this is true for  $n = 7$ ,  $n = 8$  and  $n = 9$ .

Step 2. If the proposition is true for  $n = t$  ( $t \geq 4$ ), then when  $n = t+6$ , we have  $f(t+6) - f(t) = 4$ . Note that  $\{m, m+1, \dots, m+t+5\} = \{m, m+1, m+2, m+3, m+4, m+5\} \cup \{m+6, m+7, \dots, m+t+5\}$ . We choose  $f(t+6) = f(t) + 4$  elements from it.

If we choose at least 5 elements from  $\{m, m+1, \dots, m+5\}$ , then from the conclusion of  $n = 6$ , we are done. Otherwise we must have chosen  $f(t)$  elements from  $\{m+6, m+7, \dots, m+t+5\}$ , which is true from the induction hypothesis.

Therefore the proposition is true for  $n = t+6$ .

Therefore,  $f(n) = \lfloor \frac{n+1}{2} \rfloor + \lfloor \frac{n+1}{3} \rfloor - \lfloor \frac{n+1}{6} \rfloor + 1$ . Note that this can be rewritten as:

$$f(n) = \begin{cases} 4t+1, & (n = 6t, t \in \mathbb{Z}^+), \\ 4t+2, & (n = 6t+1, t \in \mathbb{Z}^+), \\ 4t+3, & (n = 6t+2, t \in \mathbb{Z}^+), \\ 4t+4, & (n = 6t+3, t \in \mathbb{Z}^+), \\ 4t+4, & (n = 6t+4, t \in \mathbb{Z}^+), \\ 4t+5, & (n = 6t+5, t \in \mathbb{Z}^+) \end{cases}$$

Finally we solve  $f(n) = 2021$  to get  $n = 3029$  or  $3030$ . Therefore the answer is  $3029 + 3030 = \boxed{6059}$ .  $\square$

**§16.2 Week 2**