

OpenSourceEconomics

Presentation: Numerical Toolbox Integration

Learn, build, share, repeat

September 13, 2019

Visit us on



GitHub

Get material



for this presentation

Keane and Wolpin in 1994

“However, for problems of the size we would like to consider, a [...] simulation strategy is not computationally practicable.”

Keane and Wolpin (1994): Model Outline

Setting: In each of T periods an individual chooses between $K = 4$ mutually exclusive alternatives: occupation one / two, schooling, and home.

Each of the alternatives k is associated with a *per-period reward function*

$$R_k(t, s_t, x_{1t}, x_{2t}; \theta, \epsilon_{kt}) \quad \text{for } k = 1, \dots, 4.$$

Keane and Wolpin (1994): Model Outline

With $\epsilon_t \sim \mathcal{N}_4(\mu, \Sigma)$, ϵ_{kt} are alternative-specific shocks to

- ▶ skill level ($k = 1, 2$)
- ▶ consumption value of schooling ($k = 3$)
- ▶ value of non-market time ($k = 4$),

Cross-correlations admissible, but not serial correlation.

Keane and Wolpin (1994): Model Outline

With $\epsilon_t \sim \mathcal{N}_4(\mu, \Sigma)$, ϵ_{kt} are alternative-specific shocks to

- ▶ skill level ($k = 1, 2$)
- ▶ consumption value of schooling ($k = 3$)
- ▶ value of non-market time ($k = 4$),

Cross-correlations admissible, but not serial correlation.

The vector θ summarizes parameters for wage, schooling, and non-market time.

Question of interest: Does a school subsidy change the average years of schooling in the sample?

Keane and Wolpin (1994): Solution

Individual's objective function, alternative-specific value function at $t = 0, \dots, T$:

$$V_k(S(t), t) = \mathbb{E} \left[\sum_{\tau=t}^T \delta^{\tau-t} \sum_k^K R_k(\tau) d_k(\tau) \mid S(t) \right] , T < \infty.$$

Keane and Wolpin (1994): Solution

Individual's objective function, alternative-specific value function at $t = 0, \dots, T$:

$$V_k(S(t), t) = \mathbb{E} \left[\sum_{\tau=t}^T \delta^{\tau-t} \sum_k^K R_k(\tau) d_k(\tau) \mid S(t) \right] , T < \infty.$$

State Space $S(t)$: All factors, known to the individual, that affect *current* rewards or the probability distribution of any of the *future* rewards.

Keane and Wolpin (1994): Solution

Maximizing Choice: Optimal sequence of control variables $\{d_k(t)\}_{k \in K}$ for $t = 0, \dots, T$

Value function for period t :

$$V(S(t), t) = \max_{\{d_k(t)\}_{k \in K}} \{V_k(S(t), t)\},$$

Keane and Wolpin (1994): Solution

Maximizing Choice: Optimal sequence of control variables $\{d_k(t)\}_{k \in K}$ for $t = 0, \dots, T$

Value function for period t :

$$V(S(t), t) = \max_{\{d_k(t)\}_{k \in K}} \{V_k(S(t), t)\},$$

where *alternative-specific value functions* $V_k(S(t), t)$

$$V_k(S(t), t) = R_k(S(t), t) + \delta \mathbb{E}[V(S(t+1), t+1)] \quad , t \leq T-1$$

$$V_k(S(T), T) = R_k(S(T), T)$$

obey the Bellman equation (Bellman, 1957).

Keane and Wolpin (1994): Solution

Individual at $(T - 1)$: For each k , calculate

$$\begin{aligned} & \text{Emax}(R_1(T), R_2(T), R_3(T), R_4(T) \mid \bar{S}(T-1), d_k(T-1)) \\ &= \int_{\epsilon_T} \max(R(T) \mid \bar{S}(T-1), d_k(T-1)) f(\epsilon_T) d\epsilon_T \end{aligned}$$

Keane and Wolpin (1994): Solution

Individual at $(T-1)$: For each k , calculate

$$\begin{aligned} & \text{E} \max(R_1(T), R_2(T), R_3(T), R_4(T) \mid \bar{S}(T-1), d_k(T-1)) \\ &= \int_{\epsilon_T} \max(R(T) \mid \bar{S}(T-1), d_k(T-1)) f(\epsilon_T) d\epsilon_T \end{aligned}$$

$\Rightarrow V_k(S(T-1), T-1)$ are known

Keane and Wolpin (1994): Solution

Individual at $(T-1)$: For each k , calculate

$$\begin{aligned} & \text{E} \max(R_1(T), R_2(T), R_3(T), R_4(T) \mid \bar{S}(T-1), d_k(T-1)) \\ &= \int_{\epsilon_T} \max(R(T) \mid \bar{S}(T-1), d_k(T-1)) f(\epsilon_T) d\epsilon_T \end{aligned}$$

$\Rightarrow V_k(S(T-1), T-1)$ are known

→ Receive random draw of $\epsilon_{T-1} = (\epsilon_{1,T-1}, \dots, \epsilon_{4,T-1})$

→ Choose alternative k with highest value

Keane and Wolpin (1994): Solution

Individual at $t < (T - 1)$: At every t , calculate

$$\text{Emax}[V_1(S(t+1), t+1), \dots, V_4(S(t+1), t+1) \mid \bar{S}(t), d_k(t)]$$

Keane and Wolpin (1994): Solution

Individual at $t < (T - 1)$: At every t , calculate

$$\text{Emax}[V_1(S(t+1), t+1), \dots, V_4(S(t+1), t+1) \mid \bar{S}(t), d_k(t)]$$

Requirement: Alternative-specific value functions at t must have been calculated for *all* of the possible state space values at $t + 1$

⇒ Need to calculate the $V_k(\cdot)$ at each future dates,
and at all feasible state points.

⇒ Computational complexity

Keane and Wolpin (1994): Estimation

Exact solution requires to solve multiple (aka millions) K -dimensional integrals (aka $E_{\max}(\cdot)$).

A priori unknown: extent of finite sample bias as accuracy of $E_{\max}(\cdot)$ varies.

Keane and Wolpin (1994): Estimation

Exact solution requires to solve multiple (aka millions) K -dimensional integrals (aka $E_{\max}(\cdot)$).

A priori unknown: extent of finite sample bias as accuracy of $E_{\max}(\cdot)$ varies.

Our goal: Use quasi-Monte Carlo approach to lower required points for given accuracy

⇒ Use respy for a simulation study

Tool of choice: respy

respy 1.2.1 documentation »

Table Of Contents

Getting Started

Economics

Software

Development

API

Additional Information

Search

Go

Enter search terms or a module, class or function name.

Welcome to respy's documentation!

[PyPI](#) | [GitHub](#) | [Issues](#) | [Pull Requests](#)

respy is an open-source Python package for the simulation and estimation of a prototypical finite-horizon discrete choice dynamic programming model. We build on the baseline model presented in:

Keane, M. P. and Wolpin, K. I. (1994). [The Solution and Estimation of Discrete Choice Dynamic Programming Models by Simulation and Interpolation: Monte Carlo Evidence](#). *The Review of Economics and Statistics*, 76(4): 648-672.

license **MIT**

- Getting Started
 - Introduction
 - Installation
 - Tutorial - Model
 - Tutorial - Keane and Wolpin (1994) - Simulation
 - Replicating Keane and Wolpin (1997)
- Economics
 - The Economic Model
 - Solution and Estimation
- Software
 - Model Specification
 - Numerical Methods
 - Reliability
 - Scalability
 - Software Engineering
- Development
 - Docker

New option in respy

Implementation of *low-discrepancy sequences for quasi-Monte Carlo simulation*:

- ▶ Halton ((co)prime numbers)
- ▶ Sobol (primitive polynomial)
- ▶ R_d -sequence (additive recurrence method based on irrational numbers)

Halton and Sobol sequences rely on implementation from [chaospy](#).

R_d implemented based on a [blog post](#) by M. Roberts.

Background R_d sequence


Intuition: Connection of generalized golden ratio (Marohnic and Strmecki (2012)) with construction of higher-dimensional low discrepancy sequences

Construction method: The parameter-free d -dimensional infinite sequence $R_d(\phi_d)$ is constructed as follows:

$$t_n = \{n\alpha\}; \quad \alpha = \left(\frac{1}{\phi_d}, \frac{1}{\phi_d^2}, \dots, \frac{1}{\phi_d^d} \right); \quad n = 1, 2, \dots,$$


and ϕ_d is the unique positive root of $x^{d+1} = x + 1$.

Simulation Study




Take $E_{\max}(\cdot)$ for one state, iterate over points

Simulation Study




Compute relative RMSE by aggregating over all state spaces

Simulation Study



Compute relative RMSE by aggregating over all state spaces

Simulation Study



Compute differences in policy effect dependent on number of points

References

Bellman, R. E. (1957). [“Dynamic Programming”](#). Princeton University Press, Princeton, NJ.

Davis, P. J. and Rabinowitz, P. (1984). [“Methods of Numerical Integration”](#). Dover Publications, Mineola, NY.

Feinberg, J. and Langtangen, H.P. (2015). [“Chaospy: An open source tool for designing methods of uncertainty quantification”](#). *Journal of Computational Science*, Vol.11, pp. 46-57.

Halton, J. (1964). [“Algorithm 247: Radical-inverse Quasi-Random Point Sequence”](#) *Communications of the ACM*, 7, pp. 701-701.

Keane, M. P. and Wolpin, K. I. (1997). [“The Career Decisions of Young Men”](#). *Journal of Political Economy*, 105(3), pp. 473-522.

Keane, M. P. and Wolpin, K. I. (1994). [“The Solution and Estimation of Discrete Choice Dynamic Programming Models by Simulation and Interpolation: Monte Carlo Evidence”](#). *The Review of Economics and Statistics*, 76(4), pp. 648-672.

References

L'Ecuyer, P. (2017). "Randomized Quasi-Monte Carlo: An Introduction for Practitioners". 12th International Conference on Monte Carlo and Quasi-Monte Carlo Methods in Scientific Computing (MCQMC 2016), August 2016, Stanford, CA.

L'Ecuyer, P. and Lemieux, C. (2005). "Recent Advances in Randomized Quasi-Monte Carlo Methods". In: Dror M., L'Ecuyer P., Szidarovszky F. (eds) Modeling Uncertainty. *International Series in Operations Research and Management Science*, Vol. 46, Springer, Boston, MA.

Lemieux, C. (2009). "Monte Carlo and Quasi-Monte Carlo Sampling". Springer, Boston, MA.

Puterman, M. L. (1994). "Markov Decision Processes: Discrete Stochastic Dynamic Programming (1st ed.)". John Wiley & Sons, Inc., New York, NY, USA.

References

Roberts, M. (2019). "The Unreasonable Effectiveness of Quasirandom Sequences". [Extreme Learning](#).

Retrieved on 24th August, 2019 from <http://extremelearning.com.au/unreasonable-effectiveness-of-quasirandom-sequences/>.

Skrainka, B. S., and K. L. Judd. (2011). "High Performance Quadrature Rules: How Numerical Integration Affects a Popular Model of Product Differentiation". Unpublished.

Sobol, I. M. (1967). "Distribution of Points in a Cube and Approximate Evaluation of Integrals". *Zh. Vych. Mat. Mat. Fiz.*, 7, pp. 784-702 (in Russian), *U.S.S.R Comput. Maths. Math. Phys.* 7: pp. 86-112.

Wolpin, K. I. (2013). "The Limits of Inference without Theory".. MIT Press, Cambridge, MA.

*Thank you for your attention
and enjoy the workshop!*

Contact




Philipp Eisenhauer
peisenha

Philipp Eisenhauer

Mail eisenhauer@policy-lab.org

Web <http://eisenhauer.io>

Repository <https://github.com/peisenha>



Rafael Suchy

Rafael Suchy
rafaelsuchy

Mail rafael.suchy@uni-bonn.de

Repository <https://github.com/rafaelsuchy>

Appendix