

Physicochemical and metabolic evaluation of nine compounds

MMV_OSDD**Report #: CDCO_MMV_OSDD_18_001****25 May 2018****Quality Statement:**

This non-GLP study was conducted using established techniques in accordance with the relevant guidelines and standard operating procedures (SOPs) of the Centre for Drug Candidate Optimisation, Monash University. This report accurately reflects the raw data obtained during the performance of this study.

Director:	Susan Charman, PhD
Project Coordinator:	Karen White, PhD
Section Leader(s):	Michael Campbell, PhD Francis Chiu, PhD Alison Thistlethwaite, PhD
Study contributor(s):	Scott Blundell Caithlin White Helena Barker
Study number(s):	MMV_OSDD_18_001

A. Experimental Methods

a) Calculated physicochemical parameters using ChemAxon JChem software

A range of physicochemical properties evaluating drug-likeness and likely oral absorption characteristics were calculated using the ChemAxon chemistry cartridge via JChem for Excel software (version 16.4.11). A brief description of each parameter is provided below, along with a suggested ideal range based on research reported in the ADME literature from key industry and academic sources.

MW (< 500): Molecular Weight.

PSA_{pH 7.4} (< 140 Å²): Polar surface area also inversely correlates with membrane permeability.

HBD (< 5) & HBA (< 10): Number of hydrogen bond donors and acceptors gives an indication of the hydrogen bonding capacity, which is inversely related to membrane permeability.

FRB (≤ 10): Number of freely rotating bonds represents the flexibility of a molecule's conformation.

Arom. Rings (< 4): Total number of aromatic and heteroaromatic rings is also related to molecular flexibility.

Fsp³ (> 0.3): Fraction of sp³ carbons to total carbons indicates the complexity of a molecule's 3D structure.

cpKa: Ionisation constants impact solubility and permeability. Only physiologically relevant predicted values are provided here (*i.e.* 0 < pKa < 12).

cLogP/cLogD_{pH} (< 5): Partition coefficients reflect the lipophilic character of the neutral structure, while distribution coefficients reflect the partitioning properties of the ionised molecule at a specific pH.

b) Kinetic Solubility Estimation using Nephelometry (Sol_{pH})

Compound in DMSO was spiked into either pH 6.5 phosphate buffer or 0.01 M HCl (approx pH 2.0) with the final DMSO concentration being 1%. After 30 minutes had elapsed, samples were then analysed via Nephelometry to determine a solubility range. See Bevan and Lloyd (2000) *Anal Chem*, 72:1781-1787.

c) Distribution Coefficient Estimation using Chromatography (gLogD_{pH})

Partition coefficient values (LogD) of the test compounds were estimated at pH 7.4 by correlation of their chromatographic retention properties against the characteristics of a series of standard compounds with known partition coefficient values. The method employed is a gradient HPLC based derivation of the method developed by Lombardo. See Lombardo *et al.* (2001) *J Med Chem*, 44:2490-2497.

d) In Vitro Metabolic Stability

Incubation:

The metabolic stability assay was performed by incubating each test compound in liver microsomes at 37°C and a protein concentration of 0.4 mg/mL. The metabolic reaction was initiated by the addition of an NADPH-regenerating system and quenched at various time points over a 60 minute incubation period by the addition of acetonitrile containing diazepam as internal standard. Control samples (containing no NADPH) were included (and quenched at 2, 30 and 60 minutes) to monitor for potential degradation in the absence of cofactor.

The human liver microsomes used in this experiment were supplied by XenoTech, lot # 1410230. The mouse liver microsomes used in this experiment were supplied by XenoTech, lot # 1510256. Microsomal incubations were performed at a substrate concentration of 0.5 µM.










Data analysis:

Species scaling factors from Ring *et al.* (2011) *J Pharm Sci*, 100:4090-4110 were used to convert the *in vitro* CL_{int} (µL/min/mg) to an *in vivo* CL_{int} (mL/min/kg). Hepatic blood clearance and the corresponding hepatic extraction ratio (E_H) were calculated using the well stirred model of hepatic extraction in each species, according to the "*in vitro* T_{1/2}" approach described in Obach (1999) *Drug Metab. Dispos.* 27: 1350-1359. The E_H was then used to classify compounds as low (< 0.3), intermediate (0.3 - 0.7), high (0.7 - 0.95) or very high (> 0.95) extraction compounds. Predicted *in vivo* clearance values have not been corrected for microsomal or plasma protein binding. Species scaling calculations are based on two assumptions: 1) NADPH-dependent oxidative metabolism predominates over other metabolic routes (*i.e.* direct conjugative metabolism, reduction, hydrolysis, *etc.*), and; 2) rates of metabolism and enzyme activities *in vitro* are truly reflective of those that exist *in vivo*. If significant non-NADPH-mediated degradation is observed in microsome control samples, then assumption (1) is invalid and predicted clearance parameters are therefore not reported.

B. Results

Experimental results are tabulated below.










Table 1: Physicochemical evaluation of nine compounds

Compound (Batch)	Structure	MW	PSA (Å ²)	FRB	HBD	HBA	Arom. Rings	Fsp ³	Predicted pKa (0 - 12 only)	cLogP	cLogD at pH 7.4	gLogD at pH 7.4	Sol _{2.0} (µg/mL)	Sol _{6.5} (µg/mL)	Notes
MMV025100 (MNB6-7)		306.36	112.0	2	2	5	3	0.00	Basic: 3.8 Acidic: 10.1	1.3	1.3	1.4	50 - 100	6.3 - 12.5	
MMV639565 (AEW 302-1)		386.79	52.3	5	0	4	4	0.11	Basic: 1.2 Acidic: none	3.4	3.4	3.8	1.6 - 3.1	3.1 - 6.3	1
MMV669784 (EGT 302-1)		388.76	61.5	5	0	4	4	0.06	Basic: 1.2 Acidic: none	3.6	3.6	3.7	6.3 - 12.5	6.3 - 12.5	
MMV670246 (EGT 95-3)		415.78	81.4	5	1	5	4	0.05	Basic: 1.2 Acidic: none	3.0	3.0	3.1	3.1 - 6.3	1.6 - 3.1	1
MMV693155 (AEW 317-1, AEW 294-2)		412.4	81.8	8	1	6	4	0.19	Basic: 1.2 Acidic: none	2.3	2.3	2.8	25 - 50	12.5 - 25	1, 2
MMV897700 (EGT 65-1)		408.41	61.5	7	0	5	3	0.50	Basic: 1.2 Acidic: none	1.6	1.6	4.3	1.6 - 3.1	< 1.6	
MMV1576784 (EGT 257-1)		448.47	61.5	7	0	5	3	0.19	Basic: 1.2 Acidic: none	7.8	7.8	4.4	1.6 - 3.1	< 1.6	
MMV1579336 (EGT 199-5)		426.38	101.7	8	1	7	4	0.14	Basic: 1.2 Acidic: 3.7	2.7	-0.6	1.3	6.3 - 12.5	25 - 50	
MMV1579341 (MK035-1)		391.38	68.1	5	1	4	5	0.10	Basic: 1.2 Acidic: none	2.9	2.9	3.4	3.1 - 6.3	3.1 - 6.3	

Notes

- 1 - Results for this compound were originally reported in CDCO_MMV_OSDD_17_001.
- 2 - A second chromatographic peak was visible in the gLogD experiment. The reported gLogD value is calculated from the retention time of the major peak (70% by peak area).

Table 2: Metabolic evaluation of nine compounds

Compound (Batch)	Structure	Microsome Species	T _{1/2} (min)	CL _{int, in vitro} (μL/min/mg protein)	Predicted CL _{int, in vivo} (mL/min/kg)	Predicted CL _{blood} (mL/min/kg)	Predicted E _H	Clearance classification	Notes
MMV025100 (MNB6-7)		Human	> 255	< 7	< 6	< 5	< 0.22	low	1
		Mouse	23	76	196	74	0.62	intermediate	
MMV639565 (AEW 302-1)		Human	53	33	27	12	0.57	intermediate	2
		Mouse	9	193	497	97	0.81	high	
MMV669784 (EGT 302-1)		Human	> 255	< 7	< 6	< 5	< 0.22	low	1
		Mouse	46	37	96	53	0.45	intermediate	
MMV670246 (EGT 95-3)		Human	164	11	9	6	0.30	low	2
		Mouse	115	15	39	29	0.24	low	
MMV693155 (AEW 317-1, AEW 294-2)		Human	35	49	41	14	0.66	intermediate	2
		Mouse	5	361	930	106	0.89	high	
MMV897700 (EGT 65-1)		Human	9	197	162	18	0.89	high	3
		Mouse	3	573	1479	111	0.92	high	
MMV1576784 (EGT 257-1)		Human	7	249	204	19	0.91	high	3
		Mouse	< 2	> 866	> 2235	> 113	> 0.94	very high	
MMV1579336 (EGT 199-5)		Human	> 255	< 7	< 6	< 5	< 0.22	low	4
		Mouse	> 255	< 7	< 18	< 16	< 0.13	low	
MMV1579341 (MK035-1)		Human	8	205	168	18	0.89	high	
		Mouse	4	417	1076	108	0.90	high	

Notes

- 1 - This compound showed minimal microsomal degradation (<15%) in human metabolism samples over the course of the incubation.
- 2 - Results for this compound were originally reported in CDCO_MMV_OSDD_17_001.
- 3 - Calculated mouse metabolism parameters are based on the first 2 time-points (i.e. 2 & 5 minutes) due to rapid degradation, and are therefore an estimate only.
- 4 - This compound showed minimal microsomal degradation (<15%) in human and mouse metabolism samples over the course of the incubation.