

P. falciparum PfATP4 Multi-Drug Resistance Resistance to KAE609 (Cipargamin) is Present in Africa*

James McCulloch[†]

March 31, 2018

Abstract

The PfATP4 (PF3D7.1211900) multi-drug resistance mutation **G223R** is found in Africa by genetically analyzing 2640 worldwide Plasmodium falciparum blood stage isolates (the MalariaGen Pf3k resource). This mutation confers an approximate 8 fold [4] increase in the PfATP4 IC_{50} of Spiroindolones (KAE609 & KAE678) [14],[16],[4],[10] and Aminopyrazoles (GNF-Pf4492) [4]. It is postulated that the **G223R** mutation may be a consequence of the drug resistant Southeast Asian Dd2 genotype becoming more dominant in Africa [3]. The presence of this mutation has important policy implications for the eventual general deployment of the Spiroindolone KAE609 (Cipargamin) which is currently undergoing stage 2 clinical trials.

Keywords: Falciparum Spiroindolones Cipargamin KAE609 G223R Drug Resistance.

Introduction

Advances in laboratory automation, process control software and automated biological detectors have allowed malaria researchers to perform high-throughput screening (HTS) of very large chemical libraries [6] [4] to find candidate antimalarial drugs. This HTS screening has been successful in identifying novel anti-malarial drugs including KAE609 [6]. Screens often use whole-cell assays to determine likely candidates. A straightforward method of determining the location and action of the HTS drug candidates is to pressure Plasmodium falciparum (Pf) laboratory strains and map the resultant genetic mutations that confer resistance.

Many resistance conferring mutations of novel HTS candidates have mapped to the transmembrane ATPase sodium pump [15][16] PfATP4 (PF3D7.1211900) indicating that the likely HTS candidate action is the inhibition of this protein. A literature search reveals 54 mutations in PfATP4 conferring

*©James McCulloch 2018. Permission is granted to copy, distribute and quote this document.

[†]email: james.duncan.mcculloch@gmail.com

resistance. These are tabulated in table 4 for the convenience of the reader. Many of the mutations confer resistance to multiple HTS candidates from different compound classes [16].

The MalariaGen consortium Malaria 3000 (Pf3k) project has processed 2640 genetically sequenced blood stage Pf isolates (128 laboratory, 2512 field) using GATK¹ best practices into Variant Calling File (VCF) format. This wonderful genetic resource removes the need for the considerable technical and computing infrastructure needed to go from raw short read FASTQ sequence files, via sorted BAM files, to properly generated VCF files². An immediate use of the Pf3k resource is to check for the existence of drug resistance conferring mutations in geographically distinct Pf populations. This was done by Gomes et al [5] for a number of drug target proteins including PfATP4 using *Delly* software (Rausch et al [13]) to analyze Pf3k BAM files and generate variants³.

The results tabulated by Gomes et al [5] are different to the results of this research. In particular, Gomes et al [5] do not report the PfATP4 synonymous SNP **G223G** (C669A, 35% of all isolates) characteristic of the Pf strain Dd2 [14] or **G223S** (G667A, 8.9% of all isolates) present in African populations. It should be noted that the G667A and C669A SNPs (GGC→AGA) occurring together produce the **G223R** resistance mutation. The probable reason for the difference between Gomes et al [5] and this research appears to be different variant calling methodology. Whereas Gomes et al [5] use *Delly* (Rausch et al [13]) to call variants, this research does not attempt to independently discover variants, but uses the Pf3k VCF files generated by MalariaGen using GATK. The VCF files are then post-processed by `kg1_genome` software developed for this and subsequent Pf genetic research by the author (details in the Methods and Data appendix).

PfATP4 Mutations in Model Pf Organisms

The MalariaGen Pf3k whole genome sequence (WGS) dataset includes 5 laboratory model Pf reference strains and 121 mixtures and genetic crosses of these model strains (see the MalariaGen ftp website⁴ for more details). Consistent with standard practice, the 3D7⁵ strain is the genetic reference wild type (WT) for this analysis and the MalariaGen BAM/VCF Pf3k generation. The Single Nucleotide Polymorphism (SNP) mutations from these model strains are tabulated below (table 1). The SNP structure of the 121 Pf crosses and mixtures of the 5 model strains is consistent

¹The Broad Institute Genome Analysis Toolkit (GATK).

²For more information see the methods and data appendix

³*Delly* outputs variants as BCF files but these can be readily converted to VCF format files.

⁴ftp://ngs.sanger.ac.uk/production/pf3k/release_5/

⁵*P. falciparum* strain 3D7 was derived from strain NF54 from a patient living in the Netherlands.

with the pattern of crosses and mixtures of the 5 model strains documented⁶ by MalariaGen. However, the pattern of the SNPs in all isolates (128 laboratory, 2512 field) reflects the geographical bias of the Pf3k field isolates which are only sourced from Southeast Asia and Africa with no South American examples.

Table 1: Pf3k Laboratory Strain PfATP4 Mutations

Pf Strain	Location	E30G	S86N	G223G	Q1081K	G1128R	I1237V
3D7 (WT)	Netherlands						
7G8 (2)	Brazil	2	2		2	2	2
Dd2 (1)	Cambodia			1		1	
GB4 (1)	Ghana				1	1	
HB3 (2)	Honduras						
Mixtures & Crosses (121)		33	36	30	55	82	34
% of 2640 isolates		1.3%	1.5%	35.1%	36.6%	56.7%	3.4%

The Geographic Distribution of G223R

The geography of the **G223R** mutation is tabulated below in table 2. The table gives the proportion of the three mutually exclusive mutations of codon 223 in PfATP4 and the synonymous mutation **A229A** (T687A). The Southeast Asian locations that do not have **G223S** or **G223R** (G667A) mutations are not tabulated. The synonymous mutation **G223G** (C669A) is characteristic of the Dd2 model organism [14] and is, as expected, prevalent in Southeast Asia. In particular it is present in 72% of the Ramu, Bangladesh isolates tabulated below. The presence of C669A brings the Pf organism to within 1 SNP of **G223R** with the subsequent addition of G667A. However, only populations with the **G223S** (G667A) mutation have also acquired the **G223R** mutation, strongly suggesting that this mutation is acquired by the subsequent addition of C669A to an *existing* G667A genotype.

The **G223R** mutation may be a consequence of the drug resistant Southeast Asian Dd2 genotype becoming more dominant in Africa [3] with this gene flow adding the Dd2 characteristic mutation **G223G** (C669A) to *existing* **G223S** (G667A) African sub-populations.

We also find very high linkage dis-equilibrium between G667A and T687A (**A229A**). This dis-equilibrium is tabulated as a Pearson correlation (correlations with $p > 0.01$ are not shown). This may be an artifact of an ancestor genotype that has not decayed to linkage equilibrium through recombination because of the proximity of the SNPs. However, there is no similar magnitude of linkage dis-equilibrium with other nearby SNPs such as C669A.

⁶See the relevant Pf3K documentation at MalariaGen at <https://www.malariagen.net/>.

Table 2: Geographic Locations of G223S and G223R Mutations

Location	Isolates	G223G C669A	G223R G667A+C669A	G223S G667A	A229A T687A	Linkage DisEq. G667A*T687A
Nigeria, Ilorin	5	0% (0)	20% (1)	0% (0)	0% (0)	
Guinea, Nzerekore	100	14% (14)	6% (6)	16% (16)	23% (23)	0.94
Ghana, Kintampo	68	12% (8)	6% (4)	24% (16)	24% (16)	0.86
Ghana, Navrongo	549	11% (59)	6% (32)	22% (119)	23% (129)	0.89
Mali, Kollo	51	22% (11)	4% (2)	25% (13)	24% (12)	0.86
Gambia, Brikama	65	15% (10)	3% (2)	12% (8)	11% (7)	0.81
Mali, Faladje	36	14% (5)	3% (1)	11% (4)	8% (3)	
CongoDR, Kinshasa	113	18% (20)	2% (2)	13% (15)	12% (14)	0.89
Senegal, Thies	133	16% (21)	1% (1)	12% (16)	9% (12)	0.82
Malawi, Chikwawa	317	19% (61)	1% (2)	7% (23)	7% (23)	0.96
Mali, Bandiagara	9	22% (2)	0% (0)	33% (3)	22% (2)	
Bangladesh, Ramu	50	72% (36)	0% (0)	2% (1)	2% (1)	
Malawi, Zomba	52	10% (5)	0% (0)	2% (1)	2% (1)	

G223R and Multi-Drug Resistance

Rottmann et al [14] find the **G223R** mutation in one of the Dd2 strains (NITD678-RDd2 clone#1) they pressured with Spiroindolone KAE678 and speculate that the **G223R** mutation may *not* be a result of evolved resistance to KAE678. Manary et al [10] pressured Dd2 strains with KAE678 for 2-4 months and found the **G223R** mutation in the resultant drug resistant strain. Flannery et al [4] show an approximate 8 fold increase in the IC_{50} of GNF-Pf4492, KAE609 and KAE678 for G223R mutated Dd2. Spillman and Kirk [16] show that the NITD678-RDd2 clone#1 strain developed by Rottmann et al [14] has an KAE678 EC_{50} of 193 ± 39 nM compared to 21.9 ± 1.2 nM for the Dd2 wildtype. This is an approximate 9 fold increase and is consistent with results found by Flannery et al [4]. Finally, Jiménez-Díaz et al [7] mention **G223R** as a drug resistant mutation for the Dihydroisoquinolone SJ733 but provide no further detail.

The presence of the **G223R** mutation at moderate but significant levels in Africa has important policy implications for the development and deployment of the new generation of anti-malarial drugs which are active against the ATPase sodium pump PfATP4. One of the most promising of these anti-malarials, KAE609 (Cipargamin) is currently (March 2018) undergoing stage 2 clinical trials. The possibility that the **G223R** mutation may be a result of the drug resistant Southeast Asian Dd2 genotype becoming more dominant in Africa [3] and the very high linkage dis-equilibrium between the SNPs G667A and T687A are interesting topics that merit further research.

References

- [1] V. Corey, A. Lukens, E. Istvan, M. Lee, V. Franco, P. Magistrado, O. Coburn-Flynn, T. Sakata-Kato, O. Fuchs, N. Gndig, G. Goldgof, M. Linares, M. Gomez-Lorenzo, C. De CAzar, M. Lafuente-Monasterio, S. Prats, S. Meister, O. Tanaseichuk, M. Wree, Y. Zhou, P. Willis, F. Gamo, D. Goldberg, D. Fidock, D. Wirth, and E. Winzeler. A broad analysis of resistance development in the malaria parasite. *Nature Communications*, 7, 2016.
- [2] A. Dring, D. Weese, T. Rausch, and K. Reinert. SeqAn an efficient, generic C++ library for sequence analysis. *BMC Bioinformatics*, 9(11), 2008.
- [3] A. Ecker, A. Lehane, J. Clain, and D. Fidock. PfCRT and its role in antimalarial drug resistance. *Trends in Parasitology*, 28, 2012.
- [4] E. Flannery, C. McNamara, S. Kim, and et al. Mutations in the P-Type Cation-Transporter ATPase 4, PfATP4, Mediate Resistance to Both Aminopyrazole and Spiroindolone Antimalarials. *ACS Chemical Biology*, 10(2), 2015.
- [5] A. Gomes, M. Ravenhall, E. Benavente, A. Talman, C. Sutherland, C. Roper, T. Clark, and S. Campino. Genetic diversity of next generation antimalarial targets: A baseline for drug resistance surveillance programmes. *International Journal for Parasitology Drugs and Drug Resistance*, 7, 2017.
- [6] M. Hovlid and E. Winzeler. Phenotypic screens in antimalarial drug discovery. *Trends in parasitology*, 32, 2016.
- [7] M. Jiménez-Díaz, D. Ebert, Y. Salinas, A. Pradhan, A. Lehane, M. Myrand-Lapierre, K. O’Loughlin, D. Shackelford, M. Justino de Almeida, A. Carrillo, J. Clark, A. Dennis, J. Diep, X. Deng, S. Duffy, A. Endsley, G. Fedewa, W. A. Guiguemde, M. Gómez, G. Holbrook, J. Horst, C. Kim, J. Liu, M. Lee, A. Matheny, M. Martínez, G. Miller, A. Rodríguez-Alejandre, L. Sanz, M. Sigal, N. Spillman, P. Stein, Z. Wang, F. Zhu, D. Waterson, S. Knapp, A. Shelat, V. Avery, D. Fidock, F. Gamo, S. Charman, J. Mirsalis, H. Ma, S. Ferrer, K. Kirk, I. Angulo-Barturen, D. Kyle, J. DeRisi, D. Floyd, and R. Guy. (+)-SJ733, a clinical candidate for malaria that acts through ATP4 to induce rapid host-mediated clearance of Plasmodium. *Proceedings of the National Academy of Sciences*, 111(50):E5455–E5462, 2014.
- [8] A. Lee and D. Fidock. Evidence of a Mild Mutator Phenotype in Cambodian Plasmodium falciparum Malaria Parasites. *PLoS ONE*, 11(2), 2016.
- [9] A. Lehane, M. Ridgway, E. Baker, and K. Kirk. Diverse chemotypes disrupt ion homeostasis in the malaria parasite. *Molecular Microbiology*, 94(2), 2014.
- [10] M. Manary, S. Singhakul, E. Flannery, S. Bopp, V. Corey, A. Bright, C. W. McNamara, J. R. Walker, and E. A. Winzeler. Identification of pathogen genomic variants through an integrated pipeline. *BMC Bioinformatics*, 15, 2014.
- [11] M. Pinkevych, J. Petracic, S. Berezky, I. Rooth, A. Frnert, and M. Davenport. Understanding the Relationship Between Plasmodium falciparum Growth Rate and Multiplicity of Infection. *The Journal of Infectious Diseases*, 211, 2015.
- [12] R. Poplin, V. Ruano-Rubio, M. DePristo, T. Fennell, M. Carneiro, G. Van der Auwera, D. Kling, L. Gauthier, A. Levy-Moonshine, D. Roazen, K. Shakir, J. Thibault, S. Chandran, C. Whelan, M. Lek, S. Gabriel, M. Daly, B. Neale, D. MacArthur, and E. Banks. Scaling accurate genetic variant discovery to tens of thousands of samples. *bioRxiv*, 2017.
- [13] T. Rausch, T. Zichner, A. Schlattl, A. Stütz, V. Benes, and J. Korbel. DELLY: structural variant discovery by integrated paired-end and split-read analysis. *Bioinformatics*, 28, 2012.
- [14] M. Rottmann, C. McNamara, B. Yeung, M. Lee, B. Zou, B. Russell, and et al. Spiroindolones, a potent compound class for the treatment of malaria. *Science*, 329, 2010.
- [15] N. Spillman, R. Allen, C. McNamara, B. Yeung, E. Winzeler, T. Diagana, and K. Kirk. Na⁺ Regulation in the Malaria Parasite Plasmodium falciparum Involves the Cation ATPase PfATP4 and Is a Target of the Spiroindolone Antimalarials. *Cell Host & Microbe*, 13, 2013.

- [16] N. Spillman and K. Kirk. The malaria parasite cation ATPase PfATP4 and its role in the mechanism of action of a new arsenal of antimalarial drugs. *International Journal for Parasitology Drugs and Drug Resistance*, 2, 2015.

Methods and Data Appendix

This research uses genetically sequenced *Plasmodium falciparum* isolates from the MalariaGen Pf3K project (release 5, 9 Feb 2016). MalariaGen selected 2640 field (2512) and laboratory (128) isolates from publically available FASTQ short read sequence files. A spreadsheet with accession numbers, location and other useful sample information is available at the ftp website ftp://ngs.sanger.ac.uk/production/pf3k/release_5/. The selected isolates are from Southeast Asia and Africa. MalariaGen processed these isolates into aligned sorted and filtered BAM (compressed SAM) files. The prepared BAM files are then processed by the Broad Institute GATK [12] variant caller to generate VCF files. In order to account for the frequent Multiplicity of Infection (MOI) [11] in Pf blood samples, MalariaGen has assumed a diploid genotype when generating the VCF files.

These VCF files are then post-processed using a C++ software library `kg1_genome` developed for this and subsequent Pf genetic research by the author. This software is available at https://github.com/kellerberrin/KGL_Gene under an MIT license. The associated documentation on GitHub shows how to compile and use this library; the library also has a built-in help function. The `kg1_genome` library uses third party libraries (documented on GitHub) notably the Seqan [2] genome sequence library for advanced sequence comparison and manipulation. Finally, output from `kg1_genome` is examined using tools that will be familiar to readers such as `Excel` and `R` to produce the final SNP list (no indels were detected in PfATP4). The FASTA and GFF files used by `kg1_genome` were obtained from Plasmodb (<http://plasmodb.org>) release 35 (1 December 2017).

The identification of SNPs and other variants is a trade-off between sensitivity and specificity; the `kg1_genome` software parses the MalariaGen VCF files and filters variants using the following quality metrics: (1) the Phred-scaled probability that a REF/ALT polymorphism exists, $QUAL \geq 100$; (2) the log of the ratio of the probability that a called variant is true over the probability that the called variant is false ($\log [prob(TP)/prob(FP)]$), $VQSLOD^7 \geq 2$. For each filtered variant, each of the 2640 isolates are individually filtered for depth of coverage, $DP \geq 10$ and the Phred-scaled confidence that the isolate genotype assignment is correct, $GQ \geq 20$.

After parsing and filtering the VCF file the `kg1_genome` software converts the filtered isolate variant information into an internal variant format and, for each isolate, mutates the reference PfATP4 DNA sequence with the filtered variants obtained for each isolate. This reconstructed mutated DNA

⁷For a discussion of the VQS, VQSLOD statistics see <https://gatkforums.broadinstitute.org/gatk/discussion/39/variant-quality-score-recalibration-vqsr>.

sequence is then compared to the reference DNA sequence using the Seqan [2] genome sequence library and a list of variants is generated from the results of this comparison. This indirect method of extracting isolate variant information has the advantage of examining reconstructed mutated DNA using the familiar Needleman Wunsch algorithm and does not assume VCF called variants are independent or mutually exclusive.

This process generates 188 distinct SNP variants from the 2640 isolates. The majority of the distinct SNPs, 110, are found in only 1 or 2 isolates. There are 22 SNPs that are found in $\geq 1\%$ of all isolates and these are tabulated below in table 3.

Table 3: All PfATP4 SNPs $\geq 1\%$ frequency in the 2640 Pf3k Isolates.

Frequency	Chrom. 12	SNP	Codon	Mutation
56.7% (1498)	C529418T	G3382A	GGA→AGA	G1128R
36.6% (966)	G529559T	C3241A	CAA→AAA	Q1081K
35.1% (927)	G532131T	C669A	GGC→GGA	G223G
14.0% (370)	G529707T	C3093A	GGC→GGA	G1031G
9.2% (243)	A532113T	T687A	GCT→GCA	A229A
8.9% (236)	C532133T	G667A	GGC→AGC	G223S
6.8% (180)	A532113C	T687G	GCT→GCG	A229A
5.2% (138)	A529665C	T3135G	AAT→AAG	N1045K
3.5% (93)	T529453C	A3347G	GAT→GGT	D1116G
3.4% (90)	T529091C	A3709G	ATA→GTA	I1237V
2.8% (75)	G531624T	C1176A	ACC→ACA	T392T
2.8% (75)	A529429C	T3371G	CTT→CGT	L1124R
2.2% (59)	C529709T	G3091A	GGC→AGC	G1031S
2.0% (53)	C532133T	G667A	GGC→AGA	}G223R
2.0% (53)	G532131T	C669A	GGC→AGA	
1.7% (46)	C529454A	G3346T	GAT→TAT	D1116Y
1.5% (40)	C532543T	G257A	AGT→AAT	S86N
1.3% (35)	G530705T	C2095A	CAA→AAA	Q699K
1.3% (34)	T532711C	A89G	GAA→GGA	E30G
1.3% (33)	A531636G	T1164C	GAT→GAC	D388D
1.2% (32)	G531280C	C1520G	ACT→AGT	T507S
1.0% (27)	G529457T	C3343A	CAA→AAA	Q1115K

Other Resistance Mutations

A literature search found 54 PfATP4 drug resistance mutations and these are listed in table 4. The 188 distinct SNP variants from the 2640 isolates are examined to see if any occur in the same codon as the tabulated resistance mutations. There are 2 synonymous mutations, P990P (A2970T, CCA→CCT, 5 isolates) and Q172Q (A516G, CAA→CAG, 1 isolate) that map to resistance codons. However, unlike the synonymous G223G (C669A) SNP, these synonymous SNPs do not increase the probability of the corresponding resistance mutations occurring: P990R (CCA→CG*), Q172H (CAA→CAC,CAT), Q172K (CAA→AAG,AAA).

Table 4: Drug Resistance Mutations in PfATP4

SNP	Compound	Compound Class	Citations
Q172H	KAE609	Spiroindolone	[8],[9]
	MMV007275	Benzamide	[9],[16]
	SJ733	Dihydroisoquinolones	[7]
Q172K	KAE609	Spiroindolone	[9]
	MMV011567	Benzamide	[9],[16]
V178I	PA21A092	Pyrazole-amide	[16]
A184S	KAE609	Spiroindolone	[14]
	KAE678	Spiroindolone	[16],[4],[10]
A187V	GNF-Pf4492	Aminopyrazole	[16],[1]
I203M	GNF-Pf4492	Aminopyrazole	[16],[4]
	KAE609	Spiroindolone	[14],[4]
	KAE678	Spiroindolone	[16],[4],[10]
V204L	KAE609	Spiroindolone	[8]
A211T	GNF-Pf4492	Aminopyrazole	[16],[4]
G223R	KAE609	Spiroindolone	[4]
	KAE678	Spiroindolone	[14],[16],[4],[10]
	GNF-Pf4492	Aminopyrazole	[4]
	SJ733	Dihydroisoquinolones	[7]
I263V	GNF-Pf4492	Aminopyrazole	[4]
	KAE609	Spiroindolone	[14],[4]
	KAE678	Spiroindolone	[16],[4],[10]
S312P	KAE609	Spiroindolone	[8]
L350H	SJ311	Dihydroisoquinolones	[16]
	SJ733	Dihydroisoquinolones	[7],[16]
L350V	KAE609	Spiroindolone	[8]
A353E	MMV011567	Benzamide	[16]
A353Q	KAE609	Spiroindolone	[9]
	MMV011567	Benzamide	[9]
N355Y	SJ733	Dihydroisoquinolones	[7]
G358S	SJ733	Dihydroisoquinolones	[7],[16]
I379N	KAE609	Spiroindolone	[8]
I398F	GNF-Pf4492	Aminopyrazole	[16],[4]
	KAE609	Spiroindolone	[16],[4],[14],[10]
	KAE678	Spiroindolone	[10]
	SJ733	Dihydroisoquinolones	[7]
V400A	KAE609	Spiroindolone	[8]
P412L	MMV772	Undisclosed	[16]
	SJ733	Dihydroisoquinolones	[7]
P412T	SJ733	Dihydroisoquinolones	[7],[16]
V414D	KAE609	Spiroindolone	[8]
V415D	SJ733	Dihydroisoquinolones	[7],[16]
T416N	KAE609	Spiroindolone	[8]
T418N	KAE609	Spiroindolone	[1],[16],[4],[14],[10]
	SJ733	Dihydroisoquinolones	[7]
A421E	KAE609	Spiroindolone	[8]
P437S	SJ311	Dihydroisoquinolones	[16],[7]
E895K	KAE609	Spiroindolone	[8]
F917L	MMV772	Undisclosed	[16]
	SJ733	Dihydroisoquinolones	[7]
L928F	SJ733	Dihydroisoquinolones	[7],[16]
L938I	KAE609	Spiroindolone	[8]
P966A	KAE609	Spiroindolone	[8]
	SJ733	Dihydroisoquinolones	[7]
P966S	SJ247	Dihydroisoquinolones	[16]
	SJ733	Dihydroisoquinolones	[7]
P966T	SJ279	Dihydroisoquinolones	[16]
	SJ733	Dihydroisoquinolones	[7]
A967G	KAE609	Spiroindolone	[8]
P990R	GNF-Pf4492	Aminopyrazole	[1],[16],[4]
	KAE609	Spiroindolone	[16],[4],[14],[1],[10]
	KAE678	Spiroindolone	[16],[4],[10]
	SJ733	Dihydroisoquinolones	[7]
A1158V	KAE609	Spiroindolone	[8]
A1207V	KAE609	Spiroindolone	[8]
D1247Y	KAE609	Spiroindolone	[16],[4],[14],[10]