

a
T1 tumour fraction
power, %
rats per arm
90 %
80 %
8
27
5 pp
10 pp
15 pp
20 pp
25 pp
15 pp, pilot SD
pilot bound, 2.9 pp
b
T2a, T2c invasion interval, floor
interval bound
rats per arm, or control pieces
half-width, p = 0.1
half-width, p = 0.2
half-width, p = 0.35
half-width, p = 0.5
floor, 0 of n
0.15
0.10
26
39
29
c
T2b invasion, LITT vs none
power, %
rats per arm
80 %
44
58
90
0.20 vs 0.50, Holm
0.20 vs 0.50
0.20 vs 0.40
d
T3 recurrence-specific arm event
power, %
rats per arm
80 %
10
17
in 50 % of recurrences
in 60 % of recurrences
in 80 % of recurrences
60 %, unadjusted
e
T4 not computed
f
T5 not computed
g
T6 human:rat titration
replicates per level
assumed SD per mixture level
z, as pre-stated
t
31
4
SD = RMSE goal
mouse-mixture error
h
T7 host response, sham + LITT vs naive
power, %
rats per arm
12
17
80 %
d = 1.2, Holm
d = 0.8
d = 1.2
d = 1.5
i
T8 clones, share of animals
95 % lower bound
barcoded animals
0.80
0.70, N1
0.60, N2
16
9
no failure
1 failure
2 failures