
(a) variant call per subtype
sensitivity or specificity, % (95 % interval)
100
80
60
40
20
0
pre-stated sensitivity bound 85 %
PENDING G6
variant sensitivity and specificity per subtype,
held-out transitional persons, 95 % interval
sensitivity (variant called on a variant)
specificity (no call on an ordinary spine)
sixth lumbar
thirteenth thoracic
sacralised fifth lumbar
four-lumbar
both
transitional subtype (n per subtype from the G1 inventory)
(b) exact level, ordinary and transitional
exact level, % of test persons (95 % interval)
pre-stated bound, transitional 90 %
PENDING G6
exact level by method and stratum,
person-clustered 95 % interval
ordinary test persons
transitional test persons
registration ceiling*
classifier
count
coupled
method; * truth boxes carrying the CT-imported name
(c) surgeon + overlay vs coupled call
disagreement with the operated level, % of transitional test persons
PENDING G6
disagreement with the operated level,
transitional test persons
surgeon's count reconciled with the imported label
coupled call (names the variant)
(d) power: lower bound against test persons
Wilson 95 % lower bound, %
split target n ≥ 60
k = 0
k = 1
k = 2
k = 3
k = 5
exact level
lower bound ≥ 90 %
variant sensitivity
lower bound ≥ 85 %
today: 89 subjects with truth → 37 test persons
(3 real, all in test, + 40 % of 86 CT; 50 % rule: 46)
COBRA2026 planning CTs not yet screened
transitional test persons, n (k = errors among them)