Comprehensive Radiomorphometric Analysis of Lumbar Spine Pathologies: Integrating Advanced Computational Diagnostics and Standardized Clinical Grading Systems

The diagnostic framework for lumbar spinal disorders has transitioned from conventional qualitative assessment to a sophisticated paradigm of radiomorphometric quantification. This evolution is necessitated by the immense global burden of low back pain, which affected approximately 619 million individuals in 2020 and remains the primary driver of physical disability worldwide.[1] In the clinical interpretation of magnetic resonance imaging (MRI), the objective identification of morphometric markers is critical for distinguishing between normal age-related changes and clinically significant degenerative conditions.[4] The lumbar spine functions as a complex "three-joint unit," where the health of the intervertebral disc is inextricably linked to the paired posterior facet joints.[4] Consequently, structural failure in any single component often initiates a degenerative cascade that compromises the entire motion segment.[7] The advent of artificial intelligence (AI) and automated segmentation has provided clinicians with tools to perform these intricate measurements with expert-level precision, reducing the inter-reader variability that historically hampered diagnostic consistency.[10]

Vertebral Body Morphometry and Structural Integrity Assessment

The vertebral body serves as the primary load-bearing structure of the spine, and its morphometric integrity is the cornerstone of sagittal balance and structural stability. Deformities of the vertebral body, particularly height reduction, are indicative of compression fractures which often go underreported in clinical practice.[13] Radiologists utilize two primary methods for evaluating these changes: the Genant semiquantitative (SQ) method and quantitative morphometry (QM).[13] While the SQ method relies on visual grading (normal, mild, moderate, or severe), the QM method involves precise landmarking to calculate vertebral heights and indices.[13]

Landmark Logic and Height Ratios

Quantitative assessment typically requires the localization of six specific landmarks on the vertebral body: two at the anterior edges, two at the posterior edges, and two at the midpoint

of the superior and inferior endplates.[13] These points allow for the calculation of three primary heights: the anterior height (Ha), the middle height (Hm), and the posterior height (Hp).[14] These measurements are used to derive height ratios that define the type of fracture present.

Deformity Index	Mathematical Formula	Clinical Finding
Compression Ratio	$\frac{Hm}{Ha}$ or $\frac{Hm}{Hp}$	Values < 0.8 suggest biconcave fracture.
Wedge Ratio	$\frac{Ha}{Hp}$	Values < 0.8 suggest anterior wedge fracture.
Crush Ratio	$\frac{Hp}{Ha}$	Values < 0.8 suggest posterior height loss.

Automated deep learning systems, such as SpineToolKit, have demonstrated the ability to locate these landmarks across MRI, CT, and radiographic studies with a mean height measurement error of less than 3%.[13] This level of precision is vital for opportunistic screening, allowing for the early detection of osteoporotic fractures that may not be immediately obvious upon visual inspection.[10] Furthermore, the anterior vertebral body height in the lumbar region is typically greater than the posterior height, a morphometric feature that contributes to the natural lumbar lordosis angle (LLA).[5] The LLA is another critical quantitative marker, with automated systems reporting a mean error of less than 3.6° in its measurement.[13]

Segmental Deformity and Translational Instability

Beyond discrete height loss, the relationship between adjacent vertebral bodies is a primary indicator of spinal stability. Degenerative spondylolisthesis is defined by the translation of one vertebral body over another, usually diagnosed when translation is $\geq 3\ \text{mm}$ on standing lateral radiography.[16] Morphometric indices developed for automated tests can now detect both translational and vertical instabilities from flexion-extension radiographs.[18] Rotation-dependent translation (RDT) measures sagittal plane translation adjusted for the degree of rotation, providing a level-dependent assessment of mechanical irritation of the

nerve roots.[18] Similarly, sagittal rotation-dependent disc widening (RDDW) quantifies motion perpendicular to the endplates, which is a hallmark of vertical instability.[18]

Instability Marker	Measurement Logic	Normal Limit
Sagittal Translation	Linear displacement (mm)	$< 2 - 4$ mm
RDT Index	Translation per degree of rotation	Statistical deviation from mean
AVI-Index	Vertical motion perpendicular to endplate	Level-specific tables

The significance of these measurements lies in their ability to distinguish physiological movement from pathological instability, which guides the decision for surgical fusion versus simple decompression.[18]

The Disco-Vertebral Junction and Modic Phenomenon

The interface between the intervertebral disc and the vertebral endplate, known as the disco-vertebral junction (DVJ), is a common site of inflammation and structural failure.[19] Modic changes (MCs) are signal intensity variations in the vertebral bone marrow adjacent to the endplates, often associated with degenerative disc disease (DDD) and chronic low back pain.[19] These changes are considered dynamic, representing various stages of an inflammatory cascade triggered by mechanical trauma, discitis, or autoimmune reactions.[21]

Classification and Signal Characteristics

Modic changes are classified into three types based on paired T1-weighted and T2-weighted MRI signal characteristics.[21]

Modic Type	T1 Signal	T2 Signal	Pathological Basis

Type 1	Hypointense	Hyperintense	Fibrovascular tissue; bone marrow edema.
Type 2	Hyperintense	Iso/Hyperintense	Fatty replacement of bone marrow.
Type 3	Hypointense	Hypointense	Extensive subchondral bone sclerosis.

Automated detection systems for MCs utilize feature extraction techniques such as local binary patterns and Hu's moments to identify intensity variations with accuracies reaching 85.91%.[24] Clinical research has increasingly emphasized the importance of lesion burden over Modic type alone.[29] The Modic Change Grade (MCG), or Modic-Udby definition, quantifies the vertical extent of the changes within the vertebral body.[29]

- **Grade A (Minor Burden)**: Involvement of $< 25\%$ of the vertebral body height or volume.
- **Grade B (Major Burden)**: Involvement of $25–50\%$.
- **Grade C (High Burden)**: Involvement of $> 50\%$.

High-burden lesions (MCG-B and MCG-C) are more strongly predictive of patient-reported disability and are used to integrate vertebrogenic strategies into surgical planning.[18] Recent interventional studies, such as those evaluating basivertebral nerve (BVN) ablation, have shown significant sustained improvements in outcomes for patients with symptomatic MC type 1 or 2.[22]

Advanced Intervertebral Disc Radiomorphometrics

Intervertebral disc (IVD) degeneration is a hallmark of the spinal aging process and is a primary driver of spinal stenosis and radiculopathy.[1] The morphometric assessment of the IVD focuses on height maintenance, hydration status, and the extent of displacement.[33]

The Disc Height Index (DHI) Framework

The Disc Height Index (DHI) is a standardized method for communicating disc height that normalizes the measurement against the size of the vertebral column.[37] By using ratios,

variations in patient positioning and magnification factors are mitigated, allowing for reliable longitudinal analysis.[37]

The most common formula, adapted from the work of Farfan, is expressed as:

$$DHI = \frac{Ha + Hp}{Ds + Di} \times 100$$

where Ha and Hp are the anterior and posterior disc heights, and Ds and Di are the superior and inferior disc depths.[38]

Research has yielded several alternative methods for calculating DHI based on different landmark configurations:

Method ID	Calculation Formula	Contextual Application
Method 1	$DHI =$	Ratio of anterior+posterior height to disc diameter.[37]
Method 2	$DHI =$	Ratio of mid-disc height to mid-vertebral body height.[37]
Method 3	$DHI =$	Ratio of mid-disc height to disc diameter.[37]
Method 6	$DHI =$	Mean IVD height relative to mean vertebral heights.[37]

Studies have demonstrated that DHI measurements have high systematic reliability (Dice Similarity Coefficient > 90), although minor biases can occur in the setting of severe spondylolisthesis, where corner point measurements may overestimate the actual height.[38]

Pfirrmann Grading and Internal Composition

Hydration and internal structural organization are graded using the Pfirrmann classification, which relies on the distinction between the nucleus pulposus and the annulus fibrosus on T2-weighted MRI.[34]

Pfirrmann Grade	Distinction of Nucleus/Annulus	Signal Intensity	Disc Height
Grade I	Clear	Hyperintense	Normal
Grade II	Clear	Hyperintense (with horizontal bands)	Normal
Grade III	Unclear	Intermediate	Normal to slightly decreased
Grade IV	Lost	Intermediate to Hypointense	Normal to moderately decreased
Grade V	Lost	Hypointense (Black)	Collapsed disc space

Sources:.[34]

Innovative research has also integrated latent geometric features extracted by Vision Transformers (e.g., SwinUNETR-V2) with standard determinants like anteroposterior width and lateral width to enhance the prediction of disc narrowing beyond manual measures alone.[43] Automated analysis of the internal composition, such as the NP-to-CSA ratio (nucleus pulposus to cross-sectional area), has shown that computer-assisted methods provide outcomes comparable to autopsy-derived data while overcoming the labor-intensive nature of manual tracing.[46]

Central Spinal Canal Stenosis Quantification

Lumbar spinal stenosis (LSS) is diagnosed by the objective evidence of neurovascular compromise within the spinal canal.[3] Quantitative criteria for central stenosis focus on the anteroposterior (AP) diameter and the cross-sectional area of the dural sac (DSCA).[48]

Volumetric and Linear Canal Indices

The thecal sac is filled with cerebrospinal fluid (CSF), and its shape is best evaluated on axial T2-weighted images where the contrast between fluid and soft tissue is maximal.[54]

Clinical Classification	DSCA Threshold (mm2)	Mid-Sagittal AP Diameter (mm)
Normal	> 100	> 12
Relative Stenosis	$75 - 100$	$10 - 12$
Absolute Stenosis	$< 70 - 75$	$< 7 - 10$

Sources:.[49]

Automated systems like SpineLogic utilize deep learning to segment the thecal sac and measure the AP and mediolateral (ML) distances with human-level accuracy.[11] Mean absolute errors for the AP distance typically range from $0.59\ \text{mm}$ to $0.75\ \text{mm}$, which is well within the clinically acceptable range.[48] Furthermore, research into Indian populations established a critical canal dimension of $11.13\ \text{mm}$ as the threshold at which symptoms of stenosis frequently appear.[52]

Morphological Variants of the Canal

The axial shape of the spinal canal is a critical morphometric pathology that influences the risk of nerve root entrapment.[59]

Vertebral Level	Predominant Shape	Frequency (%)
L1	Oval	$85 - 95\%$

L2	Oval	90%
L3	Triangular	$80 - 95\%$
L4	Triangular	95%
L5	Trefoil	$60 - 65\%$

Sources:.[59]

A trefoil-shaped canal at the lower lumbar levels is particularly predisposed to lateral recess narrowing because the nerve roots are positioned in narrow niches bordered by the pedicle and facet joints.[31] Short pedicles, often a developmental trait, further exacerbate this narrowing by reducing the overall dimensions of the bony canal.[52]

Lateral Recess and Subarticular Zone Narrowing

Lateral recess stenosis (LRS) involves the compression of the traversing nerve root as it takes off from the thecal sac, before it enters the neural foramen.[61] Failure to adequately decompress this region is a leading cause of failed back surgery syndrome.[31]

Depth and Angular Criteria

Topographically, the lateral recess is a triangular space bounded laterally by the pedicle, posteriorly by the superior articular facet (SAP), and anteriorly by the vertebral body and disc.[31]

Parameter	Stenosis Diagnostic Threshold	Pathophysiology
Lateral Recess Depth (LRD)	$\leq 3 -$	Distance from VB dorsal surface to ventral SAF segment.[52]
Lateral Recess Angle (LRA)	$<$	Angle between the floor

		and roof of the recess.[53]
Lateral Recess Height	$\leq 2 -$	Reduced by facet hypertrophy or endplate spurring.[53]

Quantitative measurements of the lateral recess have demonstrated better repeatability than visual assessments.[64] Specifically, the lateral recess angle has been identified as a superior predictor of surgical findings compared to interfacet distance.[63] The narrowest point of the lateral spinal canal (NPLC) is typically found at a distance of $0.7 \pm 0.3\ \text{cm}$ from the dura at the L4-L5 level.[62]

Mechanisms of Lateral Entrapment

Nerve root compression in the lateral recess typically follows two morphologic pathways.[31] The first is the "trefoil narrowing," where the nerve root remains in its standard position but the recess narrows in an anteroposterior fashion due to progressive facet or endplate changes.[31] The second is the "acute angular pinch," where narrowing occurs simultaneously from all directions due to equal contribution from the disc, facet joints, and ligamentum flavum, often resulting in medial deflection of the root.[31]

Neural Foraminal Morphometry and Volumetric Analysis

Neural foraminal stenosis involves the narrowing of the bony exit where the nerve root exits the spinal canal.[65] This condition is defined by a decrease in disc height, hypertrophic facet changes, and buckling of the ligamentum flavum.[36]

The Lee Grading System for Foraminal Narrowing

The Lee system is a widely accepted qualitative framework that integrates fat obliteration patterns and nerve root compression.[50]

Lee Grade	Severity	Description
Grade 0	Normal	No foraminal stenosis; fat well-maintained.

Grade 1	Mild	Fat obliteration in two opposing directions (vertical or transverse).
Grade 2	Moderate	Fat obliteration in four directions without morphologic change in nerve root.
Grade 3	Severe	Morphologic change/compression of the nerve root; fat entirely lost.

Sources:.[37]

The Lee system has shown nearly perfect interobserver agreement ($\kappa > 0.81$) and strong clinical correlation, particularly in younger patient cohorts.[65]

Quantitative Volumetric Models

To move beyond qualitative grading, clinicians have adopted volumetric measurements of the lumbar foramina. approximating the foramen as an elliptical cylinder allows for a precise estimation of available space.[70]

The major axis (a) is measured as the longest distance across the foramen in the sagittal view, the minor axis (b) as the shortest perpendicular distance, and the depth (H) as the anterior-to-posterior distance in the axial view.[70]

Level	**Right Volume (mm3)**	**Left Volume (mm3)**	**SD**
L1/L2	579.92	594.43	$\pm 55 / \pm 44$

L2/L3	688.22	715.87	$\pm 55 / \pm 48$
L3/L4	761.70	790.30	$\pm 59 / \pm 50$
L4/L5	787.82	809.61	$\pm 29 / \pm 57$
L5/S1	824.24	N/A	± 68

Sources:.[70]

A progressive volumetric increase is observed in the craniocaudal direction, reflecting the spine's adaptation to increasing biomechanical stresses at the lower levels.[70] A novel metric, the Nerve root/Foramen (N/F) ratio, has also emerged to evaluate the relative occupancy of the exiting nerve root, providing an objective measure that correlates with subjective severity scores.[71]

Posterior Element Pathologies and Hypertrophy

The posterior spinal column, including the ligamentum flavum and the spinous processes, plays a pivotal role in the pathogenesis of stenosis and midline back pain.[7]

Ligamentum Flavum Thickness and Area

The ligamentum flavum (LF) is an elastic structure that maintains spinal alignment and limits excessive movement.[32] Chronic mechanical stress leads to the replacement of elastic fibers with fibrous scar tissue, resulting in hypertrophy and buckling into the canal.[72]

Parameter	Threshold	Clinical Significance
Normal LFT	$3.5 -$	Baseline at L4-L5 level.[73]
Hypertrophied LFT	$\geq$	Upper limit for healthy adults.[72]

Severe LFH	$\geq$	High risk of neurogenic claudication.[76]
LFA Cutoff	$105.90\ \text{mm}$	Optimal predictor for central canal stenosis.[78]

Measurements are typically taken on axial T1-weighted or T2-weighted images at the level of the facet joints.[77] Research suggests that Ligamentum Flavum Area (LFA) is a more sensitive measurement parameter than thickness alone because it encompasses the entire cross-sectional burden and remains stable despite ligamentous asymmetry.[78] LFT is consistently higher on the left side in some cohorts and increases with age across all lumbar levels.[72]

Baastrup Disease (Kissing Spine Syndrome)

Baastrup disease refers to the close approximation and impingement of adjacent posterior spinous processes.[7] This condition is most prevalent at the L4-L5 level and is strongly associated with excessive lordosis, obesity, and the natural process of aging.[7]

Morphometric Pathologies in Baastrup Disease:

- **Spinous Process Apposition**: Contact between adjacent processes leads to reactive cortical sclerosis and hypertrophy.[7]
- **Interspinous Bursitis**: Appearing as bright, high-signal-intensity areas on T2-weighted or STIR sagittal images, this represents adventitious bursa formation.[84]
- **Osseous Remodeling**: Flattening and enlargement of the articulating surfaces of the spinous processes.[7]
- **Epidural Extension**: Fluid and inflammation from the interspinous space can extend through natural clefts in the ligamentum flavum to create midline epidural cysts, resulting in posterior thecal sac compression.[84]

In symptomatic individuals over the age of 80, the incidence of radiologic findings for Baastrup disease is as high as 81%.[82] MRI is the most sensitive modality for early detection, as it can identify interspinous bursitis before the onset of osseous remodeling seen on radiographs or CT.[7]

Facet Joint Morphometrics and Osteoarthritis

The facet joints connect adjacent vertebrae and bear approximately 33% of the dynamic compressive load.[6] Degeneration of these joints, or facet joint osteoarthritis (FJOA), is present in virtually all individuals over the age of 60.[8]

Grading Systems for Facet Arthropathy

Various grading systems (Pathria, Grogan, and Weishaupt) are utilized to objectify FJOA severity across CT and MRI.[89]

Grading Criteria	Grogan (MRI)	Weishaupt (CT/MRI)
Normal (Grade 1)	Uniformly thick cartilage.	Joint space width $2-4\ \mathrm{mm}$.
Mild (Grade 2)	Eroded/irregular cartilage.	Narrowing $<2\ \mathrm{mm}$; small osteophytes.
Moderate (Grade 3)	Incomplete coverage; dense bone.	Moderate osteophytes; bone erosion.
Severe (Grade 4)	Cartilage absence; large osteophytes.	Severe OA; cysts; vacuum phenomenon.

Sources:.[4]

FJOA is significantly exacerbated by the progression of disc degeneration at the same level, reinforcing the "three-joint complex" concept where pathology in one unit influences the biomechanics of the others.[4] Facet synovial cysts, often medial to the joint, can compress the dural sac and nerve roots, appearing as focal outpouchings on axial MRI.[6]

Facet Orientation and Tropism

Facet orientation is the angle of the joint in the transverse plane relative to the midsagittal line.[6] Facet Tropism (FT) is defined as an asymmetry between the right and left facet angles.[6]

The facet angle is calculated by intersecting the line along the posterior border of the vertebral body with a line drawn through the margins of the facet joint.[16]

FT Grade (Ko et al.)	Angulation Difference	Clinical Association

Grade 0	$\leq$	Normal asymmetry.
Grade 1	$7-$	Moderate tropism; linked to disc prolapse.[90]
Grade 2	$\geq$	Severe tropism; linked to spondylolisthesis.[16]

Thresholds for defining tropism vary in the literature between $7^{\circ}, 8^{\circ}$, and 10°.[16] FT is established as a silent driver of lumbar degeneration, as the asymmetry alters rotational kinematics and creates shear forces that accelerate disc dehydration.[4]

Cervical Spine Specific Morphometrics

While lumbar pathologies dominate back pain research, specific morphometric markers are essential for the cervical spine due to its higher mobility and the delicate nature of the spinal cord in this region.[67]

Myelopathic Indices and Cord Compression

The severity of Degenerative Cervical Myelopathy (DCM) is quantified using normalized morphometric measures of cord compression.[103]

1. **Modified Spinal Cord Compression (MSCC)**: Computed by normalizing the anteroposterior cord diameter at the compression site by the diameters above and below the level.[103]
2. **Cross-Sectional Area (CSA)**: Automatic calculation of the cord's CSA from MRI scans is used to predict the therapeutic transition from conservative to operative management.[103]
3. **K-Line**: Connects the midpoints of the spinal cord at C2 and C7. A "K-line negative" condition occurs when the anterior depressor (typically a herniated disc) makes contact with this line, indicating significant compression and helping to plan surgical approach.[105]

Automated systems achieving Dice coefficients above 0.90 for cervical anatomy allow for the rapid detection of abnormal intervertebral disc gaps and vertebral-to-neural foramen distances, facilitating early identification of degenerative changes.[104]

Computational Paradigms: The Future of Automated Radiomorphometry

The development of high-quality annotated datasets has paved the way for AI systems to replace labor-intensive manual measurements in clinical workflows.[106]

The RSNA LumbarDISC Dataset and AI Challenges

The Radiological Society of North America (RSNA) published the Lumbar Degenerative Imaging Spine Classification (LumbarDISC) dataset, which is the largest publicly available collection of adult lumbar MRI examinations.[108] Comprising 2,697 patients with multisequence, multiplanar images, it has served as the "standard of truth" for training models to localize and classify conditions across levels L1/L2 through L5/S1.[1]

Key findings from AI implementations on this dataset include:

- **Multi-Task Classification**: Models can simultaneously grade central canal, neural foraminal, and subarticular stenosis into three severity grades (Normal/Mild, Moderate, or Severe) with high precision.[44]
- **Architecture Innovations**: Convolutional neural networks (CNNs) like YOLOv8 and Attention U-Nets have achieved Dice scores of 0.94 for lumbar segmentation, demonstrating superior robustness across various hardware and patient demographics.[30]
- **Weighted Log Loss Metrics**: To ensure clinical utility, AI models are evaluated using weighted metrics that penalize the misclassification of "Severe" conditions more heavily (sample weight of 4) than "Normal" ones.[2]

Integration into Clinical Workflows

Solutions such as Remedy Logic's SpineLogic are now deployed in over 100 facilities, providing automated measurements and flagging abnormalities directly within the radiologist's PACS/RIS systems.[11] These "AI co-pilots" expedite diagnostic turnaround by pre-populating reporting templates and generating summary tables for quick clinician confirmation.[11] Automated quantification not only saves time but also enables the creation of large-scale normative datasets, which are essential for distinguishing pathological narrowing from normal anatomical variation.[115]

Synthesis of Radiomorphometric Insights

The clinical relevance of lumbar spine morphometry is best understood through the synthesis of multiple structural markers. A single narrow canal dimension may be asymptomatic, but when combined with high-burden Modic changes and severe facet tropism, it represents a high-risk phenotype for degenerative spondylolisthesis and debilitating radiculopathy.[16]

Integrated Clinical Thresholds for Lumbar Pathology

Pathological Entity	Measurement Marker	Critical Threshold for Intervention

Central Stenosis	DSCA	$< 70\ \text{mm}^2$ (Absolute Stenosis)
Lateral Recess Stenosis	Depth (LRD)	$\leq 2\ \text{mm}$ (Severe)
Foraminal Stenosis	Lee Grade	Grade 3 (Nerve morphologic change)
LF Hypertrophy	Thickness	$> 5\ \text{mm}$ (Significant encroachment)
Disc Degeneration	DHI / Pfirrmann	Grade V / $> 50\%$ height loss
Vertebral Deformity	Height Ratio	< 0.75 (Moderate/Severe fracture)
Modic Burden	MCG	Grade C ($> 50\%$ vertebral volume)

Sources:.[29]

Furthermore, the relationship between clinical symptoms and imaging findings remains complex; many participants in population-based cohorts exhibit radiographic narrowing without clinical symptoms.[1] This highlights the ongoing need for "precision phenotyping," where morphometric data are integrated with clinical scoring (e.g., Oswestry Disability Index) to guide surgical decision-making and patient selection.[29]

Future Directions: Precision and Functional Imaging

The frontier of radiomorphometrics involves moving beyond static anatomical measurement toward functional and biochemical analysis.

- **Ultrashort Time to Echo (UTE) MRI**: This technique allows for the direct quantitative evaluation of short-T2 tissues such as the cartilaginous endplate (CEP) and the annulus fibrosus, offering insights into early-stage degeneration that standard sequences cannot provide.[19]
- **Upright and Load-Bearing MRI**: Morphometric changes under axial loading, such as the reduction in DSCA and narrowing of the neural foramina when standing, provide a more accurate representation of the stenosis a patient experiences during daily activity.[52]
- **Precision Phenotyping of Modic Changes**: Future research aims to distinguish between bacterial and sterile MCG subtypes using radiomics and quantitative MRI endpoints (e.g., PDFF), allowing for targeted therapies such as antibiotics versus regenerative repair.[29]

In conclusion, the integration of automated computational tools with standardized morphometric grading systems has established a new standard for spinal diagnosis. By providing objective, reproducible, and granular data, these radiomorphometric frameworks empower clinicians to deliver more accurate diagnoses, optimize treatment trajectories, and ultimately improve patient outcomes in the complex landscape of degenerative spinal disease.[1]

Works cited

1. RSNA Lumbar Spine Degenerative Classification AI Challenge (2024), accessed February 21, 2026, https://www.rsna.org/artificial-intelligence/ai-image-challenge/lumbar-spine-degenerative-classification-ai-challenge
2. RSNA 2024 Lumbar Spine Degenerative Classification | Kaggle, accessed February 21, 2026, https://www.kaggle.com/competitions/rsna-2024-lumbar-spine-degenerative-classification
3. Automated measurement of anteroposterior diameter and foraminal widths in MRI images for lumbar spinal stenosis diagnosis - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC7605707/
4. Evaluation of MRI and CT parameters to analyze the correlation between disc and facet joint degeneration in the lumbar three, accessed February 21, 2026, https://rcastoragev2.blob.core.windows.net/93003bf91d770813fb9d4c929c37e22f/PMC6783151.pdf
5. The morphology of the lumbar vertebrae: a systematic review with meta-analysis of 1481 individuals with implications for spine surgery - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11624215/
6. Facet joint syndrome: from diagnosis to interventional management - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC6206372/
7. Baastrup's disease (kissing spines syndrome): a pictorial review - PMC - NIH, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC4330238/
8. Facet Synovitis - Radsource, accessed February 21, 2026,

https://radsource.us/facet-synovitis/
9. Kissing Spine of Baastrup Syndrome: Is there Surgical Treatment? - Clinics in Surgery, accessed February 21, 2026, https://www.clinicsinsurgery.com/open-access/kissing-spine-of-baastrup-syndrome-is-there-surgical-treatment-9360.pdf
10. Artificial Intelligence in Spinal Imaging and Patient Care: A Review of Recent Advances, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11224760/
11. Remedy Logic: AI for Spine MRI Interpretation, accessed February 21, 2026, https://www.remedylogic.com/
12. Deep learning-based automated segmentation and quantification of the dural sac cross-sectional area in lumbar spine MRI - Frontiers, accessed February 21, 2026, https://www.frontiersin.org/journals/radiology/articles/10.3389/fradi.2025.1503625/full
13. Vertebral Deformity Measurements at MRI, CT, and Radiography Using Deep Learning, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC8823454/
14. An automated algorithm for quantitative morphometry of thoracic and lumbar vertebral bodies in lateral radiographs - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11911063/
15. Automatic measurement of anatomical parameters of the lumbar vertebral body and the intervertebral disc on radiographs by deep learning - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11320507/
16. A Radiological Study of the Association of Facet Joint Tropism and Facet Angulation With Degenerative Spondylolisthesis - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC6945004/
17. A Radiological Study of the Association of Facet Joint Tropism and Facet Angulation With Degenerative Spondylolisthesis - Neurospine, accessed February 21, 2026, https://www.e-neurospine.org/journal/view.php?doi=10.14245/ns.1836232.116
18. Automated radiographic metrics for diagnosing lumbar spine instability: a cross-sectional observational study, accessed February 21, 2026, https://qims.amegroups.org/article/view/135403/html
19. Quantitative magnetic resonance imaging of the lumbar intervertebral discs - PMC - NIH, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5219962/
20. Automated measurement of anteroposterior diameter and foraminal widths in MRI images for lumbar spinal stenosis diagnosis | PLOS One - Research journals, accessed February 21, 2026, https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0241309
21. Modic changes - An evidence-based, narrative review on its patho-physiology, clinical significance and role in chronic low back pain - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC7452231/
22. Modic Changes as Biomarkers for Treatment of Chronic Low Back Pain - MDPI,

accessed February 21, 2026, https://www.mdpi.com/2227-9059/13/7/1697
23. Evaluating CNN Architectures for the Automated Detection and Grading of Modic Changes in MRI: A Comparative Study - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11735353/
24. Automatic Detection and Classification of Modic Changes in MRI Images Using Deep Learning: Intelligent Assisted Diagnosis System | Request PDF - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/publication/375448849_Automatic_Detection_and_Classification_of_Modic_Changes_in_MRI_Images_Using_Deep_Learning_Intelligent_Assisted_Diagnosis_System
25. Comparing image normalization techniques in an end-to-end model for automated modic changes classification from MRI images - Research Collection, accessed February 21, 2026, https://www.research-collection.ethz.ch/bitstreams/5e50ee4b-9e46-4ff4-b379-be2fc05051ba/download
26. Automatic detection and voxel-wise mapping of lumbar spine Modic changes with deep learning - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC9238279/
27. Automatic Modic Changes Classification in Spinal MRI | Request PDF - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/publication/304664299_Automatic_Modic_Changes_Classification_in_Spinal_MRI
28. Automatic Detection and Classification of Modic Changes in MRI Images Using Deep Learning: Intelligent Assisted Diagnosis System - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC10782244/
29. Modic Changes in spine surgery: clinical update 2025 - AO Foundation, accessed February 21, 2026, https://www.aofoundation.org/spine/about-aospine/blog/2025/udby-modic-changes-in-spine-surgery
30. Automated lumbar spine segmentation in MRI using an enhanced U-Net with inception module and dual-output mechanism - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC12603128/
31. Correlation of lumbar lateral recess stenosis in magnetic resonance imaging and clinical symptoms - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5441455/
32. Ligamentum Flavum Hypertrophy: Non-Invasive KL Care - Chiropractic Specialty Center, accessed February 21, 2026, https://www.mychiro.com.my/articles/ligamentum-flavum-hypertrophy/
33. An automatic tool for lumbar deformities and intervertebral disc degeneration using deep-learning based 3D segmentation - ISMRM, accessed February 21, 2026, https://cds.ismrm.org/protected/23MProceedings/PDFfiles/1558_PEI66VR8k.html
34. Lumbar spine MRI annotation with intervertebral disc height and Pfirrmann grade predictions | PLOS One - Research journals, accessed February 21, 2026, https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0302067

35. The Quantitative Evaluation of Automatic Segmentation in Lumbar Magnetic Resonance Images - PMC - PubMed Central, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11224749/
36. AI and Deep Learning for Automated Segmentation and Quantitative Measurement of Spinal Structures in MRI - arXiv, accessed February 21, 2026, https://arxiv.org/html/2503.11281v2
37. Radiographic evaluation of lumbar intervertebral disc height index: an intra and inter-rater agreement and reliability study - medRxiv.org, accessed February 21, 2026, https://www.medrxiv.org/content/10.1101/2021.11.29.21267030v1.full-text
38. The disc height index (DHI) measurement method using a section of a... - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/figure/The-disc-height-index-DHI-measurement-method-using-a-section-of-a-lumbar-spine-standing_fig2_361954708
39. Disc height measurement. Ha: anterior disc height, Hp: posterior disc... | Download Scientific Diagram - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/figure/Disc-height-measurement-Ha-anterior-disc-height-Hp-posterior-disc-height-Ds_fig4_283741167
40. Open Lumbar Spine Image Analysis: A 3D Slicer Extension for Segmentation, Grading, and Intervertebral Disc Height Index With Multi–Data Set Validation - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC12695293/
41. Tracing the disc: The novel qualitative morphometric MRI based disc degeneration classification system - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC10945309/
42. The relationship between quantitative measures of disc height and disc signal intensity with Pfirrmann score of disc degeneration - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC4917505/
43. A CNN Autoencoder for Learning Latent Disc Geometry from Segmented Lumbar Spine MRI, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11908304/
44. RSNA 2024 Lumbar Spine Degenerative Classification Training Script - GitHub, accessed February 21, 2026, https://github.com/anto18671/lumbar-spine-degenerative-classification
45. AI and Deep Learning for Automated Segmentation and Quantitative Measurement of Spinal Structures in MRI - arXiv, accessed February 21, 2026, https://arxiv.org/html/2503.11281v1
46. Morphology and Composition of Lumbar Intervertebral Discs: Comparative Analyses of Manual Measurement and Computer-Assisted Algorithms - MDPI, accessed February 21, 2026, https://www.mdpi.com/2306-5354/11/5/466
47. Imaging spinal stenosis | Applied Radiology, accessed February 21, 2026, https://appliedradiology.com/articles/imaging-spinal-stenosis
48. AI-Based Measurement of Lumbar Spinal Stenosis on MRI: External Evaluation of a Fully Automated Model - PubMed, accessed February 21, 2026, https://pubmed.ncbi.nlm.nih.gov/38426719/
49. Quantitative radiologic criteria for the diagnosis of lumbar spinal stenosis: a

systematic literature review - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC3161920/
50. Magnetic Resonance Imaging Grading Systems for Central Canal and Neural Foraminal Stenoses of the Lumbar and Cervical Spines With a Focus on the Lee Grading System - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC9971835/
51. Lumbar Spinal Stenosis - StatPearls - NCBI Bookshelf - NIH, accessed February 21, 2026, https://www.ncbi.nlm.nih.gov/books/NBK531493/
52. Magnetic resonance imaging in the diagnosis of lumbar canal stenosis in Indian patients, accessed February 21, 2026, https://joas.org.in/magnetic-resonance-imaging-in-the-diagnosis-of-lumbar-canal-stenosis-in-indian-patients/
53. Quantitative radiologic criteria for the diagnosis of lumbar spinal stenosis: a systematic literature review - ZORA, accessed February 21, 2026, https://www.zora.uzh.ch/server/api/core/bitstreams/6a62a8c1-efd8-41a7-8b5b-c74e1f3df276/content
54. AI-Based Measurement of Lumbar Spinal Stenosis on MRI - ZORA, accessed February 21, 2026, https://www.zora.uzh.ch/server/api/core/bitstreams/37e21fe4-3087-4609-8c51-2ed53c2b5ea1/content
55. The measure method of height/depth of lateral recess: the distance... - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/figure/The-measure-method-of-height-depth-of-lateral-recess-the-distance-between-the-anterior_fig2_317346587
56. Imaging Lumbar Spinal Stenosis - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/publication/12108909_Imaging_Lumbar_Spinal_Stenosis
57. SpineLogic: AI-Driven Insights for Spine MRI | TestDynamics, accessed February 21, 2026, https://testdynamics.net/radiology-ai-software/spinelogic/
58. Exploring Variations in Lumbar Canal Width: An MRI Study on Asymptomatic Patients by Age and Gender - MDPI, accessed February 21, 2026, https://www.mdpi.com/2077-0383/13/22/6775
59. Clinical Anatomy of the Lumbar Spine: A Morphometric and Radiological Approach in a South Indian Population: Observational study - Impactfactor, accessed February 21, 2026, https://impactfactor.org/PDF/IJPCR/16/IJPCR,Vol16,Issue7,Article231.pdf
60. Lumbar Root Compression in the Lateral Recess: MR Imaging, Conventional Myelography, and CT Myelography Comparison with Surgical Confirmation - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC7973614/
61. Lateral Recess Stenosis | NSPC Brain & Spine Surgery, accessed February 21, 2026, https://nspc.com/lateral-recess-stenosis/
62. The Morphometric Study of Degenerative Lateral Canal Stenosis at L4–L5 and L5–S1 Using Magnetic Resonance Imaging (MRI): Feasibility Analysis for Posterior Surgical Decompression - PMC, accessed February 21, 2026,

https://pmc.ncbi.nlm.nih.gov/articles/PMC5349339/
63. Measurement of the lateral recess angle as a possible alternative for evaluation of the lateral recess stenosis on a CT scan | Request PDF - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/publication/8212858_Measurement_of_the_lateral_recess_angle_as_a_possible_alternative_for_evaluation_of_the_lateral_recess_stenosis_on_a_CT_scan
64. Visual and quantitative assessment of lateral lumbar spinal canal stenosis with magnetic resonance imaging | Request PDF - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/publication/51691364_Visual_and_quantitative_assessment_of_lateral_lumbar_spinal_canal_stenosis_with_magnetic_resonance_imaging
65. A Practical MRI Grading System for Lumbar Foraminal Stenosis - AJR Online, accessed February 21, 2026, https://www.ajronline.org/doi/10.2214/AJR.09.2772
66. Foraminal Stenosis: What It Is, Symptoms, Types & Treatments - Cleveland Clinic, accessed February 21, 2026, https://my.clevelandclinic.org/health/diseases/24856-foraminal-stenosis
67. Foraminal Stenosis: What Causes It and How It's Treated - WebMD, accessed February 21, 2026, https://www.webmd.com/back-pain/what-is-neural-foraminal-stenosis
68. A Practical MRI Grading System for Lumbar Foraminal Stenosis - AJR, accessed February 21, 2026, https://ajronline.org/doi/abs/10.2214/AJR.09.2772
69. Clinical Correlation of a New MR Imaging Method for Assessing Lumbar Foraminal Stenosis, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC7968829/
70. Quantifying Lumbar Foraminal Volumetric Dimensions: Normative Data and Implications for Stenosis—Part 2 of a Comprehensive Series - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11270341/
71. MRI evaluation of lumbar foraminal stenosis: correlation between a new quantitative evaluation and the qualitative Lee's classification - PubMed, accessed February 21, 2026, https://pubmed.ncbi.nlm.nih.gov/40133649/
72. Thickness of the Ligamentum Flavum: Correlation with Age and Its Asymmetry–An Magnetic Resonance Imaging Study, accessed February 21, 2026, https://www.asianspinejournal.org/upload/pdf/asj-9-245.pdf?utm_source=consensus
73. Measurement of thickness of the ligamentum flavum at different spinal levels in 200 patients, accessed February 21, 2026, https://www.researchgate.net/figure/Measurement-of-thickness-of-the-ligamentum-flavum-at-different-spinal-levels-in-200_tbl1_275361309
74. Analysis of the Relationship between Ligamentum Flavum Thickening and Lumbar Segmental Instability, Disc Degeneration, and Facet Joint Osteoarthritis in Lumbar Spinal Stenosis - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5165005/
75. Is there a relationship between low back pain and ligamentum flavum

hypertrophy?, accessed February 21, 2026, https://www.europeanreview.org/wp/wp-content/uploads/3822-3827-1.pdf
76. 99% PEOPLE DON'T KNOW THIS IN LOW BACK PAIN !! - YouTube, accessed February 21, 2026, https://www.youtube.com/shorts/6MgnjA5AMKk
77. Analysis of the Thickness of Ligamentum Flavum and Its Relationship With Degenerative Disc Changes at L3-4, L4-5, and L5-S1 Levels in Patients Undergoing Magnetic Resonance Imaging (MRI) of the Lumbosacral Spine - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11663540/
78. The Role of the Ligamentum Flavum Area as a Morphological Parameter of Lumbar Central Spinal Stenosis - :::::Pain Physician:::::, accessed February 21, 2026, https://www.painphysicianjournal.com/current/pdf?article=NDMxNg%3D%3D&journal=104
79. LIGAMENTUM FLAVUM HYPERTROPHY IN ELDERLY PATIENTS WITH LOW BACK PAIN: A MRI STUDY BEL A⁄RISI BULUNAN YAfiLI HASTALARDA L‹GAM, accessed February 21, 2026, https://geriatri.dergisi.org/uploads/pdf/pdf_TJG_934.pdf
80. The “Thickened” Ligamentum Flavum: Is It Buckling or Enlargement? - PMC - NIH, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC7964003/
81. (PDF) Thickness of the Ligamentum Flavum: Correlation with Age and Its Asymmetry-An Magnetic Resonance Imaging Study - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/publication/275361309_Thickness_of_the_Ligamentum_Flavum_Correlation_with_Age_and_Its_Asymmetry-An_Magnetic_Resonance_Imaging_Study
82. Partial Spinous Process Decompression in Baastrup's Disease: A Case Report and Literature Review - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC9944191/
83. Baastrup's Sign (Kissing Spine) and Back Pain - Health Central, accessed February 21, 2026, https://www.healthcentral.com/condition/back-pain/baastrup-s-sign-kissing-spine-back-pain
84. Baastrup's Disease - Radsource, accessed February 21, 2026, https://radsource.us/baastrups_disease/
85. Baastrup's Disease: An Often Missed Etiology for Back Pain - PMC - NIH, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC4762769/
86. Lumbar Interspinous Bursitis (Baastrup Disease) in a Symptomatic Population: Prevalence on Magnetic Resonance Imaging | Request PDF - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/publication/5474044_Lumbar_Interspinous_Bursitis_Baastrup_Disease_in_a_Symptomatic_Population_Prevalence_on_Magnetic_Resonance_Imaging
87. Baastrup Disease, accessed February 21, 2026, https://cdn.mdedge.com/files/s3fs-public/Document/September-2017/036100560.pdf
88. Baastrup's disease: The kissing spine - PMC - NIH, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC3936220/

89. Radiological and Radionuclide Imaging of Degenerative Disease of the Facet Joints - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC4479906/
90. Facet Tropism/Inclination and Its Association with Intervertebral Disc Herniation in the Lumbar Spine – A Radiological Evaluation - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC9757969/
91. The correlation between radiographic and pathologic grading of lumbar facet joint degeneration - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC4812623/
92. In vivo topographic analysis of lumbar facet joint space width distribution in healthy and symptomatic subjects - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC3685171/
93. (PDF) MR imaging and CT in osteoarthritis of lumbar facet joints - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/publication/12914195_MR_imaging_and_CT_in_osteoarthritis_of_lumbar_facet_joints
94. Comparison of computed tomography and magnetic resonance imaging in the evaluation of facet tropism and facet arthrosis in degenerative cervical spondylolisthesis - Genetics and Molecular Research, accessed February 21, 2026, https://geneticsmr.com/wp-content/uploads/2024/04/gmr3355.pdf
95. Enhanced Quantitative Assessment of Lumbar Facet Joint Degeneration: Refining the Joint Degeneration Grading System With Grayscale Curve Fitting and Imaging Parameter - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC12703409/
96. Facet joint hypertrophy is a misnomer: A retrospective study - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC6024170/
97. Differential Diagnosis of Facet Joint Disorders - RSNA Journals, accessed February 21, 2026, https://pubs.rsna.org/doi/abs/10.1148/rg.2021200079
98. Kissing spine syndrome - Baastrup's disease - - Caring Medical, accessed February 21, 2026, https://caringmedical.com/prolotherapy-news/kissing-spines-syndrome-baastrups-disease/
99. Facet orientation and tropism: associations with facet joint osteoarthritis and degenerative spondylolisthesis - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC3892429/
100. Semiautomatic Assessment of Facet Tropism From Lumbar Spine MRI Using Deep Learning: A Northern Finland Birth Cohort Study - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC10997184/
101. The linear and angular dimensions of the L4-L5 facet in healthy adults: Measurements on the axial and parasagittal-oblique mri, accessed February 21, 2026, https://garj.org/articles/pdf/5575309105042015
102. RELATIONSHIP BETWEEN FACET TROPISM, LUMBAR DEGENERATION AND FACET DEGENERATION - The Journal of Turkish Spinal Surgery, accessed February 21, 2026, https://www.jtss.org/articles/relationship-between-facet-tropism-lumbar-degene

ration-and-facet-degeneration/doi/jtss.galenos.2022.54254
103. Normalizing Spinal Cord Compression Morphometric Measures: Application in Degenerative Cervical Myelopathy | medRxiv, accessed February 21, 2026, https://www.medrxiv.org/content/10.1101/2024.03.13.24304177v1.full-text
104. Deep Learning for Cervical Spine Radiography: Automated Measurement of Intervertebral and Neural Foraminal Distances - MDPI, accessed February 21, 2026, https://www.mdpi.com/2075-4418/15/17/2162
105. Pathology-Guided AI System for Accurate Segmentation and Diagnosis of Cervical Spondylosis - arXiv, accessed February 21, 2026, https://arxiv.org/html/2503.06114v1
106. Computerized Characterization of Spinal Structures on MRI and Clinical Significance of 3D Reconstruction of Lumbosacral Intervertebral Foramen - PubMed, accessed February 21, 2026, https://pubmed.ncbi.nlm.nih.gov/35051149/
107. Computerized Characterization of Spinal Structures on MRI and Clinical Significance of 3D Reconstruction of Lumbosacral Intervertebral Foramen - :::::Pain Physician:::::, accessed February 21, 2026, https://www.painphysicianjournal.com/current/pdf?article=NzM5OQ%3D%3D&journal=141
108. RSNA releases Lumbar Degenerative Imaging Spine Classification dataset - AuntMinnie, accessed February 21, 2026, https://www.auntminnie.com/clinical-news/mri/article/15814904/rsna-releases-lumbar-degenerative-imaging-spine-classification-dataset
109. [2506.09162] The RSNA Lumbar Degenerative Imaging Spine Classification (LumbarDISC) Dataset - arXiv, accessed February 21, 2026, https://arxiv.org/abs/2506.09162
110. The RSNA Lumbar Degenerative Imaging Spine Classification (LumbarDISC) Dataset, accessed February 21, 2026, https://www.researchgate.net/publication/392597545_The_RSNA_Lumbar_Degenerative_Imaging_Spine_Classification_LumbarDISC_Dataset
111. The RSNA Lumbar Degenerative Imaging Spine Classification (LumbarDISC) Dataset - PubMed, accessed February 21, 2026, https://pubmed.ncbi.nlm.nih.gov/41532835/
112. Lumbar Spine Degenerative Classification Using YOLO v8 and DeepScoreNet - medRxiv, accessed February 21, 2026, https://www.medrxiv.org/content/10.1101/2024.12.06.24318595v1.full.pdf
113. A neural network model for detection and classification of lumbar spinal stenosis on MRI, accessed February 21, 2026, https://dspace.vnmu.edu.ua/bitstream/handle/123456789/8781/%D0%9A%D0%BE%D0%BB%D1%96%D1%81%D0%BD%D0%B8%D0%BA%20%D0%A1.%20%D0%9F.%202024%20A%20neural%20network%20model%20for%20detection%20and%20classification%20of%20lumbar%20spinal%20stenosis%20on%20MRI.pdf?sequence=1&isAllowed=y
114. Radiologist-Validated Automatic Lumbar T1-Weighted Spinal MRI Segmentation Tool via an Attention U-Net Algorithm - MDPI, accessed February 21, 2026, https://www.mdpi.com/2075-4418/15/23/3046

115. Automated Measurements of Foramina on CT - American College of Radiology, accessed February 21, 2026, https://www.acr.org/Data-Science-and-Informatics/AI-in-Your-Practice/AI-Use-Cases/Use-Cases/Automated-Measurements-of-Foramina-on-CT
116. Automated Measurements of Lumbar Spinal Canal on CT - American College of Radiology, accessed February 21, 2026, https://www.acr.org/Data-Science-and-Informatics/AI-in-Your-Practice/AI-Use-Cases/Use-Cases/Auto-Measurement-Lumbar-Spinal-Canal-CT
117. Lumbar Vertebra Interpedicular Distance at L1–L5 - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/figure/Lumbar-Vertebra-Interpedicular-Distance-at-L1-L5_tbl3_235403515
118. Measuring spinal canal size in lumbar spinal stenosis: description of method and preliminary results - PMC, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC4378259/
119. Magnetic Resonance Imaging Evaluation of Biomechanical Effects of Axial Loading on the Lumbar Spine - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/publication/365168843_Magnetic_Resonance_Imaging_Evaluation_of_Biomechanical_Effects_of_Axial_Loading_on_the_Lumbar_Spine
120. (PDF) Semi-automatic CAD for upright lumbar spine MRI: reproducibility, time efficiency, and workload reduction with Q-Spine - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/publication/396665775_Semi-automatic_CAD_for_upright_lumbar_spine_MRI_reproducibility_time_efficiency_and_workload_reduction_with_Q-Spine
121. Measurement of facet tropism. | Download Scientific Diagram - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/figure/Measurement-of-facet-tropism_fig1_324682968
122. MRI grading of lumbar disc herniation based on AFFM-YOLOv8 system - PMC - NIH, accessed February 21, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC12464187/
123. Automated Segmentation and Area Estimation of Neural Foramina with Boundary Regression Model | Request PDF - ResearchGate, accessed February 21, 2026, https://www.researchgate.net/publication/308626571_Automated_Segmentation_and_Area_Estimation_of_Neural_Foramina_with_Boundary_Regression_Model