

An automatic report for the dataset : stocks-0-samsungsds

The Automatic Statistician

July 22, 2019

Abstract

This report was produced by the Automatic Bayesian Covariance Discovery (ABCD) algorithm.

1 Executive summary

The raw data and full model posterior with extrapolations are shown in figure ??.




Figure 1: Raw data (left) and model posterior with extrapolation (right)

The structure search algorithm has identified six additive components in the data. The first 3 additive components explain 94.7% of the variation in the data as shown by the coefficient of determination (R^2) values in table ???. The 6 additive components explain 100.0% of the variation in the data. After the first 4 components the cross validated mean absolute error (MAE) does not decrease by more than 0.1%. This suggests that subsequent terms are modelling very short term trends, uncorrelated noise or are artefacts of the model or search procedure. Short summaries of the additive components are as follows:

- A very smooth function. This function applies until 21 Nov 2018 and from 21 Nov 2018 onwards.
- A smooth function. This function applies until 20 Sep 2018 and from 29 Nov 2018 onwards.

- A smooth function. This function applies from 20 Sep 2018 until 21 Nov 2018 and from 21 Nov 2018 until 29 Nov 2018.
- A smooth function. This function applies from 21 Nov 2018 until 21 Nov 2018.
- A constant.
- Uncorrelated noise.

#	R^2 (%)	ΔR^2 (%)	Residual R^2 (%)	Cross validated MAE	Reduction in MAE (%)
-	-	-	-	215427.35	-
1	12.8	12.8	12.8	14416.56	93.3
2	75.3	62.5	71.6	7701.87	46.6
3	94.7	19.4	78.6	6023.01	21.8
4	98.8	4.1	76.6	5879.25	2.4
5	98.8	-0.0	-0.0	5879.23	0.0
6	100.0	1.2	100.0	5879.23	0.0

Table 1: Summary statistics for cumulative additive fits to the data. The residual coefficient of determination (R^2) values are computed using the residuals from the previous fit as the target values; this measures how much of the residual variance is explained by each new component. The mean absolute error (MAE) is calculated using 10 fold cross validation with a contiguous block design; this measures the ability of the model to interpolate and extrapolate over moderate distances. The model is fit using the full data and the MAE values are calculated using this model; this double use of data means that the MAE values cannot be used reliably as an estimate of out-of-sample predictive performance.

Model checking statistics are summarised in table ?? in section ?? . These statistics have not revealed any inconsistencies between the model and observed data.

The rest of the document is structured as follows. In section ?? the forms of the additive components are described and their posterior distributions are displayed. In section ?? the modelling assumptions of each component are discussed with reference to how this affects the extrapolations made by the model. Section ?? discusses model checking statistics, with plots showing the form of any detected discrepancies between the model and observed data.

2 Detailed discussion of additive components

2.1 Component 1 : A very smooth function. This function applies until 21 Nov 2018 and from 21 Nov 2018 onwards

This component is a very smooth function. This component applies until 21 Nov 2018 and from 21 Nov 2018 onwards.

This component explains 12.8% of the total variance. The addition of this component reduces the cross validated MAE by 93.3% from 215427.4 to 14416.6.




Figure 2: Pointwise posterior of component 1 (left) and the posterior of the cumulative sum of components with data (right)




Figure 3: Pointwise posterior of residuals after adding component 1

2.2 Component 2 : A smooth function. This function applies until 20 Sep 2018 and from 29 Nov 2018 onwards

This component is a smooth function with a typical lengthscale of 3.1 weeks. This component applies until 20 Sep 2018 and from 29 Nov 2018 onwards.

This component explains 71.6% of the residual variance; this increases the total variance explained from 12.8% to 75.3%. The addition of this component reduces the cross validated MAE by 46.58% from 14416.56 to 7701.87.




Figure 4: Pointwise posterior of component 2 (left) and the posterior of the cumulative sum of components with data (right)




Figure 5: Pointwise posterior of residuals after adding component 2

2.3 Component 3 : A smooth function. This function applies from 20 Sep 2018 until 21 Nov 2018 and from 21 Nov 2018 until 29 Nov 2018

This component is a smooth function with a typical lengthscale of 4.9 weeks. This component applies from 20 Sep 2018 until 21 Nov 2018 and from 21 Nov 2018 until 29 Nov 2018.

This component explains 78.6% of the residual variance; this increases the total variance explained from 75.3% to 94.7%. The addition of this component reduces the cross validated MAE by 21.80% from 7701.87 to 6023.01.




Figure 6: Pointwise posterior of component 3 (left) and the posterior of the cumulative sum of components with data (right)




Figure 7: Pointwise posterior of residuals after adding component 3

2.4 Component 4 : A smooth function. This function applies from 21 Nov 2018 until 21 Nov 2018

This component is a smooth function with a typical lengthscale of 2.6 days. This component applies from 21 Nov 2018 until 21 Nov 2018.

This component explains 76.6% of the residual variance; this increases the total variance explained from 94.7% to 98.8%. The addition of this component reduces the cross validated MAE by 2.39% from 6023.01 to 5879.25.




Figure 8: Pointwise posterior of component 4 (left) and the posterior of the cumulative sum of components with data (right)




Figure 9: Pointwise posterior of residuals after adding component 4

2.5 Component 5 : A constant

This component is constant.

This component explains -0.0% of the residual variance; this reduces the total variance explained from 98.8% to 98.8%. The addition of this component reduces the cross validated MAE by 0.00% from 5879.25 to 5879.23. This component neither explains residual variance nor improves MAE and therefore is likely to be an artefact of the model or search procedure.




Figure 10: Pointwise posterior of component 5 (left) and the posterior of the cumulative sum of components with data (right)




Figure 11: Pointwise posterior of residuals after adding component 5

2.6 Component 6 : Uncorrelated noise

This component models uncorrelated noise.

This component explains 100.0% of the residual variance; this increases the total variance explained from 98.8% to 100.0%. The addition of this component reduces the cross validated MAE by 0.00% from 5879.23 to 5879.23. This component explains residual variance but does not improve MAE which suggests that this component describes very short term patterns, uncorrelated noise or is an artefact of the model or search procedure.




Figure 12: Pointwise posterior of component 6 (left) and the posterior of the cumulative sum of components with data (right)

3 Extrapolation

Summaries of the posterior distribution of the full model are shown in figure ???. The plot on the left displays the mean of the posterior together with pointwise variance. The plot on the right displays three random samples from the posterior.




Figure 13: Full model posterior with extrapolation. Mean and pointwise variance (left) and three random samples (right)

Below are descriptions of the modelling assumptions associated with each additive component and how they affect the predictive posterior. Plots of the pointwise posterior and samples from the posterior are also presented, showing extrapolations from each component and the cumulative sum of components.

3.1 Component 1 : A very smooth function. This function applies until 21 Nov 2018 and from 21 Nov 2018 onwards

This component is assumed to continue very smoothly but is also assumed to be stationary so its distribution will eventually return to the prior. The prior distribution places mass on smooth functions with a marginal mean of zero and a typical lengthscale of 2.0 years. [This is a placeholder for a description of how quickly the posterior will start to resemble the prior].




Figure 14: Posterior of component 1 (top) and cumulative sum of components (bottom) with extrapolation. Mean and pointwise variance (left) and three random samples from the posterior distribution (right).

3.2 Component 2 : A smooth function. This function applies until 20 Sep 2018 and from 29 Nov 2018 onwards

This component is assumed to continue smoothly but is also assumed to be stationary so its distribution will return to the prior. The prior distribution places mass on smooth functions with a marginal mean of zero and a typical lengthscale of 3.1 weeks. [This is a placeholder for a description of how quickly the posterior will start to resemble the prior].




Figure 15: Posterior of component 2 (top) and cumulative sum of components (bottom) with extrapolation. Mean and pointwise variance (left) and three random samples from the posterior distribution (right).

3.3 Component 3 : A smooth function. This function applies from 20 Sep 2018 until 21 Nov 2018 and from 21 Nov 2018 until 29 Nov 2018

This component is assumed to stop before the end of the data and will therefore be extrapolated as zero.




Figure 16: Posterior of component 3 (top) and cumulative sum of components (bottom) with extrapolation. Mean and pointwise variance (left) and three random samples from the posterior distribution (right).

3.4 Component 4 : A smooth function. This function applies from 21 Nov 2018 until 21 Nov 2018

This component is assumed to stop before the end of the data and will therefore be extrapolated as zero.




Figure 17: Posterior of component 4 (top) and cumulative sum of components (bottom) with extrapolation. Mean and pointwise variance (left) and three random samples from the posterior distribution (right).

3.5 Component 5 : A constant

This component is assumed to stay constant.




Figure 18: Posterior of component 5 (top) and cumulative sum of components (bottom) with extrapolation. Mean and pointwise variance (left) and three random samples from the posterior distribution (right).

3.6 Component 6 : Uncorrelated noise

This component assumes the uncorrelated noise will continue indefinitely.




Figure 19: Posterior of component 6 (top) and cumulative sum of components (bottom) with extrapolation. Mean and pointwise variance (left) and three random samples from the posterior distribution (right).

4 Model checking

Several posterior predictive checks have been performed to assess how well the model describes the observed data. These tests take the form of comparing statistics evaluated on samples from the prior and posterior distributions for each additive component. The statistics are derived from autocorrelation function (ACF) estimates, periodograms and quantile-quantile (qq) plots.

Table ?? displays cumulative probability and p -value estimates for these quantities. Cumulative probabilities near 0/1 indicate that the test statistic was lower/higher under the posterior compared to the prior unexpectedly often i.e. they contain the same information as a p -value for a two-tailed test and they also express if the test statistic was higher or lower than expected. p -values near 0 indicate that the test statistic was larger in magnitude under the posterior compared to the prior unexpectedly often.

#	ACF		Periodogram		QQ	
	min	min loc	max	max loc	max	min
1	0.313	0.104	0.844	0.434	0.059	0.904
2	0.529	0.609	0.780	0.625	0.568	0.659
3	0.666	0.598	0.486	0.408	0.696	0.307
4	0.603	0.550	0.376	0.497	0.464	0.512
5	0.495	0.519	0.495	0.482	0.504	0.496
6	0.531	0.506	0.552	0.511	0.392	0.466

Table 2: Model checking statistics for each component. Cumulative probabilities for minimum of autocorrelation function (ACF) and its location. Cumulative probabilities for maximum of periodogram and its location. p -values for maximum and minimum deviations of QQ-plot from straight line.

No statistically significant discrepancies between the data and model have been detected but model checking plots for each component are presented below.

4.1 Model checking plots for components without statistically significant discrepancies

4.1.1 Component 1 : A very smooth function. This function applies until 21 Nov 2018 and from 21 Nov 2018 onwards

No discrepancies between the prior and posterior of this component have been detected




Figure 20: ACF (top left), periodogram (top right) and quantile-quantile (bottom left) uncertainty plots. The blue line and shading are the pointwise mean and 90% confidence interval of the plots under the prior distribution for component 1. The green line and green dashed lines are the corresponding quantities under the posterior.

4.1.2 Component 2 : A smooth function. This function applies until 20 Sep 2018 and from 29 Nov 2018 onwards

No discrepancies between the prior and posterior of this component have been detected




Figure 21: ACF (top left), periodogram (top right) and quantile-quantile (bottom left) uncertainty plots. The blue line and shading are the pointwise mean and 90% confidence interval of the plots under the prior distribution for component 2. The green line and green dashed lines are the corresponding quantities under the posterior.

4.1.3 Component 3 : A smooth function. This function applies from 20 Sep 2018 until 21 Nov 2018 and from 21 Nov 2018 until 29 Nov 2018

No discrepancies between the prior and posterior of this component have been detected




Figure 22: ACF (top left), periodogram (top right) and quantile-quantile (bottom left) uncertainty plots. The blue line and shading are the pointwise mean and 90% confidence interval of the plots under the prior distribution for component 3. The green line and green dashed lines are the corresponding quantities under the posterior.

4.1.4 Component 4 : A smooth function. This function applies from 21 Nov 2018 until 21 Nov 2018

No discrepancies between the prior and posterior of this component have been detected




Figure 23: ACF (top left), periodogram (top right) and quantile-quantile (bottom left) uncertainty plots. The blue line and shading are the pointwise mean and 90% confidence interval of the plots under the prior distribution for component 4. The green line and green dashed lines are the corresponding quantities under the posterior.

4.1.5 Component 5 : A constant

No discrepancies between the prior and posterior of this component have been detected




Figure 24: ACF (top left), periodogram (top right) and quantile-quantile (bottom left) uncertainty plots. The blue line and shading are the pointwise mean and 90% confidence interval of the plots under the prior distribution for component 5. The green line and green dashed lines are the corresponding quantities under the posterior.

4.1.6 Component 6 : Uncorrelated noise

No discrepancies between the prior and posterior of this component have been detected




Figure 25: ACF (top left), periodogram (top right) and quantile-quantile (bottom left) uncertainty plots. The blue line and shading are the pointwise mean and 90% confidence interval of the plots under the prior distribution for component 6. The green line and green dashed lines are the corresponding quantities under the posterior.

5 MMD - experimental section

#	mmd
1	0.000
2	0.000
3	0.000
4	0.000
5	0.794
6	0.001

Table 3: MMD p -values

5.0.1 Component 1 : A very smooth function. This function applies until 21 Nov 2018 and from 21 Nov 2018 onwards




Figure 26: MMD plot

5.0.2 Component 2 : A smooth function. This function applies until 20 Sep 2018 and from 29 Nov 2018 onwards




Figure 27: MMD plot

5.0.3 Component 3 : A smooth function. This function applies from 20 Sep 2018 until 21 Nov 2018 and from 21 Nov 2018 until 29 Nov 2018




Figure 28: MMD plot

5.0.4 Component 4 : A smooth function. This function applies from 21 Nov 2018 until 21 Nov 2018




Figure 29: MMD plot

5.0.5 Component 5 : A constant




Figure 30: MMD plot

5.0.6 Component 6 : Uncorrelated noise




Figure 31: MMD plot