

SPAN EQUIVALENCE BETWEEN WEAK N -CATEGORIES

YUYA NISHIMURA

ABSTRACT.

1. Introduction

In [Cottrell 2015], Thomas Cottrell defined an equivalence of K -algebras on an n -globular set to show the following coherence theorem:

1.1. THEOREM. *Let K be an n -globular operad with unbiased contraction γ , and let X be n -globular set. Then the free K -algebra on X is equivalent to the free strict n -category on X .*

The equivalence in this theorem is defined as follows:

1.2. DEFINITION. *Let K be an n -globular operad. K -algebras $KX \rightarrow X$ and $KY \rightarrow Y$ are equivalent if there exists a map of K -algebras $u : X \rightarrow Y$ or $u : Y \rightarrow X$ such that u is surjective on 0-cells, full on m -cells for all $1 \leq m \leq n$, and faithful on n .*

But, as he said, this equivalence is not the best one: “This definition of equivalence is much more (and thus much less general) than ought to be.” To improve it, he suggested two approaches. The one of them is to replace the map u with a span of maps of K -algebras. In this paper, we adopt this approach and prove two theorems. The first is that we define an adequate equivalence using spans and prove this is indeed an equivalence relation. The second is that our equivalence is a generalization of ordinary equivalence.

In Section 2 we recall the preliminary definitions (globular sets, their maps, operads, algebras for an operad). In Section 3 we define *span equivalence* and prove first theorem. In Section 4 we define span equivalence of ordinary categories and equivalence fusion, then we prove second theorem.

2. Preliminary

2.1. DEFINITION. *Let $n \in \mathbb{N}$. An n -globular set is a diagram*

$$X = (X_n \begin{array}{c} \xrightarrow{s_n^X} \\ \xrightarrow{t_n^X} \end{array} X_{n-1} \begin{array}{c} \xrightarrow{s_{n-1}^X} \\ \xrightarrow{t_{n-1}^X} \end{array} \dots \begin{array}{c} \xrightarrow{s_1^X} \\ \xrightarrow{t_1^X} \end{array} X_0)$$

2010 Mathematics Subject Classification: 00A00.

Key words and phrases: Weak n -category, span equivalence.

© Yuya Nishimura, 2016. Permission to copy for private use granted.

of sets and maps such that

$$s_{k-1}^X s_k^X(x) = s_{k-1}^X t_k^X(x), \quad t_{k-1}^X s_k^X(x) = t_{k-1}^X t_k^X(x)$$

for all $k \in \{2, \dots, n\}$ and $x \in X_k$.

Elements of X_k are called k -cells of X . We defined hom-sets of X as follows:

$$\mathbf{Hom}_X(x, y) := \{\alpha \in X_k \mid s_k^X(\alpha) = x, t_k^X(\alpha) = y\}$$

for all $k \in \{1, \dots, n\}$ and $x, y \in X_{k-1}$.

Let X, Y be n -globular sets, A map of n -globular sets from X to Y is a collection $f = \{f_k : X_k \rightarrow Y_k\}_{k \in \{1, \dots, n\}}$ of maps of sets such that

$$s_k^Y f_k(x) = f_{k-1} s_k^X(x), \quad t_k^Y f_k(x) = f_{k-1} t_k^X(x)$$

for all $k \in \{1, \dots, n\}$ and $x \in X_k$.

The category of n -globular sets and their maps is denoted by $n\text{-}\mathbf{GSet}$.

2.2. DEFINITION. A category is cartesian if it has all pullbacks. A functor is cartesian if it preserves pullbacks. A natural transformation is cartesian if all of its naturality squares are pullbacks squares. A monad is cartesian if its functor part, unit and counit are cartesian. A map of monad is cartesian if its underlying natural transformation is cartesian.

2.3. DEFINITION. Let \mathcal{C} be a cartesian category with a terminal object 1 . and T be a cartesian monad on \mathcal{C} . The category of T -collections is the slice category $\mathcal{C}/T1$. The category has a monoidal structure: let $k : K \rightarrow T1, k' : K' \rightarrow T1$ be collections; then their tensor product is defined to be the composite along the top of the diagram

$$\begin{array}{ccccc} K \otimes K' & \longrightarrow & TK' & \xrightarrow{Tk'} & T^2 1 \xrightarrow{\mu_1^T} T1 \\ \downarrow \lrcorner & & \downarrow T! & & \\ K & \xrightarrow{k} & T1 & & \end{array}$$

where $!$ is the unique map $K' \rightarrow 1$. the unit for this tensor product is the collection

$$\begin{array}{c} 1 \\ \downarrow \mu_1^T \\ T1 \end{array}$$

The monoidal category is denoted by $T\text{-}\mathbf{Coll}$.

2.4. DEFINITION. Let \mathcal{C} be a cartesian category with a terminal object 1 , and T be a cartesian monad on \mathcal{C} . A T -operad is a monoid in the monoidal category $T\text{-}\mathbf{Coll}$. In the case in which T is the free strict n -category monad on $n\text{-}\mathbf{GSet}$, a T -operad is called an n -globular operad.

2.5. DEFINITION. Let \mathcal{C} be a cartesian category with a terminal object 1 , T be a cartesian monad on \mathcal{C} and K be a T -operad. Then there is an induced monad on \mathcal{C} , which by abuse of notation we denote (K, η^K, μ^K) : The endfunctor

$$K : \mathcal{C} \rightarrow \mathcal{C}$$

is defined as follows; The object part of the functor, for $X \in \mathcal{C}$, KX is defined by the pullback:


$$\begin{array}{ccc} KX & \xrightarrow{K!} & K \\ k_X \downarrow & \lrcorner & \downarrow k \\ TX & \xrightarrow{T!} & T1 \end{array}$$

The arrow part of the functor, for $Y \in \mathcal{C}, u : X \rightarrow Y$, Ku is defined by the unique property of the pullback:

$$\begin{array}{ccccc} & & K! & & \\ & \curvearrowright & & \curvearrowleft & \\ KX & \xrightarrow{\quad Ku \quad} & KY & \xrightarrow{K!} & K \\ k_X \downarrow & \lrcorner & K_Y \downarrow & \lrcorner & \downarrow k \\ TX & \xrightarrow{Tu} & TY & \xrightarrow{T!} & T1 \\ & \curvearrowleft & & \curvearrowright & \\ & & T! & & \end{array}$$

Components η_X^K, μ_X^K of the unit map $\eta^K : 1 \Rightarrow K$ and $\mu^K : K^2 \Rightarrow K$ are defined by the following diagrams:

$$\begin{array}{ccccc} X & \xrightarrow{\quad ! \quad} & 1 & & \\ \eta_X^K \searrow & & \downarrow \epsilon & & \\ & KX & \xrightarrow{K!} & K & \\ \eta_X^T \searrow & k_X \downarrow & \lrcorner & \downarrow k & \eta_1^T \\ & TX & \xrightarrow{T!} & T1 & \end{array}$$



2.6. DEFINITION. Let \mathcal{C} be a cartesian category with a terminal object 1 , T be a cartesian monad on \mathcal{C} and K be a T -operad. We define a K -algebra as an algebra for the induced monad (K, η^K, μ^K) . Similarly, a map of algebras for T -operad K is a map of algebras for the induced monad. The category of K -algebras and thier maps is denoted by $K\text{-Alg}$.

Leinster's (or Batanin's) weak n -category is an algebra for specific operad. See section 9 and 10 in [Leinster 2004] for details.

3. Span equivalence

3.1. DEFINITION. Let $f : X \rightarrow Y$ be a map of n -globular sets.

- f is surjective on k -cells $\Leftrightarrow f_k : X_k \rightarrow Y_k$ is surjective
- f is injective on k -cells $\Leftrightarrow f_k : X_k \rightarrow Y_k$ is injective
- f is full on k -cells $\Leftrightarrow \begin{cases} \forall x, x' \in X_{k-1}, g \in \mathbf{Hom}_Y(f(x), f(x')), \\ \exists h \in \mathbf{Hom}_X(x, x') \text{ s.t. } f(h) = g \end{cases}$
- f is faithful on k -cell $\Leftrightarrow \begin{cases} \forall x, x' \in X_{k-1}, g, g' \in \mathbf{Hom}_X(f(x), f(x')), \\ g \neq g' \Rightarrow f(g) \neq f(g') \end{cases}$

Let f be a map of K -algebras. f is surjective (respectively, injective, full, faithful) on k -cells if the underlying map is surjective (respectively, injective, full, faithful) on k -cells.

3.2. DEFINITION. Let K be an n -globular operad. K -algebras $KX \rightarrow X$ and $KY \rightarrow Y$ are span equivalent if there exists a triple $\langle \psi, u, v \rangle$ such that $\psi : KZ \rightarrow Z$ is a K -algebra, $u : Z \rightarrow X$ and $v : Z \rightarrow Y$ are maps of K -algebras, surjective on 0 -cells, full on m -cells for all $1 \leq m \leq n$, and faithful on n -cells. The triple $\langle \psi, u, v \rangle$ is referred to as a span equivalence of K -algebras.

3.3. PROPOSITION. *In the pullback diagram in $n\text{-GSet}$*

$$\begin{array}{ccc} P & \xrightarrow{j} & Y \\ \downarrow i & \lrcorner & \downarrow g \\ X & \xrightarrow{f} & S \end{array}$$

- f is surjective on 0-cells $\Rightarrow j$ is surjective on 0-cells
- f is full on k -cells $\Rightarrow j$ is full on k -cells
- f is faithful on k -cells $\Rightarrow j$ is faithful on k -cells

PROOF. We define an n -globular set P as follows:

$$P_k := \{(x, y) \in X_k \times Y_k \mid f_k(x) = g_k(y)\}$$

$$s_l^P := (P_l \ni (x, y) \mapsto (s_l^X(x), s_l^Y(y)) \in P_{l-1})$$

$$t_l^P := (P_l \ni (x, y) \mapsto (t_l^X(x), t_l^Y(y)) \in P_{l-1})$$

for all $k \in \{0, \dots, n\}$, $l \in \{1, \dots, n\}$, and maps of n -globular sets i, j as follows:

$$i_k := (P_k \ni (x, y) \mapsto x \in X_k), \quad j_k := (P_k \ni (x, y) \mapsto y \in Y_k)$$

for all $k \in \{0, \dots, n\}$. Then (P, i, j) is a pullback of X and Y over S . It is enough to prove the proposition that we check the claims for (P, i, j) . Firstly, we prove surjectivity on 0-cells. For $y \in Y_0$, there exists $x \in X_0$ such that $f_0(x) = g_0(y)$, So $(x, y) \in P_0$ and $j_0((x, y)) = y$. which is the condition of surjectivity. To show fullness, we suppose $(x, y), (x', y') \in P_{k-1}$, $\phi \in \mathbf{Hom}(y, y')$, we can see $s_k g_k(\phi) = g_{k-1}(y) = f_{k-1}, t_k g_k(\phi) = g_{k-1}(y') = f_{k-1}(x')$. Thus $g_k(\phi) \in \mathbf{Hom}(f_{k-1}(x), f_{k-1}(x'))$. For fullness, there exists $\psi \in \mathbf{Hom}(x, x')$ such that $f_k(\psi) = g_k(\phi)$. Then $(\psi, \phi) \in \mathbf{Hom}((x, y), (x', y'))$ and $j_k(\psi, \phi) = \phi$. Therefore j is full on k -cells. Lastly, let f be faithful on k -cells. let $(x, y), (x', y') \in P_{k-1}$, $\psi, \phi \in \mathbf{Hom}((x, y), (x', y'))$ such that $j_k(\psi) = j_k(\phi)$. Then $f_k i_k(\psi) = g_k j_k(\psi) = g_k j_k(\phi) = f_k i_k(\phi)$. From faithfulness, $i_k(\psi) = i_k(\phi)$, and $\psi = (i_k(\psi), j_k(\psi)) = (i_k(\phi), j_k(\phi)) = \phi$. Therefore j is faithful on k -cells.


3.4. REMARK. Let K be a monad on $n\text{-GSet}$. Pullbacks in $K\text{-Alg}$ are created by the forgetful functor $U : K\text{-Alg} \rightarrow n\text{-GSet}$.

By the remark, we can get similar fact in category of K -algebras to proposition 3.3.

3.5. PROPOSITION. *Let*

$$\begin{array}{ccc} & P & \\ f \swarrow & & \searrow g \\ X & & Y \end{array} \qquad \begin{array}{ccc} & Q & \\ h \swarrow & & \searrow i \\ Y & & Z \end{array}$$

be span equivalences, then



is span equivalence.

PROOF. By the fact, p, q are surjective on 0-cells, full on k -cells for $1 \leq k \leq n$ and faithful on n -cells. Therefore $f \circ p, i \circ q$ are surjective on 0-cells, full on k -cells for $1 \leq k \leq n$ and faithful on n -cells. So the span is span equivalence.

3.6. THEOREM. *Span equivalence is equivalence relation on K -algebras.*

PROOF. It is straightforward from the definition and previous proposition that span equivalence is equivalence relation.

4. Characterizing equivalence of categories via spans

4.1. DEFINITION. Let \mathcal{A} and \mathcal{B} be categories. We say that \mathcal{A} and \mathcal{B} are span equivalent if there exists a triple $\langle \mathcal{A}, u, v \rangle$ such that \mathcal{C} is a category, $u : \mathcal{C} \rightarrow \mathcal{A}$ and $v : \mathcal{C} \rightarrow \mathcal{B}$ are functors, surjective on objects, full and faithful.

4.2. DEFINITION. Let \mathcal{A} and \mathcal{B} be categories, let $\langle S : \mathcal{A} \rightarrow \mathcal{B}, T : \mathcal{B} \rightarrow \mathcal{A}, \eta : I_{\mathcal{A}} \rightarrow TS, \epsilon : ST \rightarrow I_{\mathcal{B}} \rangle$ be an adjoint equivalence between \mathcal{A} and \mathcal{B} . We define a category, equivalence fusion $\mathcal{A} \Downarrow \mathcal{B}$, as follows:

- object-set

$$\mathbf{Ob}(\mathcal{A} \Downarrow \mathcal{B}) := \mathbf{Ob}(\mathcal{A}) \bigsqcup \mathbf{Ob}(\mathcal{B}) \quad (\text{disjoint})$$

- hom-set

$$\mathbf{Hom}(x, y) := \begin{cases} \{\langle f, x, y \rangle \mid f \in \mathcal{A}(x, y)\} & (x, y \in \mathcal{A}) \\ \{\langle f, x, y \rangle \mid f \in \mathcal{B}(x, y)\} & (x, y \in \mathcal{B}) \\ \{\langle f, x, y \rangle \mid f \in \mathcal{B}(Sx, y)\} & (x \in \mathcal{A}, y \in \mathcal{B}) \\ \{\langle f, x, y \rangle \mid f \in \mathcal{B}(x, Sy)\} & (x \in \mathcal{B}, y \in \mathcal{A}) \end{cases}$$

- composition

$$\begin{aligned}
 \tilde{\circ} : \mathbf{Hom}(y, z) \times \mathbf{Hom}(x, y) &\longrightarrow \mathbf{Hom}(x, z) \\
 \langle \langle g, y, z \rangle, \langle f, x, y \rangle \rangle &\longmapsto \langle g, y, z \rangle \tilde{\circ} \langle f, x, y \rangle := \langle g \circ f, x, z \rangle
 \end{aligned}$$

$$g \circ f := \begin{cases} g \circ_{\mathcal{A}} f & (x, y, z \in \mathcal{A}) \\ g \circ_{\mathcal{B}} f & (x, y, z \in \mathcal{B}) \\ g \circ_{\mathcal{B}} Sf & (x, y \in \mathcal{A}, z \in \mathcal{B}) \\ g \circ_{\mathcal{B}} f & (x \in \mathcal{A}, y, z \in \mathcal{B}) \\ g \circ_{\mathcal{B}} f & (x, y \in \mathcal{B}, z \in \mathcal{A}) \\ Sg \circ_{\mathcal{B}} f & (x \in \mathcal{B}, y, z \in \mathcal{A}) \\ \eta_z^{-1} \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x & (x \in \mathcal{A}, y \in \mathcal{B}, z \in \mathcal{A}) \\ g \circ_{\mathcal{B}} f & (x \in \mathcal{B}, y \in \mathcal{A}, z \in \mathcal{B}) \end{cases}$$

- *identities*

$$\text{id}_x := \begin{cases} \langle \text{id}_x, x, x \rangle & (x \in \mathcal{A}, \text{id}_x \in \mathcal{A}(x, x)) \\ \langle \text{id}_x, x, x \rangle & (x \in \mathcal{B}, \text{id}_x \in \mathcal{B}(x, x)) \end{cases}$$

4.3. PROPOSITION. *The equivalence fusion $\mathcal{A} \Downarrow \mathcal{B}$ forms a category.*

PROOF. It is easy to check that the composition $\tilde{\circ}$ is map from $\mathbf{Hom}(x, y) \times \mathbf{Hom}(y, z)$ to $\mathbf{Hom}(x, z)$. Now, we prove that the composition $\tilde{\circ}$ satisfies associative law and identity law by case analysis.

- associative law

$$\begin{aligned} & - x \in \mathcal{A}, y \in \mathcal{A}, z \in \mathcal{A}, w \in \mathcal{A}, \\ & \quad h \circ (g \circ f) = h \circ_{\mathcal{A}} (g \circ_{\mathcal{A}} f) \\ & \quad (h \circ g) \circ f = (h \circ_{\mathcal{A}} g) \circ_{\mathcal{A}} f \\ & - x \in \mathcal{A}, y \in \mathcal{A}, z \in \mathcal{A}, w \in \mathcal{B}, \\ & \quad h \circ (g \circ f) = h \circ (g \circ_{\mathcal{A}} f) = h \circ_{\mathcal{B}} S(g \circ_{\mathcal{A}} f) = h \circ_{\mathcal{B}} (Sg \circ_{\mathcal{B}} Sf) \\ & \quad (h \circ g) \circ f = (h \circ_{\mathcal{B}} Sg) \circ f = (h \circ_{\mathcal{B}} Sg) \circ_{\mathcal{B}} Sf \\ & - x \in \mathcal{A}, y \in \mathcal{A}, z \in \mathcal{B}, w \in \mathcal{A}, \\ & \quad h \circ (g \circ f) = h \circ (g \circ_{\mathcal{B}} Sf) = \eta_w^{-1} \circ_{\mathcal{A}} Th \circ_{\mathcal{A}} T(g \circ_{\mathcal{B}} Sf) \circ_{\mathcal{A}} \eta_x \\ & \quad \quad \quad = \eta_w^{-1} \circ_{\mathcal{A}} Th \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} TSf \circ_{\mathcal{A}} \eta_x \\ & \quad \quad \quad = \eta_w^{-1} \circ_{\mathcal{A}} Th \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} \eta_y \circ_{\mathcal{A}} f \\ & \quad (h \circ g) \circ f = (\eta_w^{-1} \circ_{\mathcal{A}} Th \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} \eta_y) \circ f = (\eta_w^{-1} \circ_{\mathcal{A}} Th \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} \eta_y) \circ_{\mathcal{A}} f \\ & - x \in \mathcal{A}, y \in \mathcal{A}, z \in \mathcal{B}, w \in \mathcal{B}, \\ & \quad h \circ (g \circ f) = h \circ (g \circ_{\mathcal{B}} Sf) = h \circ_{\mathcal{B}} (g \circ_{\mathcal{B}} Sf) \\ & \quad (h \circ g) \circ f = (h \circ_{\mathcal{B}} g) \circ f = (h \circ_{\mathcal{B}} g) \circ_{\mathcal{B}} Sf \\ & - x \in \mathcal{A}, y \in \mathcal{B}, z \in \mathcal{A}, w \in \mathcal{A}, \\ & \quad h \circ (g \circ f) = h \circ (\eta_z^{-1} \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x) = h \circ_{\mathcal{A}} \eta_z^{-1} \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x \\ & \quad \quad \quad = \eta_w^{-1} \circ_{\mathcal{A}} TSh \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x \\ & \quad (h \circ g) \circ f = (Sh \circ_{\mathcal{B}} g) \circ f = \eta_w^{-1} \circ_{\mathcal{A}} T(Sh \circ_{\mathcal{B}} g) \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x \\ & \quad \quad \quad = \eta_w^{-1} \circ_{\mathcal{A}} TSh \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x \\ & - x \in \mathcal{A}, y \in \mathcal{B}, z \in \mathcal{A}, w \in \mathcal{B}, \\ & \quad h \circ (g \circ f) = h \circ (\eta_z^{-1} \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x) \\ & \quad \quad \quad = h \circ_{\mathcal{B}} S(\eta_z^{-1} \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x) \end{aligned}$$

$$\begin{aligned}
&= h \circ_{\mathcal{B}} S\eta_z^{-1} \circ_{\mathcal{B}} ST(g \circ_{\mathcal{B}} f) \circ_{\mathcal{B}} S\eta_x \\
&= h \circ_{\mathcal{B}} (\epsilon_{Sz} \circ_{\mathcal{B}} S\eta_z) \circ_{\mathcal{B}} S\eta_z^{-1} \circ_{\mathcal{B}} ST(g \circ_{\mathcal{B}} f) \circ_{\mathcal{B}} S\eta_x \\
&= h \circ_{\mathcal{B}} \epsilon_{Sz} \circ_{\mathcal{B}} ST(g \circ_{\mathcal{B}} f) \circ_{\mathcal{B}} S\eta_x \\
&= h \circ_{\mathcal{B}} g \circ_{\mathcal{B}} f \circ_{\mathcal{B}} \epsilon_{Sx} \circ_{\mathcal{B}} S\eta_x \\
&= h \circ_{\mathcal{B}} g \circ_{\mathcal{B}} f \\
(h \circ g) \circ f &= (h \circ_{\mathcal{B}} g) \circ f = (h \circ_{\mathcal{B}} g) \circ_{\mathcal{B}} f \\
- \quad &x \in \mathcal{A}, y \in \mathcal{B}, z \in \mathcal{B}, w \in \mathcal{A}, \\
&h \circ (g \circ f) = h \circ (g \circ_{\mathcal{B}} f) = \eta_w^{-1} \circ_{\mathcal{A}} Th \circ_{\mathcal{A}} T(g \circ_{\mathcal{B}} f) \circ_{\mathcal{A}} \eta_x \\
&(h \circ g) \circ f = (h \circ_{\mathcal{B}} g) \circ f = \eta_w^{-1} \circ_{\mathcal{A}} T(h \circ_{\mathcal{B}} g) \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x \\
- \quad &x \in \mathcal{A}, y \in \mathcal{B}, z \in \mathcal{B}, w \in \mathcal{B}, \\
&h \circ (g \circ f) = h \circ_{\mathcal{B}} (g \circ_{\mathcal{B}} f) \\
&(h \circ g) \circ f = (h \circ_{\mathcal{B}} g) \circ_{\mathcal{B}} f \\
- \quad &x \in \mathcal{B}, y \in \mathcal{A}, z \in \mathcal{A}, w \in \mathcal{A}, \\
&h \circ (g \circ f) = h \circ (Sg \circ_{\mathcal{B}} f) = Sh \circ_{\mathcal{B}} (Sg \circ_{\mathcal{B}} f) \\
&(h \circ g) \circ f = (h \circ_{\mathcal{A}} g) \circ f = S(h \circ_{\mathcal{A}} g) \circ_{\mathcal{B}} f = (Sh \circ_{\mathcal{B}} Sg) \circ_{\mathcal{B}} f \\
- \quad &x \in \mathcal{B}, y \in \mathcal{A}, z \in \mathcal{A}, w \in \mathcal{B}, \\
&h \circ (g \circ f) = h \circ (Sg \circ_{\mathcal{B}} f) = h \circ_{\mathcal{B}} (Sg \circ_{\mathcal{B}} f) \\
&(h \circ g) \circ f = (h \circ_{\mathcal{B}} Sg) \circ f = (h \circ_{\mathcal{B}} Sg) \circ_{\mathcal{B}} f \\
- \quad &x \in \mathcal{B}, y \in \mathcal{A}, z \in \mathcal{B}, w \in \mathcal{A}, \\
&h \circ (g \circ f) = h \circ (g \circ_{\mathcal{B}} f) = h \circ_{\mathcal{B}} (g \circ_{\mathcal{B}} f) \\
&(h \circ g) \circ f = (\eta_w^{-1} \circ_{\mathcal{A}} Th \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} \eta_y) \circ f \\
&\quad = S(\eta_w^{-1} \circ_{\mathcal{A}} Th \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} \eta_y) \circ_{\mathcal{B}} f \\
&\quad = S\eta_w^{-1} \circ_{\mathcal{B}} ST(h \circ_{\mathcal{B}} g) \circ_{\mathcal{B}} S\eta_y \circ_{\mathcal{B}} f \\
&\quad = (\epsilon_{Sw} \circ_{\mathcal{B}} S\eta_w) \circ_{\mathcal{B}} S\eta_w^{-1} \circ_{\mathcal{B}} ST(h \circ_{\mathcal{B}} g) \circ_{\mathcal{B}} S\eta_y \circ_{\mathcal{B}} f \\
&\quad = \epsilon_{Sw} \circ_{\mathcal{B}} ST(h \circ_{\mathcal{B}} g) \circ_{\mathcal{B}} S\eta_y \circ_{\mathcal{B}} f \\
&\quad = h \circ_{\mathcal{B}} g \circ_{\mathcal{B}} \epsilon_{Sy} \circ_{\mathcal{B}} S\eta_y \circ_{\mathcal{B}} f \\
&\quad = h \circ_{\mathcal{B}} g \circ_{\mathcal{B}} f \\
- \quad &x \in \mathcal{B}, y \in \mathcal{A}, z \in \mathcal{B}, w \in \mathcal{B}, \\
&h \circ (g \circ f) = h \circ_{\mathcal{B}} (g \circ_{\mathcal{B}} f) \\
&(h \circ g) \circ f = (h \circ_{\mathcal{B}} g) \circ_{\mathcal{B}} f \\
- \quad &x \in \mathcal{B}, y \in \mathcal{B}, z \in \mathcal{A}, w \in \mathcal{A}, \\
&h \circ (g \circ f) = h \circ (g \circ_{\mathcal{B}} f) = Sh \circ_{\mathcal{B}} (g \circ_{\mathcal{B}} f) \\
&(h \circ g) \circ f = (Sh \circ_{\mathcal{B}} g) \circ f = (Sh \circ_{\mathcal{B}} g) \circ_{\mathcal{B}} f \\
- \quad &x \in \mathcal{B}, y \in \mathcal{B}, z \in \mathcal{A}, w \in \mathcal{B}, \\
&h \circ (g \circ f) = h \circ_{\mathcal{B}} (g \circ_{\mathcal{B}} f) \\
&(h \circ g) \circ f = (h \circ_{\mathcal{B}} g) \circ_{\mathcal{B}} f \\
- \quad &x \in \mathcal{B}, y \in \mathcal{B}, z \in \mathcal{B}, w \in \mathcal{A}, \\
&h \circ (g \circ f) = h \circ_{\mathcal{B}} (g \circ_{\mathcal{B}} f) \\
&(h \circ g) \circ f = (h \circ_{\mathcal{B}} g) \circ_{\mathcal{B}} f
\end{aligned}$$

- $x \in \mathcal{B}, y \in \mathcal{B}, z \in \mathcal{B}, w \in \mathcal{B}$,
 $h \circ (g \circ f) = h \circ_{\mathcal{B}} (g \circ_{\mathcal{B}} f)$
 $(h \circ g) \circ f = (h \circ_{\mathcal{B}} g) \circ_{\mathcal{B}} f$

- identity law

- $x \in \mathcal{A}, y \in \mathcal{A}$,
 $f \circ \text{id}_x = f \circ_{\mathcal{A}} \text{id}_x = f$
 $\text{id}_y \circ f = \text{id}_y \circ_{\mathcal{A}} f = f$
- $x \in \mathcal{A}, y \in \mathcal{B}$,
 $f \circ \text{id}_x = f \circ_{\mathcal{B}} S\text{id}_x = f \circ_{\mathcal{B}} \text{id}_{Sx} = f$
 $\text{id}_y \circ f = \text{id}_y \circ_{\mathcal{B}} f = f$
- $x \in \mathcal{B}, y \in \mathcal{A}$,
 $f \circ \text{id}_x = f \circ_{\mathcal{B}} \text{id}_x = f$
 $\text{id}_y \circ f = S\text{id}_y \circ_{\mathcal{B}} f = \text{id}_{Sy} \circ_{\mathcal{B}} f = f$
- $x \in \mathcal{B}, y \in \mathcal{B}$,
 $f \circ \text{id}_x = f \circ_{\mathcal{B}} \text{id}_x = f$
 $\text{id}_y \circ f = \text{id}_y \circ_{\mathcal{B}} f = f$

4.4. DEFINITION. Let $\langle S : \mathcal{A} \rightarrow \mathcal{B}, T : \mathcal{B} \rightarrow \mathcal{A}, \eta : I_{\mathcal{A}} \rightarrow TS, \epsilon : ST \rightarrow I_{\mathcal{B}} \rangle$ be an adjoint equivalence, let $\mathcal{A} \Downarrow \mathcal{B}$ be the equivalence fusion. We define the projections u, v as follows:

- $u : \mathcal{A} \Downarrow \mathcal{B} \longrightarrow \mathcal{A}$

object-function $u : \mathbf{Ob}(\mathcal{A} \Downarrow \mathcal{B}) \longrightarrow \mathbf{Ob}(\mathcal{A})$

$$x \longmapsto ux := \begin{cases} x & (x \in \mathcal{A}) \\ Tx & (x \in \mathcal{B}) \end{cases}$$

hom-functions $u : \mathbf{Hom}(x, y) \longrightarrow \mathcal{A}(ux, uy)$

$$\langle f, x, y \rangle \longmapsto uf := \begin{cases} f & (x, y \in \mathcal{A}) \\ Tf & (x, y \in \mathcal{B}) \\ Tf \circ_{\mathcal{A}} \eta_x & (x \in \mathcal{A}, y \in \mathcal{B}) \\ \eta_y^{-1} \circ_{\mathcal{A}} Tf & (x \in \mathcal{B}, y \in \mathcal{A}) \end{cases}$$

- $v : \mathcal{A} \Downarrow \mathcal{B} \longrightarrow \mathcal{B}$

object-function $v : \mathbf{Ob}(\mathcal{A} \Downarrow \mathcal{B}) \longrightarrow \mathbf{Ob}(\mathcal{B})$

$$x \longmapsto vx := \begin{cases} Sx & (x \in \mathcal{A}) \\ x & (x \in \mathcal{B}) \end{cases}$$

hom-functions $v : \mathbf{Hom}(x, y) \longrightarrow \mathcal{B}(ux, uy)$

$$\langle f, x, y \rangle \longmapsto vf := \begin{cases} Sf & (x, y \in \mathcal{A}) \\ f & (\text{others}) \end{cases}$$

4.5. PROPOSITION. The projections u, v are functors.

PROOF. We show that u, v preserve composition of morphisms and identity morphism by case analysis.

- u preserves composition of morphisms

- $x \in \mathcal{A}, y \in \mathcal{A}, z \in \mathcal{A}$,
 $u(g \circ f) = u(g \circ_{\mathcal{A}} f) = g \circ_{\mathcal{A}} f$
 $ug \circ_{\mathcal{A}} uf = g \circ_{\mathcal{A}} f$
- $x \in \mathcal{A}, y \in \mathcal{A}, z \in \mathcal{B}$,
 $u(g \circ f) = u(g \circ_{\mathcal{B}} Sf) = T(g \circ_{\mathcal{B}} Sf) \circ_{\mathcal{A}} \eta_x = Tg \circ_{\mathcal{A}} TSf \circ_{\mathcal{A}} \eta_x$
 $ug \circ_{\mathcal{A}} uf = (Tg \circ_{\mathcal{A}} \eta_y) \circ_{\mathcal{A}} f = Tg \circ_{\mathcal{A}} TSf \circ_{\mathcal{A}} \eta_x$
- $x \in \mathcal{A}, y \in \mathcal{B}, z \in \mathcal{A}$,
 $u(g \circ f) = u(\eta_z^{-1} \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x) = \eta_z^{-1} \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x$
 $ug \circ_{\mathcal{A}} uf = (\eta_z^{-1} \circ_{\mathcal{A}} Tg) \circ_{\mathcal{A}} (Tf \circ_{\mathcal{A}} \eta_x)$
- $x \in \mathcal{A}, y \in \mathcal{B}, z \in \mathcal{B}$,
 $u(g \circ f) = u(g \circ_{\mathcal{B}} f) = T(g \circ_{\mathcal{B}} f) \circ_{\mathcal{A}} \eta_x = Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x$
 $ug \circ_{\mathcal{A}} uf = Tg \circ_{\mathcal{A}} (Tf \circ_{\mathcal{A}} \eta_x)$
- $x \in \mathcal{B}, y \in \mathcal{A}, z \in \mathcal{A}$,
 $u(g \circ f) = u(Sg \circ_{\mathcal{B}} f) = \eta_z^{-1} \circ_{\mathcal{A}} T(Sg \circ_{\mathcal{B}} f) = \eta_z^{-1} \circ_{\mathcal{A}} TSg \circ_{\mathcal{B}} Tf$
 $ug \circ_{\mathcal{A}} uf = g \circ_{\mathcal{A}} (\eta_y^{-1} \circ_{\mathcal{A}} Tf) = \eta_z^{-1} \circ_{\mathcal{A}} TSg \circ_{\mathcal{A}} Tf$
- $x \in \mathcal{B}, y \in \mathcal{A}, z \in \mathcal{B}$,
 $u(g \circ f) = u(g \circ_{\mathcal{B}} f) = T(g \circ_{\mathcal{B}} f) = Tg \circ_{\mathcal{A}} Tf$
 $ug \circ_{\mathcal{A}} uf = (Tg \circ_{\mathcal{A}} \eta_y) \circ_{\mathcal{A}} (\eta_y^{-1} \circ_{\mathcal{A}} Tf) = Tg \circ_{\mathcal{A}} Tf$
- $x \in \mathcal{B}, y \in \mathcal{B}, z \in \mathcal{A}$,
 $u(g \circ f) = u(g \circ_{\mathcal{B}} f) = \eta_z^{-1} \circ_{\mathcal{A}} T(g \circ_{\mathcal{B}} f) = \eta_z^{-1} \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf$
 $ug \circ_{\mathcal{A}} uf = (\eta_z^{-1} \circ_{\mathcal{A}} Tg) \circ_{\mathcal{A}} Tf$
- $x \in \mathcal{B}, y \in \mathcal{B}, z \in \mathcal{B}$,
 $u(g \circ f) = u(g \circ_{\mathcal{B}} f) = T(g \circ_{\mathcal{B}} f) = Tg \circ_{\mathcal{A}} Tf$
 $ug \circ_{\mathcal{A}} uf = Tg \circ_{\mathcal{A}} Tf$

- u preserves identity morphisms

- $x \in \mathcal{A}$,
 $u(\text{id}_x) = \text{id}_x = \text{id}_{ux}$
- $x \in \mathcal{B}$,
 $u(\text{id}_x) = T\text{id}_x = \text{id}_{Tx} = \text{id}_{ux}$

- v preserves composition of morphisms

- $x \in \mathcal{A}, y \in \mathcal{A}, z \in \mathcal{A}$,
 $v(g \circ f) = v(g \circ_{\mathcal{A}} f) = S(g \circ_{\mathcal{A}} f) = Sg \circ_{\mathcal{B}} Sf$
 $vg \circ_{\mathcal{B}} vf = Sg \circ_{\mathcal{A}} Sf$

- $x \in \mathcal{A}, y \in \mathcal{A}, z \in \mathcal{B}$,
 $v(g \circ f) = v(g \circ_{\mathcal{B}} Sf) = g \circ_{\mathcal{B}} Sf$
 $vg \circ_{\mathcal{B}} vf = g \circ_{\mathcal{B}} Sf$
- $x \in \mathcal{A}, y \in \mathcal{B}, z \in \mathcal{A}$,
 $v(g \circ f) = v(\eta_z^{-1} \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x)$
 $= S(\eta_z^{-1} \circ_{\mathcal{A}} Tg \circ_{\mathcal{A}} Tf \circ_{\mathcal{A}} \eta_x)$
 $= S\eta_z^{-1} \circ_{\mathcal{B}} ST(g \circ_{\mathcal{B}} f) \circ_{\mathcal{B}} S\eta_x$
 $= g \circ_{\mathcal{B}} f$
 $vg \circ_{\mathcal{B}} vf = g \circ_{\mathcal{B}} f$
- $x \in \mathcal{A}, y \in \mathcal{B}, z \in \mathcal{B}$,
 $v(g \circ f) = v(g \circ_{\mathcal{B}} f) = g \circ_{\mathcal{B}} f$
 $vg \circ_{\mathcal{B}} vf = g \circ_{\mathcal{B}} f$
- $x \in \mathcal{B}, y \in \mathcal{A}, z \in \mathcal{A}$,
 $v(g \circ f) = v(Sg \circ_{\mathcal{B}} f) = Sg \circ_{\mathcal{B}} f$
 $vg \circ_{\mathcal{B}} vf = Sg \circ_{\mathcal{B}} f$
- $x \in \mathcal{B}, y \in \mathcal{A}, z \in \mathcal{B}$,
 $v(g \circ f) = v(g \circ_{\mathcal{B}} f) = g \circ_{\mathcal{B}} f$
 $vg \circ_{\mathcal{B}} vf = g \circ_{\mathcal{B}} f$
- $x \in \mathcal{B}, y \in \mathcal{B}, z \in \mathcal{A}$,
 $v(g \circ f) = v(g \circ_{\mathcal{B}} f) = g \circ_{\mathcal{B}} f$
 $vg \circ_{\mathcal{B}} vf = g \circ_{\mathcal{B}} f$
- $x \in \mathcal{B}, y \in \mathcal{B}, z \in \mathcal{B}$,
 $v(g \circ f) = v(g \circ_{\mathcal{B}} f) = g \circ_{\mathcal{B}} f$
 $vg \circ_{\mathcal{B}} vf = g \circ_{\mathcal{B}} f$

- v preserves identity morphisms

- $x \in \mathcal{A}$
 $v(\text{id}_x) = S\text{id}_x = \text{id}_{Sx} = \text{id}_{vx}$
- $x \in \mathcal{B}$
 $v(\text{id}_x) = \text{id}_x \text{id}_{vx}$

4.6. PROPOSITION. *The projections u, v are surjective on objects, full and faithful.*

PROOF. It's trivial by definitions that u, v are surjective on objects. So we check fullness and faithfulness.

- u is full and faithful

- $x, y \in \mathcal{A}$,
 $u : \mathbf{Hom}(x, y) = \{\langle f, x, y \rangle \mid f \in \mathcal{A}(x, y)\} \ni \langle f, x, y \rangle \mapsto f \in \mathcal{A}(x, y)$ is bijective.

- $x, y \in \mathcal{B}$,
 $T : \mathcal{B}(x, y) \rightarrow \mathcal{A}(Tx, Ty)$ is bijective. Therefore $u : \mathbf{Hom}(x, y) = \{\langle f, x, y \rangle \mid f \in \mathcal{B}(x, y)\} \ni \langle f, x, y \rangle \mapsto f \in \mathcal{A}(x, y) \ni \langle f, x, y \rangle \mapsto Tf \in \mathcal{A}(Tx, Ty) = \mathcal{A}(ux, uy)$ is bijective.
 - $x \in \mathcal{A}, y \in \mathcal{B}$,
 $\mathcal{B}(Sx, y) \ni f \mapsto Tf \circ_{\mathcal{A}} \eta_x \in \mathcal{A}(x, Ty)$ is the right adjunct of each f , and bijective. Therefore $u : \mathbf{Hom}(x, y) = \{\langle f, x, y \rangle \mid f \in \mathcal{B}(Sx, y)\} \ni \langle f, x, y \rangle \mapsto Tf \circ_{\mathcal{A}} \eta_x \in \mathcal{A}(x, Ty) = \mathcal{A}(ux, uy)$ is bijective.
 - $x \in \mathcal{B}, y \in \mathcal{A}$,
 $\mathcal{B}(x, Sy) \ni f \mapsto \eta_y^{-1} \circ_{\mathcal{A}} Tf \in \mathcal{A}(Tx, y)$ is the left adjunct of each f , and bijective. Therefore $u : \mathbf{Hom}(x, y) = \{\langle f, x, y \rangle \mid f \in \mathcal{B}(x, Sy)\} \ni \langle f, x, y \rangle \mapsto \eta_y^{-1} \circ_{\mathcal{A}} Tf \in \mathcal{A}(Tx, y) = \mathcal{A}(ux, uy)$
- v is full and faithful
- $x, y \in \mathcal{A}$,
 $S : \mathcal{A}(x, y) \rightarrow \mathcal{B}(Sx, Sy)$ is bijective. Therefore $v : \mathbf{Hom}(x, y) = \{\langle f, x, y \rangle \mid f \in \mathcal{A}(x, y)\} \ni \langle f, x, y \rangle \mapsto Sf \in \mathcal{B}(Sx, Sy) = \mathcal{B}(vx, vy)$ is bijective.
 - $x, y \in \mathcal{B}$,
 $v : \mathbf{Hom}(x, y) = \{\langle f, x, y \rangle \mid f \in \mathcal{B}(x, y)\} \ni \langle f, x, y \rangle \mapsto f \in \mathcal{B}(x, y) = \mathcal{B}(vx, vy)$ is bijective.
 - $x \in \mathcal{A}, y \in \mathcal{B}$,
 $v : \mathbf{Hom}(x, y) = \{\langle f, x, y \rangle \mid f \in \mathcal{B}(Sx, y)\} \ni \langle f, x, y \rangle \mapsto f \in \mathcal{B}(Sx, y) = \mathcal{B}(vx, vy)$ is bijective.
 - $x \in \mathcal{B}, y \in \mathcal{A}$,
 $v : \mathbf{Hom}(x, y) = \{\langle f, x, y \rangle \mid f \in \mathcal{B}(x, Sy)\} \ni \langle f, x, y \rangle \mapsto f \in \mathcal{B}(x, Sy) = \mathcal{B}(vx, vy)$ is bijective.

4.7. THEOREM. Let \mathcal{A} and \mathcal{B} be categories. \mathcal{A} is ordinary equivalent to \mathcal{B} if and only if \mathcal{A} is span equivalent to \mathcal{B} .

PROOF. Let \mathcal{A} be ordinary equivalent to \mathcal{B} , then \mathcal{A} is adjoint equivalent to \mathcal{B} . Thus there exists a adjoint equivalence between \mathcal{A} and \mathcal{B} . So we can construct the equivalence fusion and the projections. By Propositions, they are span equivalence. Therefore \mathcal{A} is span equivalent to \mathcal{B} .

On the other hand, let \mathcal{A} be span equivalent to \mathcal{B} . Then there exists a span equivalence $\langle \mathcal{C}, u, v \rangle$ between \mathcal{A} and \mathcal{B} , and \mathcal{C} is ordinary equivalent to both \mathcal{A} and \mathcal{B} . Therefore \mathcal{A} is ordinary equivalent to \mathcal{B} .

4.8. REMARK. Let \mathcal{A} be presheaf category. The forgetful functor

$$U : \mathcal{A}\text{-Cat} \longrightarrow \mathcal{A}\text{-Gph}$$

is monadic. (Proposition F 1.1 in [Leinster 2004])

Let $\mathcal{A} = \mathbf{Set}$, we can see $\mathbf{Set-Cat} = \mathbf{Cat}$, $\mathbf{Set-Grp} = \mathbf{1-GSet}$. and the induced monad T_1 is the free strict 1-category monad on $\mathbf{1-GSet}$. by the remark, the comparison functor

$$K : \mathbf{Cat} \longrightarrow T_1\text{-}\mathbf{Alg}$$

is isomorphic and arrow part of the functor is

$$K : f \longmapsto Uf.$$

Moreover, the category $\mathbf{Wk-1-Cat}$ of Leinster's weak 1 category is the category $T_1\text{-}\mathbf{Alg}$ of algebras for the monad. (As for details, refer to the proof of Theorem 9.1.4 in [Leinster 2004].) So the isomorphism $K : \mathbf{Cat} \rightarrow \mathbf{Wk-1-Cat}$ preserve surjectivity, fullness and faithfullness. Hence,

4.9. THEOREM. *Let $K : \mathbf{Cat} \rightarrow \mathbf{Wk-1-Cat}$ be the isomorphism above. let \mathcal{A} and \mathcal{B} be categories. \mathcal{A} is span equivalent to \mathcal{B} in \mathbf{Cat} if and only if $K(\mathcal{A})$ is span equivalent to $K(\mathcal{B})$ in $\mathbf{Wk-1-Cat}$.*

References

- [Leinster 2004] Tom Leinster, Higher operads, higher categories, volume 298 of London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 2004.
- [Cottrell 2015] Thomas Cottrell, OPERADIC DEFINITIONS OF WEAK N-CATEGORY: COHERENCE AND COMPARISONS, Theory and Applications of Categories, Vol. 30, No. 13, 2015, pp. 433-488.

Email: