

Statistical Theory - Solutions 5

Spring 2023. Email corrections to Kolyan Ray: kolyan.ray@ic.ac.uk

1. By the Neyman-Pearson lemma, this is given by the likelihood ratio test. The likelihood ratio equals

$$\Lambda(x) = \frac{f_{\sigma_1}(x)}{f_{\sigma_0}(x)} = \frac{\prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma_1^2}} e^{-x_i^2/(2\sigma_1^2)}}{\prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma_0^2}} e^{-x_i^2/(2\sigma_0^2)}} = (\sigma_0^2/\sigma_1^2)^{n/2} \exp\left(\left(\frac{1}{2\sigma_0^2} - \frac{1}{2\sigma_1^2}\right) \sum_{i=1}^n x_i^2\right),$$

and we reject H_0 if and only if $\Lambda(x) \geq k$, where $P_0(\Lambda(X) \geq k) = \alpha$. Rearranging and using that $\sigma_1^2 > \sigma_0^2$,

$$\begin{aligned} \{\Lambda(x) \geq k\} &= \left\{ \exp\left(\frac{\sigma_1^2 - \sigma_0^2}{2\sigma_0^2\sigma_1^2} \sum_{i=1}^n x_i^2\right) \geq \left(\frac{\sigma_1^2}{\sigma_0^2}\right)^{n/2} k \right\} \\ &= \left\{ \sum_{i=1}^n x_i^2 \geq \frac{2\sigma_0^2\sigma_1^2}{\sigma_1^2 - \sigma_0^2} \left(\frac{n}{2} \log(\sigma_1^2/\sigma_0^2) + \log k\right) \right\}. \end{aligned}$$

But under H_0 , $\frac{1}{\sigma_0^2} \sum_{i=1}^n X_i^2 \sim \chi_n^2$, so we know $P_0(\sum_i X_i^2 \geq \sigma_0^2 \chi_n^2(\alpha)) = \alpha$ for $\chi_n^2(\alpha)$ the upper α -tail of the χ_n^2 distribution, i.e. satisfying $P(\chi_n^2 \geq \chi_n^2(\alpha)) = \alpha$. Thus we reject H_0 if and only if $\sum_i x_i^2 \geq \sigma_0^2 \chi_n^2(\alpha)$. [You can also rearrange to find k in terms of $\chi_n^2(\alpha)$].

2. We have likelihood ratio

$$\Lambda(x) = \frac{f_2(x)}{f_1(x)} = \frac{2}{(x+2)^2} \frac{(x+1)^2}{1} = 2 \left(1 - \frac{1}{x+2}\right)^2,$$

and the LRT rejects H_0 if and only if $\Lambda(x) \geq k$. But since $x \mapsto (1 - \frac{1}{x+2})^2$ is increasing for $x > 0$, $\Lambda(x) \geq k \iff x \geq c$ for some $c = c_\alpha$ determined by the size of the test. For the type I error,

$$P_{H_0}(X \geq c) = \int_c^\infty \frac{1}{(x+1)^2} dx = \frac{1}{c+1} = \alpha,$$

i.e. $c = 1/\alpha - 1 = \frac{1-\alpha}{\alpha}$. Thus the likelihood ratio test rejects H_0 if and only if $X \geq \frac{1-\alpha}{\alpha}$. The type II error is the probability of accepting H_0 when H_1 is true:

$$P_{H_1}(X < c_\alpha) = \int_0^{c_\alpha} \frac{2}{(x+2)^2} dx = \left[-\frac{2}{x+2}\right]_0^{c_\alpha} = 1 - \frac{2}{c_\alpha + 2} = \frac{1-\alpha}{1+\alpha}.$$

Hence for $\alpha = 0.05$, the probability of a type II error is $\frac{1-0.05}{1+0.05} = 19/21$.

For testing against $H_1 : \theta > 1$, we check that the family has a monotone likelihood ratio. For $\theta_2 > \theta_1$,

$$\frac{f_{\theta_2}(x)}{f_{\theta_1}(x)} = \frac{\theta_2}{\theta_1} \left(\frac{x+\theta_1}{x+\theta_2}\right)^2 = \frac{\theta_2}{\theta_1} \left(1 - \frac{\theta_2 - \theta_1}{x+\theta_2}\right)^2,$$

which is a strictly increasing function of $x > 0$. Thus the likelihood ratio test is also uniformly most powerful in this one-sided setting and we do not change the test (see the Karlin-Rubin theorem).

3. The likelihood ratio test statistics are given by

$$\Lambda_n(x) = \frac{\sup_{\theta \in \Theta} \prod_{i=1}^n f_{\theta}(X_i)}{\prod_{i=1}^n f_{\theta_0}(X_i)} = \frac{\prod_{i=1}^n f_{\hat{\theta}_{MLE}}(X_i)}{\prod_{i=1}^n f_{\theta_0}(X_i)}$$

and can be found in the following table (see Q1 on PS2 for the MLEs):

Distribution	$\prod_{i=1}^n f_{\theta}(x_i)$	$\Lambda_n(x)$
(a) Bernoulli(θ)	$\theta^{n\bar{X}_n} (1-\theta)^{n-n\bar{X}_n}$	$(\frac{\bar{X}_n}{\theta_0})^{n\bar{X}_n} (\frac{1-\bar{X}_n}{1-\theta_0})^{n-n\bar{X}_n}$
(b) N($\theta, 1$)	$(2\pi)^{-n/2} e^{-\frac{1}{2} \sum (x_i - \theta)^2}$	$e^{\frac{1}{2} n (\bar{X}_n - \theta_0)^2}$
(c) N($0, \theta$)	$(2\pi\theta)^{-n/2} e^{-\frac{1}{2\theta} \sum x_i^2}$	$(\frac{\theta_0}{\bar{X}_n})^{n/2} e^{\frac{n}{2} (\frac{\bar{X}_n^2}{\theta_0} - 1)}$
(d) N(μ, σ^2)	$(2\pi\sigma^2)^{-n/2} e^{-\frac{1}{2\sigma^2} \sum (x_i - \mu)^2}$	$(\frac{\sigma_0^2}{\bar{\sigma}^2})^{n/2} \exp(\frac{n}{2} (\frac{\bar{\sigma}^2}{\sigma_0^2} - 1))$
(e) Poisson(θ)	$\theta^{n\bar{X}_n} e^{-n\theta} / (\prod x_i!)$	$(\frac{\bar{X}_n}{\theta_0})^{n\bar{X}_n} e^{n(\theta_0 - \bar{X}_n)}$
(f) $(1/\theta)e^{-x/\theta}$	$\frac{1}{\theta^n} e^{-n\bar{X}_n/\theta}$	$(\frac{\theta_0}{\bar{X}_n})^n \exp(n(\frac{\bar{X}_n}{\theta_0} - 1))$
(g) $\theta e^{-\theta x}$	$\theta^n e^{-n\theta \bar{X}_n}$	$(\frac{1}{\theta_0 \bar{X}_n})^n e^{n(\theta_0 \bar{X}_n - 1)}$

4. (a) For two simple hypotheses, the Neyman-Pearson lemma says the likelihood ratio test is UMP. Since

$$x \mapsto \frac{f_1(x)}{f_0(x)} = \frac{x}{1-x}$$

is an increasing function of $x \in [0, 1]$ (it has derivative $(1-x)^{-2} > 0$ on $[0, 1]$),

$$\frac{x}{1-x} \geq k \iff x \geq B$$

for some $B = B(k)$. Thus the UMP test has critical region of the desired form.

(b) The risks equal

$$\begin{aligned} R(\delta_B, H_0) &= E_{H_0} 1\{\delta_B \neq 0\} = P_{H_0}(X \geq B) = \int_B^1 2(x-1)dx = (1-B)^2, \\ R(\delta_B, H_1) &= E_{H_1} 1\{\delta_B \neq 1\} = P_{H_1}(X < B) = \int_0^B 2xdx = B^2. \end{aligned}$$

The minimax test procedure minimizes the worst case risk over the whole parameter space $\{H_0, H_1\}$, i.e. we want to minimize

$$\begin{aligned} \sup_{\theta \in \{0,1\}} R(\delta_B, H_{\theta}) &= \max\{R(\delta_B, H_0), R(\delta_B, H_1)\} = \max\{(1-B)^2, B^2\} \\ &= \begin{cases} (1-B)^2 & \text{if } B \leq 1/2, \\ B^2 & \text{if } B \geq 1/2. \end{cases} \end{aligned}$$

This is minimized at $B = 1/2$, so $\delta_{1/2}$ is the minimax test procedure.

(c) We compute the Bayes risk directly:

$$R_{\pi}(\delta_B) = E_{\theta \sim \pi} R(\delta_B, H_{\theta}) = \nu(1-B)^2 + (1-\nu)B^2.$$

Minimizing this quadratic in B yields $B(\nu) = \nu$, i.e. δ_{ν} is the Bayes test procedure. [Note that if we give more prior weight to H_0 by taking ν closer to 1, then we take $B(\nu) = \nu$ larger to reduce the chance of incorrectly rejecting H_0].

Equating the risks for $\delta_{B(\nu)} = \delta_{\nu}$,

$$(1-\nu)^2 = R(\delta_{B(\nu)}, H_0) = R(\delta_{B(\nu)}, H_1) = \nu^2,$$

which implies $\nu = 1/2$. Thus for the prior with $\nu = 1/2$, the Bayes rule $\delta_{1/2}$ has constant risk and hence is minimax. Note this is the same estimator we computed directly in (b).

5. By the CLT, $\frac{\sqrt{n}\bar{X}_n}{\sigma} \rightarrow^d N(0, 1)$. Furthermore,

$$S_n^2 = \frac{n}{n-1} \left(\frac{1}{n} \sum_{j=1}^n (X_j - \bar{X}_n)^2 \right) = \frac{n}{n-1} \left(\frac{1}{n} \sum_{j=1}^n X_j^2 - (\bar{X}_n)^2 \right) \rightarrow^p \sigma^2$$

since $\frac{1}{n} \sum_{j=1}^n X_j^2 \rightarrow^p \sigma^2$ and $\bar{X}_n \rightarrow^p 0$, both by the WLLN. Slutsky's lemma yields

$$t_n = \frac{\sqrt{n}\bar{X}_n}{S_n} \rightarrow^d N(0, 1).$$

This implies that $\frac{\sqrt{n}(\bar{X}_n - EX_1)}{S_n}$ is an asymptotically pivotal quantity, i.e. its asymptotic distribution does not depend on EX_1 . Let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of the normal distribution. Then we have

$$\lim_{n \rightarrow \infty} P(EX_1 \in [\bar{X}_n - n^{-1/2} S_n q_{1-\alpha/2}, \bar{X}_n + n^{-1/2} S_n q_{1-\alpha/2}]) = 1 - \alpha.$$

6. For $\theta_2 > \theta_1$,

$$\frac{f_{\theta_2}(x)}{f_{\theta_1}(x)} = \frac{\prod_i \theta_2 e^{-\theta_2 x_i}}{\prod_i \theta_1 e^{-\theta_1 x_i}} = (\theta_2/\theta_1)^n e^{(\theta_1 - \theta_2) \sum_i x_i}$$

is a *decreasing* function of $T(x) = \sum_i x_i$. Thus we have a monotone likelihood ratio (note the same proofs hold as long as the function is either increasing or decreasing). So by the Karlin-Rubin theorem, the UMP test is to reject H_0 if and only if $T(X) \leq k$, where $P_0(T(X) \leq k) = \alpha$. Since $T(X) \sim \Gamma(n, \theta_0)$ under H_0 , the hint gives $2\theta_0 T(X) \sim \chi_{2n}^2$ under H_0 . Therefore, take $k = \frac{1}{2\theta_0} q_{2n}(\alpha)$ the α -quantile of the χ_{2n}^2 distribution, i.e. such that $P(\chi_{2n}^2 \leq q_{2n}(\alpha)) = \alpha$.

The power function equals

$$\pi_\phi(\theta) = P_\theta(\text{reject } H_0) = P_\theta(T(X) \leq k) = P_\theta(2\theta T(X) \leq 2\theta k) = P\left(\chi_{2n}^2 \leq \frac{\theta}{\theta_0} q_{2n}(\alpha)\right).$$

We now invert the hypothesis test. Let

$$A(\theta') = \{(x_1, \dots, x_n) : \sum x_i \geq \frac{1}{2\theta'} q_{2n}(\alpha)\}$$

be the non-rejection region for $H'_0 : \theta = \theta'$. Then a $(1 - \alpha)100\%$ confidence region is

$$C(X) = \{\theta : (X_1, \dots, X_n) \in A(\theta)\} = \{\theta : \sum X_i \geq \frac{1}{2\theta} q_{2n}(\alpha)\} = \left[\frac{q_{2n}(\alpha)}{2 \sum_i X_i}, \infty \right).$$

7. (a) The log-likelihood equals

$$\ell_n(\theta, \nu) = \log \left(\prod_{i=1}^n \frac{\theta \nu^\theta}{x_i^{\theta+1}} 1_{[\nu, \infty)}(x_i) \right) = n \log \theta + n \log \nu - (\theta + 1) \log \left(\prod x_i \right) - \infty 1\{\min_i x_i < \nu\},$$

where $\infty \times 0 = 0$. For any $\theta > 0$, ℓ_n is an increasing function of ν until the indicator function becomes 1, so the log-likelihood is maximized at $\hat{\nu} = \min_i x_i$. For the MLE of θ ,

$$\begin{aligned} \frac{\partial \ell_n(\theta, \hat{\nu})}{\partial \theta} &= \frac{n}{\theta} + n \log \hat{\nu} - \log \left(\prod x_i \right) = 0 \\ \frac{\partial^2 \ell_n(\theta, \hat{\nu})}{\partial \theta^2} &= -\frac{n}{\theta^2} < 0, \end{aligned}$$

which is solved by

$$\hat{\theta} = \frac{n}{\log(\prod_i x_i) - n \log \hat{\nu}} = \frac{n}{\log(\prod_i x_i / (\min_i x_i)^n)} = \frac{n}{T}.$$

- (b) Under H_0 , the MLE for θ is $\hat{\theta}_0 = 1$ and the MLE for ν is unchanged. Since $T = \log((\prod_i x_i)/\hat{\nu}^n)$, writing the likelihood ratio statistic as a function of T ,

$$\Lambda(x) = \frac{L_n(\hat{\theta}, \hat{\nu})}{L_n(\hat{\theta}_0, \hat{\nu})} = \frac{\prod_i \frac{\hat{\theta} \hat{\nu}^{\hat{\theta}}}{x_i^{\hat{\theta}+1}}}{\prod_i \frac{\hat{\nu}}{x_i^2}} = \hat{\theta}^n \left(\frac{\hat{\nu}^n}{\prod_i x_i} \right)^{\hat{\theta}-1} = (n/T)^n (e^{-T})^{n/T-1} = \left(\frac{n}{T} \right)^n e^{T-n}.$$

Differentiating,

$$\frac{\partial}{\partial T} \log \Lambda(x) = \frac{\partial}{\partial T} (n \log n - n \log T + T - n) = 1 - n/T.$$

Hence $\Lambda(x)$ is increasing if $T \geq n$ and decreasing if $T \leq n$. So $\Lambda(x) \geq k$ is equivalent to $T \leq c_1$ or $T \geq c_2$ for suitably chosen constants c_1, c_2 .

- (c) Expanding out T :

$$T(X) = \sum_{i=1}^n [\log(X_i) - \log(\min_i X_i)].$$

We first consider the distribution of $Y_i = \log X_i$: for $e^t \geq \nu$,

$$P(\log X_i \leq t) = P(X_i \leq e^t) = \int_{\nu}^{e^t} \frac{\theta \nu^{\theta}}{x^{\theta+1}} dx = [-\nu^{\theta} x^{-\theta}]_{\nu}^{e^t} = 1 - (\nu e^{-t})^{\theta}.$$

Therefore, setting $\nu e^{-t} = e^{-s}$ gives for all $s \geq 0$,

$$P(\log(X_i/\nu) \leq s) = P(X_i \leq \nu e^{-s}) = 1 - e^{-s\theta},$$

i.e. $Y_i = \log(X_i/\nu) \sim^{iid} \text{Exp}(\theta) = \text{Exp}(1)$ since $\theta = 1$ under H_0 . We now establish the joint distribution of $Y_i - \min_j Y_j$ for $i = 1, \dots, n$. Since the Y_i are i.i.d., the minimum is attained by each Y_j with equal probability $1/n$. Suppose $Y_1 = \min_j Y_j$ attains the minimum. Then $Y_1 - \min_j Y_j = Y_1 - Y_1 = 0$ and for $t_2, \dots, t_n \geq 0$,

$$P(Y_i - \min_j Y_j \geq t_i, i = 2, \dots, n | Y_1 = \min_j Y_j) = \frac{P(Y_i - Y_1 \geq t_i, i = 2, \dots, n)}{P(Y_1 = \min_j Y_j)}.$$

Since $Y_1, \dots, Y_n \sim \text{Exp}(1)$,

$$\begin{aligned} P(Y_i - Y_1 \geq t_i, i = 2, \dots, n) &= \int_{y_1=0}^{\infty} e^{-y_1} \prod_{i=2}^n \int_{y_i=y_1+t_i}^{\infty} e^{-y_i} dy_i dy_1 \\ &= \int_{y_1=0}^{\infty} e^{-y_1} \prod_{i=2}^n e^{-(y_1+t_i)} dy_1 \\ &= e^{-\sum_{i=2}^n t_i} \int_{y_1=0}^{\infty} e^{-ny_1} dy_1 \\ &= \frac{1}{n} e^{-\sum_{i=2}^n t_i} = \frac{1}{n} \prod_{i=2}^n e^{-t_i} = \frac{1}{n} \prod_{i=2}^n P(\text{Exp}(1) \geq t_i). \end{aligned}$$

In particular,

$$P(Y_i - \min_j Y_j \geq t_i, i = 2, \dots, n | Y_1 = \min_j Y_j) = \prod_{i=2}^n e^{-t_i}.$$

Thus conditional on $Y_1 = \min_j Y_j$, we have that $(Y_1 - \min_j Y_j, \dots, Y_n - \min_j Y_j)$ has the first coordinate equal to zero and the other $i = 2, \dots, n$ coordinates equal to independent $\text{Exp}(1)$ distributions, so that

$$\sum_{i=1}^n [Y_i - \min_j Y_j] \Big| Y_1 = \min_j Y_j =^d \sum_{i=2}^n \text{Exp}(1) \sim \Gamma(n-1, 1).$$

However, the same argument applies to any of Y_2, \dots, Y_n attaining the minimum, and so $\sum_{i=1}^n Y_i - \min_j Y_j \sim \Gamma(n-1, 1)$ irrespective of which Y_i attains the minimum.

Finally, note that if $Z \sim \chi_r^2$ and $c > 0$, then $cZ \sim \Gamma(r/2, 1/(2c))$. Setting $r = 2(n-1)$ and $c = 1/2$ yields if $Z \sim \chi_{2(n-1)}^2$ then $\frac{1}{2}Z \sim \Gamma(n-1, 1)$. Setting $Z = 2T = 2 \sum_{i=1}^n [Y_i - \min_j Y_j] \sim \chi_{2(n-1)}^2$ gives the result.