

Imperial College
London

BSc, MSci and MSc EXAMINATIONS (MATHEMATICS)

May – June 2015

This paper is also taken for the relevant examination for the Associateship of the
Royal College of Science.

Statistical Theory I

Date: Monday, 11 May 2015. Time: 2.00pm – 4.00pm. Time allowed: 2 hours.

This paper has FOUR questions.

Candidates should use ONE main answer book.

Supplementary books may only be used after the main book is full.

Statistical tables are provided on pages 5 & 6.

- DO NOT OPEN THIS PAPER UNTIL THE INVIGILATOR TELLS YOU TO.
- Affix one of the labels provided to each answer book that you use, but DO NOT USE THE LABEL WITH YOUR NAME ON IT.
- Credit will be given for all questions attempted, but extra credit will be given for complete or nearly complete answers to each question as per the table below.

Raw mark	up to 12	13	14	15	16	17	18	19	20
Extra credit	0	$\frac{1}{2}$	1	$1\frac{1}{2}$	2	$2\frac{1}{2}$	3	$3\frac{1}{2}$	4

- Each question carries equal weight.
- Calculators may not be used.

1. (a) Let X denote the observed data, whose distribution depends on an unknown parameter $\theta \in \Theta$. Let $T = t(X)$ be some statistic. Write down the definition for each of the following concepts:
 - (i) T is sufficient.
 - (ii) T is minimal sufficient.
 - (iii) T is complete.
- (b) Let X_1, \dots, X_n be i.i.d. $\text{Exponential}(\theta)$ random variables.
 - (i) Show that \bar{X} is the Cramér-Rao Unbiased estimator of some function $\mu(\theta)$ of θ , and write down its variance.
 - (ii) Find a variance stabilising transformation g such that

$$\sqrt{n}(g(\bar{X}) - g(\mu)) \xrightarrow{d} \mathcal{N}(0, 1).$$
 - (iii) Show that for such $\text{Exponential}(\theta)$ observations, the $\text{Gamma}(\alpha, \beta)$ prior is a conjugate Bayesian prior.
 - (iv) Let the prior distribution of θ be $\text{Gamma}(\alpha, \beta)$. Compute the posterior mode.
2. (a) Let $\theta \in \Theta$ be an unknown parameter and let X denote the observed data. Consider the null hypothesis $H_0 : \theta \in \Theta_0$ and alternative hypothesis $H_1 : \theta \in \Theta_1 = \Theta \setminus \Theta_0$.
 - (i) Give the definitions for the size α and the power function β of a hypothesis test with critical region R . You may use the notation P_θ to denote the dependence of the probability measure on θ .
 - (ii) Give the definition of an unbiased test.
 - (iii) Explain how one can construct a $100(1 - \alpha)\%$ confidence interval for θ by first considering, for various θ_0 , size α tests of $H_0 : \theta = \theta_0$ v.s. $H_1 : \theta \neq \theta_0$.
- (b) Let Y_1, \dots, Y_n be independent with $Y_i \sim \mathcal{N}(\theta x_i, \sigma^2)$, where σ^2 and the x_i are known constants.
 - (i) Find the Cramér-Rao unbiased estimator for θ , and write down the corresponding Cramér-Rao lower bound.
 - (ii) Find an unbiased estimator of θ which is a function of $\bar{Y} = \frac{1}{n} \sum_{i=1}^n Y_i$. What is the efficiency of this estimator?
- (c) Suppose that we observe both $X \sim \text{Geometric}(1 - \theta)$ and $Y \sim \text{Poisson}(\theta)$. X and Y are independent. $\theta \in (0, 1)$ is an unknown parameter.
 - (i) Write down a minimal sufficient statistic $t(x, y)$ for θ .
 - (ii) Show that the likelihood satisfies the monotone likelihood ratio criterion.
 - (iii) Does a similar uniformly most powerful randomised test of size $\alpha = 0.01$ for testing $H_0 : \theta = 0.5$ against $H_1 : \theta < 0.5$ exist? Write one sentence justifying your answer.

3. Let X_1, \dots, X_n be i.i.d. samples with PMF $f_X(x) = \theta(1 - \theta)^x \mathbb{I}_{x \geq 0}$, and unknown parameter $\theta \in (0, 1)$. This distribution is an alternative version of the geometric distribution with range $\{0, 1, 2, \dots\}$.
- (a) Justify without proof why $S = \sum_{i=1}^n X_i$ is a complete sufficient statistic for θ .
 - (b) Find an unbiased estimator for θ^2 in the case where $n = 1$, by comparing coefficients of $(1 - \theta)^k$ in a suitable expansion.
 - (c) Compute the total score function $U_\bullet(\theta)$ and the total Fisher information $I_\bullet(\theta)$.
 - (d) Explain why there is no Cramér-Rao unbiased estimator of θ .
 - (e) $\mathbb{I}_{X_1=0}$ is an unbiased estimator for θ . Assuming $n > 1$, obtain an improved estimator by applying the Rao-Blackwell procedure using the sufficient statistic S .
[Hint: S follows a Negative-Binomial distribution with range $\{0, 1, 2, \dots\}$ because it is a sum of n Geometric random variables with ranges $\{0, 1, 2, \dots\}$.]
 - (f) Is the improved estimator obtained in (e) the minimum-variance unbiased estimator for θ ? Justify your answer.

4. Let X_1, \dots, X_n be i.i.d. samples from a $\text{Uniform}(0, \theta_X)$ distribution, and let Y_1, \dots, Y_n be independent i.i.d. samples from a $\text{Uniform}(0, \theta_Y)$ distribution. We are interested in the hypotheses $H_0 : \theta_X = \theta_Y$ and $H_1 : \theta_X \neq \theta_Y$. Let $X_{(n)} = \max(X_1, \dots, X_n)$, let $Y_{(n)} = \max(Y_1, \dots, Y_n)$ and let $T = \max(X_1, \dots, X_n, Y_1, \dots, Y_n) = \max(X_{(n)}, Y_{(n)})$.
- (a) Show that $(X_{(n)}, Y_{(n)})$ is a sufficient statistic for (θ_X, θ_Y) .
- (b) Is the hypothesis H_0 simple or composite? Is the hypothesis H_1 simple or composite? You are not required to justify your answers.
- (c) (i) Let $\Lambda = \Lambda(X_1, \dots, X_n, Y_1, \dots, Y_n) = 2 \log(\lambda)$, where $\lambda = \frac{\sup_{\theta_X, \theta_Y \in (0, \infty)} L(\theta_X, \theta_Y)}{\sup_{\theta_X = \theta_Y \in (0, \infty)} L(\theta_X, \theta_Y)}$ is the (generalised) likelihood ratio. Assuming H_0 is true, show that Λ is ancillary for $\theta = \theta_X = \theta_Y$.
- (ii) Assuming H_0 is true, show that the distribution of Λ is χ^2_2 . You may use the following facts without proof:
1. If $A \sim \chi^2_m$ is independent of $B \sim \chi^2_k$ then $A + B \sim \chi^2_{m+k}$.
 2. If A is independent of $B \sim \chi^2_k$ and $A + B \sim \chi^2_{m+k}$ then $A \sim \chi^2_m$.
 3. If $Z \sim \text{Beta}(1, \beta)$ then $-2\beta \log(Z) \sim \chi^2_2$.
 4. $\frac{X_{(n)}}{\theta_X} \sim \text{Beta}(1, n)$, independently of $\frac{Y_{(n)}}{\theta_Y} \sim \text{Beta}(1, n)$.
 5. If $\theta = \theta_X = \theta_Y$ then T is a complete sufficient statistic for θ and $\frac{T}{\theta} \sim \text{Beta}(1, 2n)$.
- [Hint: First show that T is independent of Λ .]
- (iii) Compute the critical region for a likelihood ratio test of size α of H_0 v.s. H_1 . [Hint: The χ^2_2 distribution, the Exponential $(\frac{1}{2})$ distribution and the Gamma $(1, \frac{1}{2})$ distribution are all identical.]
- (d) You may assume without loss of generality that n is large. Comment on how the distribution in (c)(ii) relates to Wilks' Theorem. Explain why this test violates the regularity conditions for Wilks' Theorem. [Hint: The regularity conditions for Wilks' Theorem are the same as those given for the asymptotic normality of maximum likelihood estimators. The regularity condition which is violated here is also one of the conditions needed to prove the Cramér-Rao lower bound.]

DISCRETE DISTRIBUTIONS							
	RANGE \mathbb{X}	PARAMETERS	MASS FUNCTION f_X	CDF F_X	$E_{f_X}[X]$	$\text{Var}_{f_X}[X]$	MGF M_X
<i>Bernoulli</i> (θ)	$\{0, 1\}$	$\theta \in (0, 1)$	$\theta^x(1 - \theta)^{1-x}$		θ	$\theta(1 - \theta)$	$1 - \theta + \theta e^t$
<i>Binomial</i> (n, θ)	$\{0, 1, \dots, n\}$	$n \in \mathbb{Z}^+, \theta \in (0, 1)$	$\binom{n}{x} \theta^x(1 - \theta)^{n-x}$		$n\theta$	$n\theta(1 - \theta)$	$(1 - \theta + \theta e^t)^n$
<i>Poisson</i> (λ)	$\{0, 1, 2, \dots\}$	$\lambda \in \mathbb{R}^+$	$\frac{e^{-\lambda} \lambda^x}{x!}$		λ	λ	$\exp\{\lambda(e^t - 1)\}$
<i>Geometric</i> (θ)	$\{1, 2, \dots\}$	$\theta \in (0, 1)$	$(1 - \theta)^{x-1} \theta$	$1 - (1 - \theta)^x$	$\frac{1}{\theta}$	$\frac{(1 - \theta)}{\theta^2}$	$\frac{\theta e^t}{1 - e^t(1 - \theta)}$
<i>NegBinomial</i> (n, θ)	$\{n, n + 1, \dots\}$	$n \in \mathbb{Z}^+, \theta \in (0, 1)$	$\binom{x-1}{n-1} \theta^n(1 - \theta)^{x-n}$		$\frac{n}{\theta}$	$\frac{n(1 - \theta)}{\theta^2}$	$\left(\frac{\theta e^t}{1 - e^t(1 - \theta)}\right)^n$
or	$\{0, 1, 2, \dots\}$	$n \in \mathbb{Z}_+^+, \theta \in (0, 1)$	$\binom{n+x-1}{x} \theta^n(1 - \theta)^x$		$\frac{n(1 - \theta)}{\theta}$	$\frac{n(1 - \theta)}{\theta^2}$	$\left(\frac{\theta}{1 - e^t(1 - \theta)}\right)^n$

For **CONTINUOUS** distributions (see over), define the **GAMMA FUNCTION**

$$\Gamma(\alpha) = \int_0^\infty x^{\alpha-1} e^{-x} dx$$

and the **LOCATION/SCALE** transformation $Y = \mu + \sigma X$ gives

$$f_Y(y) = f_X\left(\frac{y - \mu}{\sigma}\right) \frac{1}{\sigma} \quad F_Y(y) = F_X\left(\frac{y - \mu}{\sigma}\right) \quad M_Y(t) = e^{t\mu} M_X(\sigma t) \quad E_{f_Y}[Y] = \mu + \sigma E_{f_X}[X] \quad \text{Var}_{f_Y}[Y] = \sigma^2 \text{Var}_{f_X}[X]$$

CONTINUOUS DISTRIBUTIONS							
	X	PARAMS.	PDF f_X	CDF F_X	$E_{f_X}[X]$	$\text{Var}_{f_X}[X]$	MGF M_X
<i>Uniform</i> (α, β) (standard model $\alpha = 0, \beta = 1$)	(α, β)	$\alpha < \beta \in \mathbb{R}$	$\frac{1}{\beta - \alpha}$	$\frac{x - \alpha}{\beta - \alpha}$	$\frac{(\alpha + \beta)}{2}$	$\frac{(\beta - \alpha)^2}{12}$	$\frac{e^{\beta t} - e^{\alpha t}}{t(\beta - \alpha)}$
<i>Exponential</i> (λ) (standard model $\lambda = 1$)	\mathbb{R}^+	$\lambda \in \mathbb{R}^+$	$\lambda e^{-\lambda x}$	$1 - e^{-\lambda x}$	$\frac{1}{\lambda}$	$\frac{1}{\lambda^2}$	$\left(\frac{\lambda}{\lambda - t}\right)$
<i>Gamma</i> (α, β) (standard model $\beta = 1$)	\mathbb{R}^+	$\alpha, \beta \in \mathbb{R}^+$	$\frac{\beta^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\beta x}$		$\frac{\alpha}{\beta}$	$\frac{\alpha}{\beta^2}$	$\left(\frac{\beta}{\beta - t}\right)^\alpha$
<i>Weibull</i> (α, β) (standard model $\beta = 1$)	\mathbb{R}^+	$\alpha, \beta \in \mathbb{R}^+$	$\alpha \beta x^{\alpha-1} e^{-\beta x^\alpha}$	$1 - e^{-\beta x^\alpha}$	$\frac{\Gamma(1 + 1/\alpha)}{\beta^{1/\alpha}}$	$\frac{\Gamma(1 + 2/\alpha) - \Gamma(1 + 1/\alpha)^2}{\beta^{2/\alpha}}$	
<i>Normal</i> (μ, σ^2) (standard model $\mu = 0, \sigma = 1$)	\mathbb{R}	$\mu \in \mathbb{R}, \sigma \in \mathbb{R}^+$	$\frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(x - \mu)^2}{2\sigma^2}\right\}$		μ	σ^2	$e\{\mu t + \sigma^2 t^2/2\}$
<i>Student</i> (ν)	\mathbb{R}	$\nu \in \mathbb{R}^+$	$\frac{(\pi\nu)^{-1/2} \Gamma\left(\frac{\nu+1}{2}\right)}{\Gamma\left(\frac{\nu}{2}\right) \left\{1 + \frac{x^2}{\nu}\right\}^{(\nu+1)/2}}$		0 (if $\nu > 1$)	$\frac{\nu}{\nu - 2}$ (if $\nu > 2$)	
<i>Pareto</i> (θ, α)	\mathbb{R}^+	$\theta, \alpha \in \mathbb{R}^+$	$\frac{\alpha \theta^\alpha}{(\theta + x)^{\alpha+1}}$	$1 - \left(\frac{\theta}{\theta + x}\right)^\alpha$	$\frac{\theta}{\alpha - 1}$ (if $\alpha > 1$)	$\frac{\alpha \theta^2}{(\alpha - 1)(\alpha - 2)}$ (if $\alpha > 2$)	
<i>Beta</i> (α, β)	$(0, 1)$	$\alpha, \beta \in \mathbb{R}^+$	$\frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1 - x)^{\beta-1}$		$\frac{\alpha}{\alpha + \beta}$	$\frac{\alpha\beta}{(\alpha + \beta)^2(\alpha + \beta + 1)}$	