

# LINEAR ALGEBRA, MATH 50003: Lecture Notes

Lecturer: Martin Liebeck

## 1 Course Overview

Most of the course will consist of basic results on matrices, vector spaces and linear maps. The last part of the course will have a more geometrical flavour.

### 1.1 Matrix results

Let's begin with a survey of some of the highlights among the matrix results in the course. We start with a definition.

**Definition** Let  $A, B$  be  $n \times n$  matrices over a field  $F$ . We say  $A$  is *similar* to  $B$  if there exists an invertible  $n \times n$  matrix  $P$  such that  $B = P^{-1}AP$ .

Note that if we define a relation  $\sim$  on  $n \times n$  matrices by

$$A \sim B \Leftrightarrow A \text{ is similar to } B,$$

then  $\sim$  is an equivalence relation (question on Problem Sheet 1).

Two similar matrices  $A, B$  share many basic properties: for example, they have

- the same determinant
- the same characteristic polynomial
- the same eigenvalues
- the same rank
- the same trace

(question on Problem Sheet 1). One of the major aims of the subject is:

**Major Aim** For an arbitrary  $n \times n$  matrix  $A$ , find a “nice” matrix  $B$  such that  $A \sim B$ .

In the course we'll prove three famous theorems, in each of which the meaning of the word “nice” will be apparent.

**Example** Probably the nicest matrices are the diagonal ones. Recall that an  $n \times n$  matrix  $A$  is *diagonalisable* if it is similar to a diagonal matrix  $D = \text{diag}(\lambda_1, \dots, \lambda_n)$  (the diagonal matrix with diagonal entries  $\lambda_1, \dots, \lambda_n$ , the eigenvalues of  $A$ ). This property can be used to do many computations with  $A$ , such as calculating any power  $A^k$ : a matrix  $P$  such that  $D = P^{-1}AP$  can be computed (its columns are a basis of eigenvectors of  $A$ ). Then  $A = PDP^{-1}$ , so

$$A^k = (PDP^{-1})(PDP^{-1}) \cdots (PDP^{-1}) = PD^kP^{-1},$$

and  $D^k$  is the diagonal matrix  $D^k = \text{diag}(\lambda_1^k, \dots, \lambda_n^k)$ .

However, many matrices are not diagonalisable, for example

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

To see this, suppose that  $A$  is diagonalisable. Then since the only eigenvalue is 1, there exist  $P$  such that  $P^{-1}AP = \text{diag}(1, 1) = I$ , so  $A = PIP^{-1} = I$ , a contradiction.

So not every matrix can be diagonalised. However, every complex matrix can be *triangularised*. This is one of the first main results of the course:

**Triangularisation Theorem** *If  $A$  is an  $n \times n$  matrix over  $\mathbb{C}$ , then  $A$  is similar to an upper triangular matrix, i.e. there exists  $P$  such that*

$$P^{-1}AP = \begin{pmatrix} \lambda_1 & & & \\ 0 & \lambda_2 & & * \\ & & \ddots & \\ 0 & 0 & & \lambda_n \end{pmatrix}.$$

Note that this result does not hold for matrices over arbitrary fields: for example over the real numbers  $\mathbb{R}$ , the matrix  $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  has complex eigenvalues  $\pm i$ , so is not similar to a real upper triangular matrix.

The theorem has a more serious drawback though: there is nothing unique about an upper triangular matrix similar to  $A$ . For example, for any  $a, b, a', b' \neq 0$ ,

$$\begin{pmatrix} 1 & a & b \\ & 1 & 0 \\ & & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & a' & b' \\ & 1 & 0 \\ & & 1 \end{pmatrix},$$

(question on Sheet 1), so if  $A$  is similar to one such matrix, it is similar to all of them.

It is very desirable to have a *unique* matrix of a nice form that is similar to  $A$ , and that is provided by the next main result.

**Jordan Canonical Form Theorem** *If  $A$  is an  $n \times n$  matrix over  $\mathbb{C}$ , then  $A$  is similar to a matrix of the form*

$$J = \begin{pmatrix} J_1 & & \\ & J_2 & \\ & & \ddots \\ & & & J_k \end{pmatrix},$$

*a block-diagonal matrix with blocks*

$$J_i = \begin{pmatrix} \lambda_i & 1 & 0 & \dots & 0 & 0 \\ 0 & \lambda_i & 1 & \dots & 0 & 0 \\ 0 & 0 & \lambda_i & \dots & 0 & 0 \\ & & & \ddots & & \\ 0 & 0 & 0 & \dots & \lambda_i & 1 \\ 0 & 0 & 0 & \dots & 0 & \lambda_i \end{pmatrix}$$

*(these are called Jordan blocks). The collection of Jordan blocks  $J_1, \dots, J_k$  is uniquely determined by  $A$ .*

We call the matrix  $J$  the Jordan Canonical Form (JCF) of  $A$ . Its uniqueness is a vital part of the theorem, since it gives a powerful test for the similarity of two arbitrary complex matrices  $A$  and  $B$ : find the JCFs of  $A$  and  $B$ , call them  $J$  and  $J'$ . If  $J$  and  $J'$  are the same (apart from changing the order in which the Jordan blocks appear), then  $A \sim B$ ; if not, then  $A \not\sim B$ . This test can be programmed very efficiently, and can be used for huge matrices.

The Jordan Canonical Form Theorem is an ideal result for complex matrices. But what about matrices over other fields, such as  $\mathbb{R}$  or  $\mathbb{Q}$  or the finite field  $\mathbb{F}_p$  (the field of prime order  $p$  consisting of the integers  $0, 1, \dots, p-1$  with addition and multiplication modulo  $p$ )? The JCF theorem does not hold for arbitrary matrices over these fields, for the same reason that the Triangularisation theorem does not hold.

However we will prove another canonical form theorem – the Rational Canonical Form – that holds over arbitrary fields. To state this, we need a bit of notation. Let  $F$  be a field, and denote by  $F[x]$  the set of polynomials in  $x$  over  $F$ . We can add and multiply polynomials (indeed, under addition and multiplication they form what is called a *ring*).

We call a polynomial  $p(x) \in F[x]$  *monic* if it has degree  $r \geq 1$  and its leading coefficient is 1, i.e.

$$p(x) = x^r + a_{r-1}x^{r-1} + \dots + a_0. \quad (1)$$

**Definition** Let  $p(x)$  be a monic polynomial of degree  $r$  as in (1). The *companion matrix* of  $p(x)$  is the  $r \times r$  matrix  $C(p(x))$  defined as follows:

$$C(p(x)) = \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & -a_0 \\ 1 & 0 & 0 & \dots & 0 & -a_1 \\ 0 & 1 & 0 & \dots & 0 & -a_2 \\ & & & \dots & & \\ 0 & 0 & 0 & \dots & 1 & -a_{r-1} \end{pmatrix}.$$

For example,

$$C(x^3 - x + 1) = \begin{pmatrix} 0 & 0 & -1 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

Note that  $C(p(x))$  has characteristic polynomial  $p(x)$  (question on Sheet 1).

**Rational Canonical Form Theorem** Let  $A$  be an  $n \times n$  matrix over  $F$ , with characteristic polynomial  $p(x)$ .

- (i) There exists a factorization  $p(x) = p_1(x) \dots p_k(x)$  such that  $A$  is similar to a block-diagonal matrix with blocks  $C(p_i(x))$  for  $i = 1, \dots, k$ .
- (ii) Under some conditions, the polynomials  $p_1(x), \dots, p_k(x)$  are uniquely determined by  $A$ .

The “conditions” in part (ii) will be spelled out when we state and prove the theorem in the lectures.

## 1.2 Geometry

The last part of the course will be concerned with some geometrical aspects of linear algebra.

Recall the *dot product* on  $\mathbb{R}^n$ : if  $u = (u_1, \dots, u_n)$  and  $v = (v_1, \dots, v_n) \in \mathbb{R}^n$ , then

$$u.v = \sum_{i=1}^n u_i v_i.$$

Much of the geometry of  $\mathbb{R}^n$  is based on the dot product. For example, the length  $\|u\| = \sqrt{u.u}$ , and the distance between  $u$  and  $v$  is  $\|u - v\|$ . Various types of  $n \times n$  matrices fit naturally into this geometrical picture, for example

- $P$  is *orthogonal* if  $P^T P = I$  (which implies that  $Pu.Pv = u.v$  for all  $u, v$ )
- $A$  is *symmetric* if  $A^T = A$  (which implies that  $Au.v = u.Av$  for all  $u, v$ ).

It is useful to axiomatise the basic properties of the dot product, to obtain the theory of *inner product spaces*: an inner product space is a real vector space with a map sending any pair of vectors  $u, v$  to a scalar  $(u, v)$  satisfying the following axioms:

- (1) the map is linear in each variable  $u, v$
- (2) the map is symmetric, i.e.  $(v, u) = (u, v)$  for all  $u, v$
- (3)  $(u, u) > 0$  for all nonzero vectors  $u$ .

We shall develop the theory of inner product spaces. In order to extend the geometrical notions to vector spaces over arbitrary fields, we shall also develop the theory of bilinear forms.

## 2 Some revision from 1st Year Linear Algebra

This chapter is a summary of some of the theory of matrices and linear maps from the 1st year course that we'll need.

Let  $V$  be a finite dimensional vector space over a field  $F$  and  $T : V \rightarrow V$  a linear map. If  $B = \{v_1, \dots, v_n\}$  is a basis of  $V$ , let

$$\begin{aligned} T(v_1) &= a_{11}v_1 + \dots + a_{n1}v_n, \\ &\vdots \\ T(v_n) &= a_{1n}v_1 + \dots + a_{nn}v_n \end{aligned}$$

where all the coefficients  $a_{ij} \in F$ . The *matrix of  $T$  with respect to  $B$*  is

$$[T]_B = (a_{ij}) = \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{n1} & \cdots & a_{nn} \end{pmatrix}.$$

**Proposition 2.1** *Let  $S : V \rightarrow V$  and  $T : V \rightarrow V$  be linear transformations and let  $B$  be a basis of  $V$ . Then*

$$[ST]_B = [S]_B[T]_B,$$

*where  $ST$  is the composition of  $S$  and  $T$ .*

As a consequence of the proposition, the map  $T \rightarrow [T]_B$  from linear maps to  $n \times n$  matrices has many nice properties. For example, if  $[T]_B = A$  then  $[T^2]_B = A^2$  and similarly  $[T^k]_B = A^k$  for any positive integer  $k$ . More generally, for a polynomial  $q(x) = a_r x^r + \cdots + a_1 x + a_0$  ( $a_i \in F$ ), define

$$q(A) = a_r A^r + \cdots + a_1 A + a_0 I$$

and

$$q(T) = a_r T^r + \cdots + a_1 T + a_0 I_V$$

where  $I_V : V \rightarrow V$  is the identity map. Then Proposition 2.1 implies that

$$[q(T)]_B = q(A).$$

### Change of basis

Let  $V$  be  $n$ -dimensional, and let bases  $E = \{e_1, \dots, e_n\}$  and  $F = \{f_1, \dots, f_n\}$  be two bases of  $V$ . Write

$$\begin{aligned} f_1 &= p_{11}e_1 + \cdots + p_{n1}e_n, \\ &\vdots \\ f_n &= p_{1n}e_1 + \cdots + p_{nn}e_n. \end{aligned}$$

and define  $P$  to be the  $n \times n$  matrix  $(p_{ij})$ . We call  $P$  the *change of basis matrix* from  $E$  to  $F$ .

**Proposition 2.2** (i) *The change of basis matrix  $P$  is invertible.*

(ii) *If  $T : V \rightarrow V$  is a linear map, then  $[T]_F = P^{-1}[T]_E P$  (so  $[T]_E$  and  $[T]_F$  are similar matrices).*

### Determinants

As we already noted in Chapter 1, if  $A, B$  are similar  $n \times n$  matrices, then they have the same determinant. Hence if  $T : V \rightarrow V$  is a linear map, and  $E, F$  are two bases of  $V$ , then the matrices  $[T]_E$  and  $[T]_F$  have the same determinant (by Proposition 2.2(ii)). Therefore we can define the determinant  $\det(T)$  of a linear map  $T$  to be the determinant of the matrix  $[T]_E$  for any basis  $E$  of  $V$ . The *characteristic polynomial* of  $T$  is defined to be  $\det(xI_V - T)$ . This is a polynomial in  $x$  of degree  $n = \dim V$ .

**Proposition 2.3** (i) *The eigenvalues of  $T$  are the roots of the characteristic polynomial of  $T$ .*

(ii) *If  $\lambda$  is an eigenvalue of  $T$ , the eigenvectors corresponding to  $\lambda$  are the nonzero vectors in*

$$E_\lambda = \{v \in V : (\lambda I_V - T)(v) = 0\} = \ker(\lambda I_V - T).$$

(iii) *The matrix  $[T]_B$  is a diagonal matrix iff  $B$  consists of eigenvectors of  $T$ .*

**Definition** We call  $E_\lambda$  the  $\lambda$ -*eigenspace* of  $T$ . Note that  $E_\lambda$  is a subspace of  $V$  (since it is the kernel of the linear map  $\lambda I_V - T$ ).

**Proposition 2.4** *Let  $V$  a finite-dimensional vector space over  $\mathbb{C}$ , and let  $T : V \rightarrow V$  be a linear map. Then  $T$  has an eigenvalue  $\lambda \in \mathbb{C}$ .*

*Proof* The characteristic polynomial of  $T$  has a root  $\lambda \in \mathbb{C}$  by the Fundamental theorem of Algebra.  $\square$

Note that Proposition 2.4 is not necessarily true for vector spaces over other fields. For example  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by  $T(x_1, x_2) = (x_2, -x_1)$  has characteristic polynomial  $x^2 + 1$ , which has no real roots.

## Diagonalisation

Recall that a linear map  $T : V \rightarrow V$  is diagonalisable iff there exists a basis of  $V$  consisting of eigenvectors of  $T$ . Here is a very useful result on eigenvectors.

**Proposition 2.5** *Let  $T : V \rightarrow V$  be a linear map. Suppose  $v_1, \dots, v_k$  are eigenvectors of  $T$  corresponding to distinct eigenvalues  $\lambda_1, \dots, \lambda_k$ . Then  $v_1, \dots, v_k$  are linearly independent.*

**Corollary 2.6** *Let  $V$  be  $n$ -dimensional over  $F$ , and let  $T : V \rightarrow V$  a linear map. Suppose the characteristic polynomial of  $T$  has  $n$  distinct roots in  $F$ . Then  $T$  is diagonalisable.*

**Example** Let

$$A = \begin{pmatrix} \lambda_1 & & & \\ 0 & \lambda_2 & & \\ \vdots & & \ddots & \\ 0 & \dots & 0 & \lambda_n \end{pmatrix}$$

be upper triangular, with diagonal entries  $\lambda_1, \dots, \lambda_n$ , all distinct. The characteristic polynomial of  $A$  is  $\prod_{i=1}^n (x - \lambda_i)$ , which has roots  $\lambda_1, \dots, \lambda_n$ . Hence by Corollary 2.6,  $A$  is diagonalisable, so there exists  $P$  such that  $P^{-1}AP = \text{diag}(\lambda_1, \dots, \lambda_n)$ .

Note that this is not necessarily true if the diagonal entries are not distinct, e.g.  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  is not diagonalisable.

As a final point about diagonalisation, it is sometimes important to specify which field we are working over. If  $A$  is an  $n \times n$  matrix over a field  $F$ , we say  $A$  is diagonalisable over  $F$  if it is similar to a diagonal matrix with entries in  $F$ . For example, the matrix  $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  is not diagonalisable over  $\mathbb{R}$ , but it *is* diagonalisable over  $\mathbb{C}$ .

## 3 Algebraic and geometric multiplicities of eigenvalues

In this chapter we introduce and study two types of eigenvalue multiplicity.

**Definition** Let  $T : V \rightarrow V$  be a linear map with characteristic polynomial  $p(x)$ . Let  $\lambda$  be a root of  $p(x)$  (i.e. an eigenvalue of  $T$ ). Then there is a positive integer  $a(\lambda)$  such that

$$p(x) = (x - \lambda)^{a(\lambda)} q(x),$$

where  $\lambda$  is not a root of  $q(x)$ . We call  $a(\lambda)$  the *algebraic multiplicity* of  $\lambda$  as an eigenvalue of  $T$ .

The *geometric multiplicity* of  $\lambda$  is defined to be

$$g(\lambda) = \dim E_\lambda,$$

where  $E_\lambda$  is the  $\lambda$ -eigenspace of  $T$ .

We adopt similar definitions for  $n \times n$  matrices.

**Example** For  $A = \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix}$ , we have

$$a(1) = g(1) = 1, \quad a(2) = g(2) = 1.$$

And for  $B = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ , we have

$$a(1) = 2, g(1) = 1.$$

**Proposition 3.1** *If  $\lambda$  is an eigenvalue of  $T : V \rightarrow V$ , then  $g(\lambda) \leq a(\lambda)$ .*

*Proof* Let  $r = g(\lambda) = \dim E_\lambda$  and let  $v_1, \dots, v_r$  be a basis of  $E_\lambda$ . Extend to a basis of  $V$ :

$$B = \{v_1, \dots, v_r, w_1, \dots, w_s\}.$$

We work out the matrix  $[T]_B$ :

$$\begin{aligned} T(v_1) &= \lambda v_1, \\ &\vdots \\ T(v_r) &= \lambda v_r, \\ T(w_1) &= a_{11}v_1 + \dots + a_{r1}v_r + b_{11}w_1 + \dots + b_{s1}w_s, \\ &\vdots \\ T(w_s) &= a_{1s}v_1 + \dots + a_{rs}v_r + b_{1s}w_1 + \dots + b_{ss}w_s. \end{aligned}$$

So

$$[T]_B = \left( \begin{array}{cccc|ccc} \lambda & 0 & \cdots & 0 & a_{11} & \cdots & a_{1s} \\ 0 & \lambda & \cdots & 0 & \vdots & & \vdots \\ \vdots & \vdots & \ddots & \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & \lambda & a_{r1} & \cdots & a_{rs} \\ 0 & \cdots & \cdots & 0 & b_{11} & \cdots & b_{1s} \\ \vdots & & & \vdots & \vdots & & \vdots \\ \vdots & & & \vdots & \vdots & & \vdots \\ 0 & \cdots & \cdots & 0 & b_{s1} & \cdots & b_{ss} \end{array} \right) = \begin{pmatrix} \lambda I_r & A \\ 0 & B \end{pmatrix}.$$

The characteristic polynomial of this is

$$p(x) = \det \left( \begin{array}{c|c} (x - \lambda)I_r & -A \\ \hline 0 & xI_s - B \end{array} \right).$$

Using Q4 on Sheet 1, this is

$$p(x) = \det((x - \lambda)I_r) \det(xI_s - B) = (x - \lambda)^r s(x),$$

where  $s(x)$  is the characteristic polynomial of  $B$ . Hence the algebraic multiplicity  $a(\lambda) \geq r = g(\lambda)$ .  $\square$

Using this we can prove the following basic criterion for diagonalisation.

**Theorem 3.2** Let  $\dim V = n$ , let  $T : V \rightarrow V$  be a linear map, let  $\lambda_1, \dots, \lambda_r$  be the distinct eigenvalues of  $T$ , and let the characteristic polynomial of  $T$  be

$$p(x) = \prod_{i=1}^r (x - \lambda_i)^{a(\lambda_i)}$$

(so  $\sum_{i=1}^r a(\lambda_i) = n$ ). The following statements are equivalent:

- (1)  $T$  is diagonalisable.
- (2)  $\sum_{i=1}^r g(\lambda_i) = n$ .
- (3)  $g(\lambda_i) = a(\lambda_i)$  for all  $i$ .

*Proof* We first prove (1)  $\Rightarrow$  (2). Suppose (1) holds, so  $V$  has a basis  $B$  consisting of eigenvectors of  $T$ . Each vector in  $B$  is in some eigenspace  $E_{\lambda_i}$ , so

$$\sum_{i=1}^r g(\lambda_i) = \sum_{i=1}^r \dim E_{\lambda_i} \geq |B| = n.$$

By 3.1,  $\sum_{i=1}^r g(\lambda_i) \leq \sum_{i=1}^r a(\lambda_i) = n$ . Hence  $\sum g(\lambda_i) = n$ .

Next we show that (2)  $\Leftrightarrow$  (3). This is easy, as

$$\sum g(\lambda_i) = n \Leftrightarrow \sum g(\lambda_i) = \sum a(\lambda_i) \Leftrightarrow g(\lambda_i) = a(\lambda_i) \forall i$$

(using 3.1 for the last implication).

To complete the proof, we show that (2)  $\Rightarrow$  (1). Suppose (2) holds, so  $\sum_{i=1}^r \dim E_{\lambda_i} = n$ . Let  $B_i$  be a basis of  $E_{\lambda_i}$  and let  $B = \bigcup_{i=1}^r B_i$ , so  $|B| = n$  (the  $B_i$ 's are disjoint as they consist of eigenvectors for different eigenvalues).

We claim that  $B$  is a basis of  $V$  (hence (1) holds). Since  $|B| = n = \dim V$ , it is enough to show that  $B$  is linearly independent. Suppose there is a linear relation on the vectors in  $B$ , and write it as

$$\sum_{a \in B_1} \alpha_a a + \dots + \sum_{z \in B_r} \alpha_z z = 0. \quad (2)$$

Write

$$\begin{aligned} v_1 &= \sum_{a \in B_1} \alpha_a a, \\ &\vdots \\ v_r &= \sum_{z \in B_r} \alpha_z z, \end{aligned}$$

so  $v_i \in E_{\lambda_i}$  and  $v_1 + \dots + v_r = 0$ . As  $\lambda_1, \dots, \lambda_r$  are distinct, the set of nonzero  $v_i$ 's is linearly independent by 2.5. Therefore there can't be any nonzero  $v_i$ 's, and so  $v_i = 0$  for all  $i$ . Then  $v_1 = \sum_{a \in B_1} \alpha_a a = 0$ , so as  $B_1$  is linearly independent (it is a basis of  $E_{\lambda_1}$ ) all the coefficients  $\alpha_a = 0$ . Similarly all the other  $\alpha$ 's in (2) are 0. This completes the proof that  $B$  is linearly independent, hence a basis of  $V$ .  $\square$

Using 3.2 we obtain a test to check whether a given  $n \times n$  matrix or linear map is diagonalisable:

1. Find the characteristic polynomial, and factorise it as

$$\prod_{i=1}^r (x - \lambda_i)^{a(\lambda_i)}.$$



2. Calculate each  $g(\lambda_i) = \dim E_{\lambda_i}$ .
3. If  $g(\lambda_i) = a(\lambda_i)$  for all  $i$ , YES.  
If  $g(\lambda_i) < a(\lambda_i)$  for some  $i$ , NO.

**Example** Let  $A = \begin{pmatrix} -3 & 1 & -1 \\ -7 & 5 & -1 \\ -6 & 6 & -2 \end{pmatrix}$ . Check that

- (1) Characteristic polynomial is  $(x+2)^2(x-4)$ .
- (2) For eigenvalue 4:  $a(4) = 1, g(4) = 1$  (as it is  $\leq a(4)$ ).  
For eigenvalue  $-2$ :  $a(-2) = 2, g(-2) = \dim E_{-2} = 1$ .

So  $g(-2) < a(-2)$  and  $A$  is not diagonalisable.

## 4 Direct sums

Recall that if  $U_1, \dots, U_k$  are subspaces of a vector space  $V$ , we can form their *sum*

$$U_1 + \dots + U_k = \{u_1 + \dots + u_k : u_i \in U_i \text{ for all } i\},$$

which is another subspace of  $V$ . A *direct sum* of subspaces is a particular case of this, defined as follows.

**Definition** Let  $V$  be a vector space, and let  $V_1, \dots, V_k$  be subspaces of  $V$ . We write

$$V = V_1 \oplus V_2 \oplus \dots \oplus V_k \tag{3}$$

if every vector  $v \in V$  can be expressed as  $v = v_1 + \dots + v_k$  for *unique* vectors  $v_i \in V_i$ . The uniqueness statement means that if  $v_1 + \dots + v_k = v'_1 + \dots + v'_k$  with  $v_i, v'_i \in V_i$ , then  $v_i = v'_i$  for all  $i$ . If (3) holds, we say that  $V$  is the *direct sum* of the subspaces  $V_1, \dots, V_k$ .

As an obvious first example,  $\mathbb{R}^2 = \text{Sp}(1, 0) \oplus \text{Sp}(0, 1)$ . (Here, and throughout these notes, “Sp” is an abbreviation for “Span”.)

It will be important for us to be able to check quickly whether the direct sum condition (3) holds. For a direct sum of two subspaces (the case  $k = 2$ ), this is easy:

**Proposition 4.1** *The following statements are equivalent:*

- (1)  $V = V_1 \oplus V_2$ .
- (2)  $V_1 \cap V_2 = \{0\}$  and  $\dim V_1 + \dim V_2 = \dim V$ .

*Proof* First we show (1)  $\Rightarrow$  (2). Assume (1), so that  $V = V_1 \oplus V_2$ . If there exists  $0 \neq v \in V_1 \cap V_2$ , then

$$v = v + 0 = 0 + v$$

gives two different expressions for  $v$  as a sum of vectors in  $V_1$  and  $V_2$ , contradicting the uniqueness statement in the definition of a direct sum. Therefore  $V_1 \cap V_2 = \{0\}$ . It follows that

$$\dim V = \dim(V_1 + V_2) = \dim V_1 + \dim V_2 - \dim V_1 \cap V_2 = \dim V_1 + \dim V_2.$$

Hence (2) holds.

Now we show (2)  $\Rightarrow$  (1). Assume that (2) holds. Then

$$\dim(V_1 + V_2) = \dim V_1 + \dim V_2 - \dim V_1 \cap V_2 = \dim V_1 + \dim V_2 = \dim V.$$

Hence  $V = V_1 + V_2$ . To show uniqueness, suppose  $v_1 + v_2 = v'_1 + v'_2$  with  $v_i, v'_i \in V_i$ . Then

$$v_1 - v'_1 = v'_2 - v_2 \in V_1 \cap V_2.$$

Since  $V_1 \cap V_2 = \{0\}$ , this implies that  $v_1 = v'_1, v_2 = v'_2$ . Hence  $V = V_1 \oplus V_2$ .  $\square$

The next result shows how to check the direct sum condition (3) for arbitrary values of  $k$ .

**Proposition 4.2** *The following statements are equivalent:*

- (1)  $V = V_1 \oplus \cdots \oplus V_k$ .
- (2)  $\dim V = \sum_{i=1}^k \dim V_i$ , and if  $B_i$  is a basis for  $V_i$  for  $1 \leq i \leq k$ , then  $B = B_1 \cup \cdots \cup B_k$  is a basis of  $V$ .

*Proof* First we prove (1)  $\Rightarrow$  (2). Assume that  $V = V_1 \oplus \cdots \oplus V_k$ . Let  $B_i$  be a basis of  $V_i$  for  $1 \leq i \leq k$ , and let  $B = B_1 \cup \cdots \cup B_k$ .

**Claim**  $B$  is a basis of  $V$ .

Proof of Claim: Clearly  $B$  spans  $V$ , since  $V = V_1 + \cdots + V_k$ . Now we show linear independence. Suppose there is a linear relation on the vectors in  $B$ , and write this as

$$\sum_{a \in B_1} \alpha_a a + \cdots + \sum_{z \in B_r} \alpha_z z = 0. \quad (4)$$

Now  $V = V_1 \oplus \cdots \oplus V_k$ , hence  $0 = 0 + \cdots + 0$  is the *unique* expression for the zero vector as a sum of vectors in  $V_1, \dots, V_k$ . Hence each sum in the left hand side of (4) is equal to 0, and so all the  $\alpha$ 's in (4) are 0. This proves that  $B$  is linearly independent, hence is a basis, proving the Claim.

As in the proof of 4.1 we see that  $V_i \cap V_j = \{0\}$  for  $i \neq j$ , and hence  $B_i \cap B_j = \emptyset$  and  $B$  is the disjoint union of the  $B_i$ . By the Claim, therefore, we have

$$\dim V = |B| = \sum_{i=1}^k |B_i| = \sum_{i=1}^k \dim V_i,$$

so that (2) holds.

Now we prove that (2)  $\Rightarrow$  (1). Assume that (2) holds. For each  $i$  let  $B_i$  be a basis of  $V_i$ , and let  $B = \bigcup_1^k B_i$ , a basis of  $V$ . As  $\dim V = \sum_1^k \dim V_i$ , we have  $|B| = \sum |B_i|$ , so the  $B_i$ 's are disjoint sets. Every vector in  $V$  is in the span of  $B$ , hence is a sum of vectors in  $V_1, \dots, V_k$ , so  $V = V_1 + \cdots + V_k$ . To prove uniqueness, suppose that

$$v_1 + \cdots + v_k = v'_1 + \cdots + v'_k$$

where each  $v_i, v'_i \in V_i$ . Then

$$0 = (v_1 - v'_1) + \cdots + (v_k - v'_k).$$

If any term  $v_i - v'_i$  is nonzero, this equation will give a nontrivial linear relation on the vectors in the basis  $B$ , a contradiction. Hence  $v_i = v'_i$  for all  $i$ , proving uniqueness, and so  $V = V_1 \oplus \cdots \oplus V_k$ .  $\square$

**Example** In  $\mathbb{R}^4$  let  $V_1 = \text{sp}((1, 1, 0, 0), (0, -1, 1, 0))$ ,  $V_2 = \text{sp}(2, 1, 2, 1)$ ,  $V_3 = \text{sp}(0, 0, 1, 1)$ . Is  $\mathbb{R}^4 = V_1 \oplus V_2 \oplus V_3$ ?

Answer: no, as  $\{(1, 1, 0, 0), (0, -1, 1, 0), (2, 1, 2, 1), (0, 0, 1, 1)\}$  is not a basis of  $\mathbb{R}^4$ . (The simplest way to check this is to write the vectors as the rows of a  $4 \times 4$  matrix and show that this can be reduced by row operations to a matrix with a zero row.)

To complete this chapter, we demonstrate an important link between direct sums and linear maps. First we need a definition.

**Definition** Let  $T : V \rightarrow V$  be a linear map, and  $W$  a subspace of  $V$ . We say that  $W$  is *T-invariant* if  $T(W) \subseteq W$ , where  $T(W) = \{T(w) : w \in W\}$  (in other words,  $T$  maps  $W \rightarrow W$ ). If  $W$  is  $T$ -invariant, write  $T_W : W \rightarrow W$  for the *restriction* of  $T$  to  $W$ . Thus  $T_W$  is the linear map  $W \rightarrow W$  defined by  $T_W(w) = T(w)$  for all  $w \in W$ .

**Proposition 4.3** Let  $T : V \rightarrow V$  be a linear map, and suppose that  $V = V_1 \oplus \cdots \oplus V_k$ , where each subspace  $V_i$  is  $T$ -invariant. For each  $i$  let  $B_i$  be a basis of  $V_i$ , and let  $A_i$  be the matrix of the restriction  $[T_{V_i}]_{B_i}$ . Then if  $B$  is the basis  $\bigcup_1^k B_i$  of  $V$ , the matrix  $[T]_B$  is the block-diagonal matrix

$$[T]_B = \begin{pmatrix} A_1 & & \\ & \ddots & \\ & & A_k \end{pmatrix}. \quad (5)$$

**Proof** Let  $B_1 = \{v_1, \dots, v_r\}$ . Then  $T(v_1) = T_{V_1}(v_1)$  is a vector in  $V_1$ , say  $T(v_1) = a_{11}v_1 + \cdots + a_{r1}v_r$ . Similarly for  $T(v_2), \dots$ , up to  $T(v_r) = T_{V_1}(v_r) = a_{1r}v_1 + \cdots + a_{rr}v_r$ . So we see that the top left hand block of  $[T]_B$  is the  $r \times r$  matrix  $(a_{ij})$ , which is  $[T_{V_1}]_{B_1}$ . Carrying on like this, we see that the next diagonal block is  $[T_{V_2}]_{B_2}$ , and so on.  $\square$

**Notation** In view of the proposition, and for convenience of notation, we shall denote the block-diagonal matrix in (5) by  $A_1 \oplus \cdots \oplus A_k$ . Thus for  $n_i \times n_i$  matrices  $A_i$  ( $1 \leq i \leq k$ ), we write

$$A_1 \oplus \cdots \oplus A_k = \begin{pmatrix} A_1 & & \\ & \ddots & \\ & & A_k \end{pmatrix},$$

an  $n \times n$  block-diagonal matrix, where  $n = \sum_{i=1}^k n_i$ .

## 5 Quotient spaces

Let  $V$  be a vector space over a field  $F$ , and  $W$  a subspace of  $V$ . In this section we define the *quotient space*  $V/W$ . Its vectors are the *cosets*  $W + v$  for  $v \in V$ , where

$$W + v = \{w + v : w \in W\}.$$

These are just cosets of the additive subgroup  $W$  of the group  $(V, +)$ , as seen in 1st Year Group Theory. (They are *right* cosets, but the right coset  $W + v$  is the same as

the left coset  $v + W$  because addition is commutative, so we just call them cosets.) It is of course possible to have  $W + v = W + v'$  for different vectors  $v, v'$ ; it is easy to tell when this happens:

$$W + v = W + v' \Leftrightarrow v - v' \in W.$$

You will have seen this fact in the 1st Year, but I have also set it as a question on Sheet 2 to make sure.

To make  $V/W$  into a vector space, we need to define addition and scalar multiplication of cosets. The natural definitions are:

$$(A) \quad (W + v_1) + (W + v_2) = W + v_1 + v_2$$

$$(S) \quad \lambda(W + v) = W + \lambda v$$

for all  $v_i, v \in V$ ,  $\lambda \in F$ . We must check that these operations are well-defined. Here is the check for (A):

$$\begin{aligned} W + v_1 = W + v'_1, W + v_2 = W + v'_2 &\Rightarrow v_1 - v'_1, v_2 - v'_2 \in W \\ &\Rightarrow v_1 + v_2 - (v'_1 + v'_2) \in W \\ &\Rightarrow W + v_1 + v_2 = W + v'_1 + v'_2. \end{aligned}$$

And here is the check for (S):

$$\begin{aligned} W + v = W + v' &\Rightarrow v - v' \in W \\ &\Rightarrow \lambda(v - v') \in W \\ &\Rightarrow \lambda v - \lambda v' \in W \\ &\Rightarrow W + \lambda v = W + \lambda v'. \end{aligned}$$

**Proposition 5.1** *Let  $V/W$  be the set of cosets  $W + v$  for  $v \in V$ . Then with addition and scalar multiplication defined by (A) and (S) as above,  $V/W$  is a vector space over  $F$ .*

*Proof.* We need to check the vector space axioms for  $V/W$ . These are:

Addition axioms: these amount to saying that  $(V/W, +)$  is an abelian group, with identity element the zero vector  $W + 0 = W$ .

Scalar multiplication axioms – these are

$$(S1) \quad \lambda((W + v_1) + (W + v_2)) = \lambda(W + v_1) + \lambda(W + v_2)$$

$$(S2) \quad (\lambda + \mu)(W + v) = \lambda(W + v) + \mu(W + v)$$

$$(S3) \quad (\lambda(\mu)(W + v)) = (\lambda\mu)(W + v)$$

$$(S4) \quad 1(W + v) = W + v.$$

Checking all the axioms is a routine exercise. I will just do (S1) and leave the rest to you to check:

$$\begin{aligned} \lambda((W + v_1) + (W + v_2)) &= \lambda(W + v_1 + v_2) \\ &= W + \lambda(v_1 + v_2) \\ &= W + \lambda v_1 + \lambda v_2 \\ &= (W + \lambda v_1) + (W + \lambda v_2) \\ &= \lambda(W + v_1) + \lambda(W + v_2). \quad \square \end{aligned}$$

We call the vector space  $V/W$  the *quotient space* of  $V$  by  $W$ . Its dimension is given by the next result.

**Proposition 5.2** *Let  $V$  be finite-dimensional, and let  $W$  be a subspace of  $V$ . Then  $\dim V/W = \dim V - \dim W$ .*

*Proof.* Let  $w_1, \dots, w_r$  be a basis of  $W$ . Extend this to a basis of  $V$ :

$$w_1, \dots, w_r, v_1, \dots, v_s.$$

So  $\dim W = r$  and  $\dim V = r + s$ .

**Claim**  $W + v_1, \dots, W + v_s$  is a basis of  $V/W$ .

*Proof of Claim* We first show the given set of vectors is linearly independent. Suppose

$$\sum_{i=1}^s \lambda_i (W + v_i) = W \quad (\text{the zero vector of } V/W).$$

Then  $\text{LHS} = W + \sum \lambda_i v_i = W$ , so  $\sum \lambda_i v_i \in W$ . Hence there exist scalars  $\mu_j$  such that

$$\sum_{i=1}^s \lambda_i v_i = \sum_{j=1}^r \mu_j w_j.$$

As  $w_1, \dots, w_r, v_1, \dots, v_s$  is a basis, this implies that  $\lambda_i = 0$  for all  $i$ , proving that the set of vectors in the Claim is linearly independent.

Now we prove the set spans  $V/W$ . Let  $W + v \in V/W$ . There are scalars  $\lambda_i, \mu_j$  such that

$$v = \sum_{j=1}^r \mu_j w_j + \sum_{i=1}^s \lambda_i v_i = w + \sum_{i=1}^s \lambda_i v_i,$$

where  $w \in W$  is the first sum. Hence

$$W + v = W + \sum_{i=1}^s \lambda_i v_i = \sum_{i=1}^s \lambda_i (W + v_i).$$

This proves the spanning assertion, and so the Claim is proved.

By the Claim, we have

$$\dim V/W = s = \dim V - \dim W. \quad \square$$

**Example** Let  $V = \mathbb{R}^3$  and  $W = \text{Sp}(e_1 + e_2 + e_3)$ . To find a basis of  $V/W$ , extend the basis  $w = e_1 + e_2 + e_3$  of  $W$  to a basis of  $V$  – say  $w, e_1, e_2$ . Then by the Claim in the above proof,  $W + e_1, W + e_2$  is a basis of  $V/W$ .

### Quotient spaces and linear maps

Let  $T : V \rightarrow V$  be a linear map. Suppose that  $W$  is a  $T$ -invariant subspace of  $V$  (recall this means that  $T(W) \subseteq W$ ). Then we can define the restriction  $T_W : W \rightarrow W$ . We can also define a *quotient map*  $\bar{T} : V/W \rightarrow V/W$  as follows:

$$\bar{T}(W + v) = W + T(v) \quad \forall v \in V.$$

We need to check that  $\bar{T}$  is well-defined; here is the check:

$$\begin{aligned} W + v = W + v' &\Rightarrow v - v' \in W \\ &\Rightarrow T(v - v') \in W \quad (\text{since } T(W) \subseteq W) \\ &\Rightarrow T(v) - T(v') \in W \\ &\Rightarrow W + T(v) = W + T(v') \\ &\Rightarrow \bar{T}(W + v) = \bar{T}(W + v'). \end{aligned}$$

We now show that there is close relationship between the matrices of  $T$ ,  $T_W$  and  $\bar{T}$  with respect to certain bases. Choose of basis  $B_W$  of  $W$ :

$$B_W = \{w_1, \dots, w_r\}.$$

Extend this to a basis  $B$  of  $V$ :

$$B = \{w_1, \dots, w_r, v_1, \dots, v_s\}.$$

As in 5.2, we have a basis  $\bar{B}$  of  $V/W$ :

$$\bar{B} = \{W + v_1, \dots, W + v_s\}.$$

**Proposition 5.3** *Let  $X = [T_W]_{B_W}$  (an  $r \times r$  matrix) and  $Y = [\bar{T}]_{\bar{B}}$  (an  $s \times s$  matrix). Then*

$$[T]_B = \begin{pmatrix} X & Z \\ 0 & Y \end{pmatrix},$$

where  $Z$  is  $r \times s$ .

*Proof.* Let

$$\begin{aligned} T(w_i) &= \sum_{j=1}^r x_{ji} w_j \quad (1 \leq i \leq r), \\ T(v_i) &= \sum_{j=1}^r z_{ji} w_j + \sum_{j=1}^s y_{ji} v_j \quad (1 \leq i \leq s) \end{aligned}$$

Then

$$\begin{aligned} \bar{T}(W + v_i) &= W + \sum_{j=1}^r z_{ji} w_j + \sum_{j=1}^s y_{ji} v_j \\ &= W + \sum_{j=1}^s y_{ji} v_j \\ &= \sum_{j=1}^s y_{ji} (W + v_j). \end{aligned}$$

Hence  $[T_W]_{B_W} = (x_{ij}) = X$ ,  $[\bar{T}]_{\bar{B}} = (y_{ij}) = Y$  and

$$[T]_B = \begin{pmatrix} X & Z \\ 0 & Y \end{pmatrix},$$

where  $Z = (z_{ij})$ .  $\square$

**Example** Let  $V = \mathbb{R}^3$  and  $T : V \rightarrow V$  be given by  $T(v) = Av$  for all  $v \in V$ , where

$$A = \begin{pmatrix} 1 & -2 & 1 \\ -2 & 0 & 2 \\ 1 & 1 & -2 \end{pmatrix}.$$

Let  $w = (1, 1, 1)^T$ . Then  $T(w) = 0$ , so  $W = \text{Sp}(w)$  is a  $T$ -invariant subspace. We extend the basis  $\{w\}$  of  $W$  to a basis  $B = \{w, e_1, e_2\}$  of  $V$ , so we have a basis  $\bar{B} = \{W + e_1, W + e_2\}$  of  $V/W$ . Check that

$$T(e_1) = (1, -2, 1)^T = w - 3e_2, \quad T(e_2) = (-2, 0, 1)^T = w - 3e_1 - e_2.$$

Hence  $\bar{T}(W + e_1) = W - 3e_2$ ,  $\bar{T}(W + e_2) = W - 3e_1 - e_2$ , and so

$$[\bar{T}]_{\bar{B}} = \begin{pmatrix} 0 & -3 \\ -3 & -1 \end{pmatrix}.$$

Finally,

$$[T]_B = \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & -3 \\ 0 & -3 & -1 \end{pmatrix} = \begin{pmatrix} [T_W]_{B_W} & Z \\ 0 & [\bar{T}]_{\bar{B}} \end{pmatrix},$$

where  $Z = (1, 1)$ .

**Corollary 5.4** Let  $T : V \rightarrow V$  be a linear map, and let  $W$  be a  $T$ -invariant subspace of  $V$ . Let  $c(x)$ ,  $c_1(x)$  and  $c_2(x)$  be the characteristic polynomials of  $T$ ,  $T_W$  and  $\bar{T}$ , respectively. Then  $c(x) = c_1(x)c_2(x)$ .

*Proof.* In the notation of Prop. 5.3,

$$\begin{aligned} c(x) &= \det \begin{pmatrix} xI_r - X & -Z \\ 0 & xI_s - Y \end{pmatrix} \\ &= \det(xI_r - X) \det(xI_s - Y) \\ &= c_1(x)c_2(x). \quad \square \end{aligned}$$

## 6 Triangularisation

Triangular matrices are not as easy to compute with as diagonal matrices, but they do have many nice properties. Here are a couple that will be familiar to you from 1st Year.

**Proposition 6.1** Let  $A$  and  $B$  be upper triangular  $n \times n$  matrices:

$$A = \begin{pmatrix} \lambda_1 & & & \\ 0 & \lambda_2 & & * \\ & & \ddots & \\ 0 & 0 & & \lambda_n \end{pmatrix}, \quad B = \begin{pmatrix} \mu_1 & & & \\ 0 & \mu_2 & & * \\ & & \ddots & \\ 0 & 0 & & \mu_n \end{pmatrix}.$$

- (i) The characteristic polynomial of  $A$  is  $\prod_{i=1}^n (x - \lambda_i)$ , the eigenvalues are  $\lambda_1, \dots, \lambda_n$  and the determinant is  $\prod_{i=1}^n \lambda_i$ .
- (ii) The product  $AB$  is also upper triangular, with diagonal entries  $\lambda_1\mu_1, \dots, \lambda_n\mu_n$ .

So the characteristic polynomial of a triangular matrix is  $\prod_1^n (x - \lambda_i)$ , a product of linear factors. The triangularisation theorem shows that the converse is true:

**Theorem 6.2 (Triangularisation Theorem)** Let  $V$  be an  $n$ -dimensional vector space over a field  $F$  and let  $T : V \rightarrow V$  be a linear map. Suppose that characteristic polynomial  $c(x)$  of  $T$  factorizes as a product of linear factors, so that  $c(x) = \prod_1^n (x - \lambda_i)$  with all  $\lambda_i \in F$ . Then there is a basis  $B$  of  $V$  such that the matrix  $[T]_B$  is upper triangular.

We will prove this after making a few remarks on it. First we state the corresponding matrix version:

**Corollary 6.3** Let  $A$  be an  $n \times n$  matrix over a field  $F$ , and suppose the characteristic polynomial of  $A$  factorizes as a product of linear factors. Then  $A$  is similar to an upper triangular matrix over  $F$ .

*Proof.* Let  $V = F^n$  and apply 6.2 to the linear map  $T : V \rightarrow V$  given by  $T(v) = Av$  for all  $v \in V$ .  $\square$

**Remarks** (1) If  $F = \mathbb{C}$  then by the Fundamental Theorem of Algebra, every polynomial over  $F$  factorizes as a product of linear factors. So Corollary 6.3 shows that every  $n \times n$  matrix over  $\mathbb{C}$  can be triangularised.

(2) For other fields this may not be the case; for example for  $F = \mathbb{R}$ , the matrix  $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  has characteristic polynomial  $x^2 + 1$  which has no roots in  $\mathbb{R}$ , hence is not similar to a real triangular matrix.

*Proof of Theorem 6.2.* The proof goes by induction on  $n = \dim V$ . The result is obvious for  $\dim V = 1$ .

Now assume the result for vector spaces of dimension  $n - 1$ . Let  $n = \dim V$ , and  $T : V \rightarrow V$  a linear map whose characteristic polynomial  $c(x)$  factorizes as a product of linear factors. Then  $c(x)$  has a root  $\lambda \in F$ . Let  $w_1 \in V$  be a corresponding eigenvector with  $T(w_1) = \lambda w_1$ , and let  $W = \text{Sp}(w_1)$ , a  $T$ -invariant subspace.

The quotient space  $V/W$  has dimension  $n - 1$  by Prop. 5.2. Consider the quotient map  $\bar{T} : V/W \rightarrow V/W$  (defined by  $\bar{T}(W + v) = W + T(v)$  for  $v \in V$ ). By Cor. 5.4, the characteristic polynomial of  $\bar{T}$  divides  $c(x)$ , hence is also a product of linear factors. Hence by the induction assumption,  $V/W$  has a basis

$$\bar{B} = \{W + v_2, \dots, W + v_n\}$$

such that the matrix  $[\bar{T}]_{\bar{B}}$  is upper triangular. Let  $Y = [\bar{T}]_{\bar{B}}$ . Then  $B = \{w_1, v_2, \dots, v_n\}$  is a basis of  $V$ , and by Prop. 5.3,

$$[T]_B = \begin{pmatrix} \lambda & Z \\ 0 & Y \end{pmatrix}$$

(where  $Z$  is  $1 \times n - 1$  and  $0$  is  $n - 1 \times 1$ ). This matrix  $[T]_B$  is upper triangular, so the induction proof is complete.  $\square$

The above proof gives an algorithm for triangularising a linear map  $T : V \rightarrow V$  (assuming its characteristic polynomial factorizes):

- (1) Find an eigenvector  $w_1$  for  $T$ ; let  $W = \text{Sp}(w_1)$ .
- (2) Find an eigenvector  $W + w_2$  for  $\bar{T} : V/W \rightarrow V/W$ . Let  $W' = \text{Sp}(w_1, w_2)$ .
- (3) Find an eigenvector  $W + w_3$  for  $\bar{\bar{T}} : V/W' \rightarrow V/W'$ .
- (4) Continue, until we have a basis  $B = \{w_1, w_2, w_3, \dots, w_n\}$  of  $V$ . Then  $[T]_B$  is upper triangular.

Here is an example.

**Example** Let  $V = \mathbb{R}^3$  and let  $T : V \rightarrow V$  be defined by  $T(v) = Av$  for all  $v \in V$ , where

$$A = \begin{pmatrix} 3 & 2 & 1 \\ -1 & 0 & 0 \\ -1 & -1 & 0 \end{pmatrix}.$$

Check that the characteristic polynomial of  $T$  is  $(x - 1)^3$ .

- (1) We find an eigenvector  $w_1 = (1, -1, 0)^T$ . Let  $W = \text{Sp}(w_1)$ .
- (2) Extend  $w_1$  to a basis  $C = \{w_1, e_2, e_3\}$  of  $V$ . Then  $\bar{C} = \{W + e_2, W + e_3\}$  is a basis of  $V/W$ . Compute that

$$[\bar{T}]_{\bar{C}} = \begin{pmatrix} 2 & 1 \\ -1 & 0 \end{pmatrix}.$$



This matrix has an eigenvector  $(1, -1)^T$ , which corresponds to an eigenvector  $W + e_2 - e_3$  of  $\bar{T}$ . So in the algorithm we can take  $w_2 = e_2 - e_3$ .

(3) Thus our final triangularising basis is  $B = \{w_1, w_2, e_3\}$  (the third vector can be any vector that makes a basis with  $w_1, w_2$ ): the matrix  $[T]_B$  is upper triangular (with 1's on the diagonal, as 1 is the only eigenvalue of  $T$ ). Also, if  $P$  is the matrix with columns  $w_1, w_2, e_3$ , then  $P^{-1}AP$  is upper triangular.