

ii) $f(\underline{x}) = \sqrt{1 + \underline{x}^T Q \underline{x}}$ is convex, given $Q \succeq 0$

In previous weeks, we have shown that

$h(y) = \sqrt{1 + y^2}$ is convex. ✓

$$h'(y) = \frac{1}{2} \frac{2y}{\sqrt{1+y^2}} > 0 \text{ for } y > 0 \rightarrow \text{non-decreasing}$$

$$h''(y) = \frac{\sqrt{1+y^2} - \frac{y^2}{\sqrt{1+y^2}}}{(1+y^2)} = \frac{1}{(1+y^2)^{3/2}} > 0$$

$$f(\underline{x}) = \sqrt{1 + \underline{x}^T Q \underline{x}} = \sqrt{1 + y^2}, \text{ where } y^2 = \underline{x}^T Q \underline{x}$$
$$\Rightarrow y = g(\underline{x}) = \sqrt{\underline{x}^T Q \underline{x}}$$

but $Q \succeq 0$, and we have seen in Week 2

$$Q = U^T \underbrace{D}_{\text{Diagonal}} U \rightarrow D = \begin{bmatrix} d_1 & & 0 \\ & \ddots & \\ 0 & & d_n \end{bmatrix} \quad d_i \geq 0$$

$$Q = U^T \sqrt{D} \sqrt{D} U \Rightarrow \underline{x}^T U^T \sqrt{D} \sqrt{D} U \underline{x}$$

$$\stackrel{\parallel}{\underbrace{\underline{x}^T Q \underline{x}}_{\parallel \sqrt{\underline{x}^T U^T \sqrt{D} \sqrt{D} U \underline{x}}}}$$

$$\stackrel{\parallel}{\parallel L \underline{x} \parallel}$$

↳ where $L = \sqrt{D} U$

But we know that $\|\cdot\|$ is convex function and

$\underline{x} \longrightarrow L\underline{x}$ is a linear transformation

$\Rightarrow \|L\underline{x}\|$ is convex in \underline{x}

$g(\underline{x}) : \mathbb{R}^M \longrightarrow \mathbb{R}^+$ $g := \|L\underline{x}\|$ is convex

$$f(\underline{x}) = \sqrt{\underline{x}^T Q \underline{x} + 1} = \underbrace{h \circ g}_{\substack{\text{"} \\ \sqrt{1+y^2} \\ \text{"}}}(\underline{x}) \quad \text{convex}$$

$\underbrace{\hspace{1.5cm}}_{\text{"} \|\underline{x}\| \text{"}}$

because of composing $h \circ g$ w/ g convex and h convex and non-decreasing.

$$\text{min} \quad -x_1 x_2 x_3$$

$$\text{s.t.} \quad x_1 + 3x_2 + 6x_3 \leq 48$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0$$

$$-x_3 \leq 0$$