



Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Eric Li
Assignment title: Coursework 1 Autumn 24 Drop Box
Submission title: 60005.pdf
File name: 60005.pdf
File size: 262.79K
Page count: 9
Word count: 2,372
Character count: 9,144
Submission date: 14-Nov-2024 11:54AM (UTC+0000)
Submission ID: 242604333

MATH60005 Optimisation Coursework

CID: 02216531

14 November 2024

1. Unconstrained Optimisation

a) We are given the function $f(x_1, x_2) = 8x_1 + 12x_2 + x_1^2 - 2x_2^2$.

i) Let $\mathbf{x} = (x_1, x_2)$. We can rewrite f in standard form $f(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x} + 2\mathbf{b}^T \mathbf{x} + c$ as

$$f(\mathbf{x}) = \mathbf{x}^T \begin{pmatrix} 1 & 0 \\ 0 & -2 \end{pmatrix} \mathbf{x} + 2 \begin{pmatrix} 4 \\ 6 \end{pmatrix}^T \mathbf{x} + 0.$$

The matrix $\mathbf{A} = \begin{pmatrix} 1 & 0 \\ 0 & -2 \end{pmatrix}$ has eigenvalues 1, -2. Since have opposite signs, \mathbf{A} is not positive definite, and by a lemma in lecture notes, f is not coercive.

Alternatively, by definition, f is coercive if $f \rightarrow \infty$ when $\|\mathbf{x}\| \rightarrow \infty$. However, we note that when $x_2 \rightarrow \infty$ while keeping x_1 constant, we have $f \rightarrow -\infty$ due to the $-2x_2^2$ term. Therefore, f is not coercive.

ii) We compute the first order partial derivatives of f as

$$\frac{\partial f}{\partial x_1} = 8 + 2x_1, \quad \frac{\partial f}{\partial x_2} = 12 - 4x_2.$$

Setting them to zero gives $x_1 = -4, x_2 = 3$. Therefore, the stationary point of f is $(-4, 3)$.

To classify this point, we compute the Hessian of f . We have

$$\frac{\partial^2 f}{\partial x_1^2} = 2, \quad \frac{\partial^2 f}{\partial x_1^2} = -4, \quad \frac{\partial^2 f}{\partial x_1 \partial x_2} = 0, \quad \frac{\partial^2 f}{\partial x_2 \partial x_1} = 0.$$

The Hessian is therefore given by $\begin{pmatrix} 2 & 0 \\ 0 & -4 \end{pmatrix}$ which has eigenvalues 2, -4. Since they have opposite signs, the stationary point $(-4, 3)$ is a saddle point.

b) Yes, it is. We first notice that the matrix $\mathbf{A} \mathbf{x}$ can be rewritten as $\mathbf{A} = \text{diag}(\mathbf{d}) - \mathbf{d} \mathbf{d}^T$.

Therefore, for every $\mathbf{x} \in \mathbb{R}^n$, we have $\mathbf{x}^T \mathbf{A} \mathbf{x} = \mathbf{x}^T \text{diag}(\mathbf{d}) \mathbf{x} - \mathbf{x}^T \mathbf{d} \mathbf{d}^T \mathbf{x} = \mathbf{x}^T \text{diag}(\mathbf{d}) \mathbf{x} - (\mathbf{x}^T \mathbf{d})^2$.

Multiplying out these two terms, we get

$$\mathbf{x}^T \text{diag}(\mathbf{d}) \mathbf{x} = \sum_{i=1}^n d_i x_i^2, \quad (\mathbf{x}^T \mathbf{d})^2 = \left(\sum_{i=1}^n d_i x_i \right)^2.$$

Since $\mathbf{d} \in \Delta_n$, we have $d_i \geq 0$ and $\sum_{i=1}^n d_i = 1$. Therefore, $d_i x_i^2 \geq 0$ and we could apply the Cauchy-Schwarz inequality to get