

Background and Scope
oooooo

Statistical Models
oooooooo

Using Models
oooo



Lecture 01: Statistical Models

Statistical Modelling I

Dr. Riccardo Passeggeri

Background and Scope
ooooooo

Statistical Models
oooooooo

Using Models
oooo

Outline

1. Background and Scope

2. Statistical Models

3. Using Models

Background and Scope
●○○○○○

Statistical Models
○○○○○○○

Using Models
○○○○

Background and Scope

Background and Scope
●●○○○

Statistical Models
○○○○○○○


Using Models
○○○○

Science

How scientists do science?

- ▶ Guess
- ▶ Compute the consequences of the guess
- ▶ Compare them with experiments

The Scientific Method and Statistics




Statistical modelling

Where does the randomness come from?

- ▶ measurement errors and technical noises (Higgs boson discovery announcement
<https://www.youtube.com/watch?v=0CugLD9HF94>)
- ▶ impossibility of repeating the exact same experiment
- ▶ impossibility of repeating the experiment

Statistical Answers to Scientific Questions (for this Module!)

Quantify distributions





SUMMARY
STATISTICS

Compare distributions



Predict observations



Other common tasks: clustering observations or variables.

Module Outline

- ▶ **1st half:** deriving, evaluating and applying estimators, confidence intervals and hypothesis tests based on parametric models.
- ▶ **2nd half:** deriving, evaluating and applying estimators, confidence intervals and hypothesis tests based on the theory of linear models.

Background and Scope
oooooooo

Statistical Models
●oooooooo

Using Models
ooooo

Statistical Models


Some Conventions

Key: observed data y is realization of random Y .
 $y = (y_1, \dots, y_m)$ $Y = (Y_1, \dots, Y_m)$


- ▶ Random variables: X, Y, Z
- ▶ Data: x, y, z
- ▶ Parameters: θ, α, β
- ▶ Estimators: $\hat{\theta}, \hat{\alpha}, \hat{\beta}$

- ▶ Outcome: Y, y
- ▶ Covariate: X, x

DEPENDENT
INDEPENDENT



$$Y = \beta_0 + \beta_1 X + \varepsilon$$



$$Y : (\Omega, \mathcal{F}, P) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))$$

$$Y : \omega \mapsto Y(\omega)$$

Parametric Models

$$N(\theta, 1), \theta \in \mathbb{R}$$

Let P_θ be a probability distribution with parameter θ . For example,

SINCE OBSERVED DATA ARE THE OUTCOME OF AN UNDERLYING R.V. Y , THEN
A STATISTICAL MODEL IS THE SPECIFICATION OF THE DISTRIBUTION OF Y
UP TO A PARAMETER $\theta \in \Theta$

Definition:

A **statistical model** is a collection of probability distributions $\{P_\theta : \theta \in \Theta\}$ over a sample space. The set Θ of all possible parameter values is called the **parameter space**. In this module, $\Theta \subseteq \mathbb{R}^p$, so that we consider **parametric models**.

A statistical model is generally required to be such that distinct parameter values give rise to distinct distributions, i.e.

$$P_{\theta_1} = P_{\theta_2} \Rightarrow \theta_1 = \theta_2$$

$$N(\theta^2, 1), \theta \in \mathbb{R}$$

Independent and Identically Distributed (iid)

Important special case:

- ▶ $y = (y_1, \dots, y_n) \in \mathbb{R}^n \Rightarrow Y = (Y_1, \dots, Y_n)$ is a random vector
- ▶ Then statistical model specifies the joint distribution of Y_1, \dots, Y_n up to $\theta \in \Theta$
- ▶ Y_1, \dots, Y_n is a **random sample** if Y_i 's are **iid**

Example: Guess the Weight

Observed data




Statistical model

$$N(543.4, \sigma^2), \sigma \in [0, \infty)$$

- ▶ People guess the weight of an ox
- ▶ n guesses y_1, \dots, y_n
- ▶ The true weight is 543.4 kg

Example: Clinical Trial

Observed data



- ▶ Compare new and old treatment
 ⇒ or vaccine
- ▶ n treatment assignments x_1, \dots, x_n
- ▶ n times until recovery y_1, \dots, y_n

Statistical model

$$x_i = \begin{cases} 0 & \text{NO DRUG} \\ 1 & \text{DRUG} \end{cases}$$

$$y_i | x_i = 0 \sim ? N(\mu_0, 1)$$

$$y_i | x_i = 1 \sim ? N(\mu_1, 1)$$

SCIENTIFIC Q: DOES THE DRUG HAVE AN EFFECT?

STATISTICAL Q: DOES μ_0 AND μ_1 DIFFER "SIGNIFICANTLY"?

Example: Do Taller People Have Higher Incomes?


Observed data



- ▶ Compare income across heights
- ▶ n heights x_1, \dots, x_n
- ▶ n incomes y_1, \dots, y_n

Statistical model

$$Y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$



STAT Q: IS β_1 SIGNIFICANTLY
DIFFERENT FROM 0?

Background and Scope
○○○○○○○

Statistical Models
○○○○○○○

Using Models
●○○○

Using Models

Remember: Statistical Answers to Scientific Questions

Quantify distributions

OX WEIGHT
OUESS

Compare distributions

CLINICAL
TRIAL

Predict observations

TALLER PEOPLE
HIGHER INCOME

Assessing Statistical Models

"All models are wrong, but some are useful."

We want our parametric model to

- ▶ agree well with observed data
- ▶ be simple and interpretable

Background and Scope
○○○○○○

Statistical Models
○○○○○○○

Using Models
○○○●

Looking Ahead

How do we “know” which estimators, confidence intervals, and hypothesis tests to use?