

# Analysis 1A

## Lecture 13

Cauchy Theorem  $\iff$  BW - Theorem

Ajay Chandra

### Proposition 3.39

If  $a_n \rightarrow a$  then any subsequence  $a_{n(i)} \rightarrow a$  as  $i \rightarrow \infty$ .

Next, we are going to look at the relationship between these two theorems:

### Theorem 3.27 - Cauchy Theorem

If  $(a_n)$  is a Cauchy sequence of real numbers then  $a_n$  converges.

### Theorem 3.34 - Bolzano-Weierstrass

If  $(a_n)$  is a *bounded* sequence of real numbers then it has a *convergent subsequence*.

**Bolzano-Weierstrass  $\Rightarrow$  the Cauchy Theorem**

Before continuing, a basic but useful trick:

### Lemma 3.40

Fix  $c > 0$ . Then  $a_n \rightarrow a$  if and only if

$$\forall \epsilon > 0 \exists N_\epsilon \in \mathbb{N}_{>0} \text{ such that } n \geq N_\epsilon \implies |a_n - a| < c\epsilon \quad (\star)$$

Before continuing, a basic but useful trick:

### Lemma 3.40

Fix  $c > 0$ . Then  $a_n \rightarrow a$  if and only if

$$\forall \epsilon > 0 \exists N_\epsilon \in \mathbb{N}_{>0} \text{ such that } n \geq N_\epsilon \implies |a_n - a| < c\epsilon \quad (\star)$$

**Proof:**  $\Rightarrow$ . Assume  $a_n \rightarrow a$ , we prove  $(\star)$ .

Before continuing, a basic but useful trick:

### Lemma 3.40

Fix  $c > 0$ . Then  $a_n \rightarrow a$  if and only if

$$\forall \epsilon > 0 \exists N_\epsilon \in \mathbb{N}_{>0} \text{ such that } n \geq N_\epsilon \implies |a_n - a| < c\epsilon \quad (\star)$$

**Proof:**  $\Rightarrow$ . Assume  $a_n \rightarrow a$ , we prove  $(\star)$ . Fix  $\epsilon > 0$

Before continuing, a basic but useful trick:

### Lemma 3.40

Fix  $c > 0$ . Then  $a_n \rightarrow a$  if and only if

$$\forall \epsilon > 0 \exists N_\epsilon \in \mathbb{N}_{>0} \text{ such that } n \geq N_\epsilon \implies |a_n - a| < c\epsilon \quad (\star)$$

**Proof:**  $\Rightarrow$ . Assume  $a_n \rightarrow a$ , we prove  $(\star)$ . Fix  $\epsilon > 0$  and let  $\epsilon' := c\epsilon$ . Then by the definition of convergence applied to  $\epsilon' > 0$  we find

$$\exists N \in \mathbb{N}_{>0} : n \geq N \implies |a_n - a| < \epsilon',$$

which is  $(\star)$ .



Before continuing, a basic but useful trick:

### Lemma 3.40

Fix  $c > 0$ . Then  $a_n \rightarrow a$  if and only if

$$\forall \epsilon > 0 \exists N_\epsilon \in \mathbb{N}_{>0} \text{ such that } n \geq N_\epsilon \implies |a_n - a| < c\epsilon \quad (\star)$$

**Proof:**  $\Rightarrow$ . Assume  $a_n \rightarrow a$ , we prove  $(\star)$ . Fix  $\epsilon > 0$  and let  $\epsilon' := c\epsilon$ . Then by the definition of convergence applied to  $\epsilon' > 0$  we find

$$\exists N \in \mathbb{N}_{>0} : n \geq N \implies |a_n - a| < \epsilon',$$

which is  $(\star)$ .

$\Leftarrow$ . We assume  $(\star)$  and show  $a_n \rightarrow a$ .

Before continuing, a basic but useful trick:

### Lemma 3.40

Fix  $c > 0$ . Then  $a_n \rightarrow a$  if and only if

$$\forall \epsilon > 0 \exists N_\epsilon \in \mathbb{N}_{>0} \text{ such that } n \geq N_\epsilon \implies |a_n - a| < c\epsilon \quad (\star)$$

**Proof:**  $\Rightarrow$ . Assume  $a_n \rightarrow a$ , we prove  $(\star)$ . Fix  $\epsilon > 0$  and let  $\epsilon' := c\epsilon$ . Then by the definition of convergence applied to  $\epsilon' > 0$  we find

$$\exists N \in \mathbb{N}_{>0} : n \geq N \implies |a_n - a| < \epsilon',$$

which is  $(\star)$ .

$\Leftarrow$ . We assume  $(\star)$  and show  $a_n \rightarrow a$ . Fix  $\epsilon > 0$ .

Before continuing, a basic but useful trick:

### Lemma 3.40

Fix  $c > 0$ . Then  $a_n \rightarrow a$  if and only if

$$\forall \epsilon > 0 \exists N_\epsilon \in \mathbb{N}_{>0} \text{ such that } n \geq N_\epsilon \implies |a_n - a| < c\epsilon \quad (\star)$$

**Proof:**  $\Rightarrow$ . Assume  $a_n \rightarrow a$ , we prove  $(\star)$ . Fix  $\epsilon > 0$  and let  $\epsilon' := c\epsilon$ . Then by the definition of convergence applied to  $\epsilon' > 0$  we find

$$\exists N \in \mathbb{N}_{>0} : n \geq N \implies |a_n - a| < \epsilon',$$

which is  $(\star)$ .

$\Leftarrow$ . We assume  $(\star)$  and show  $a_n \rightarrow a$ . Fix  $\epsilon > 0$ . Set  $\epsilon' = \epsilon/c > 0$ . Then  $(\star)$  applied to  $\epsilon' > 0$  implies

$$\exists N \in \mathbb{N}_{>0} \text{ such that } n \geq N_\epsilon \implies |a_n - a| < c\epsilon' = \epsilon.$$

Before continuing, a basic but useful trick:

### Lemma 3.40

Fix  $c > 0$ . Then  $a_n \rightarrow a$  if and only if

$$\forall \epsilon > 0 \exists N_\epsilon \in \mathbb{N}_{>0} \text{ such that } n \geq N_\epsilon \implies |a_n - a| < c\epsilon \quad (\star)$$

**Proof:**  $\Rightarrow$ . Assume  $a_n \rightarrow a$ , we prove  $(\star)$ . Fix  $\epsilon > 0$  and let  $\epsilon' := c\epsilon$ . Then by the definition of convergence applied to  $\epsilon' > 0$  we find

$$\exists N \in \mathbb{N}_{>0} : n \geq N \implies |a_n - a| < \epsilon',$$

which is  $(\star)$ .

$\Leftarrow$ . We assume  $(\star)$  and show  $a_n \rightarrow a$ . Fix  $\epsilon > 0$ . Set  $\epsilon' = \epsilon/c > 0$ . Then  $(\star)$  applied to  $\epsilon' > 0$  implies

$$\exists N \in \mathbb{N}_{>0} \text{ such that } n \geq N_\epsilon \implies |a_n - a| < c\epsilon' = \epsilon.$$

### Warning!

Do not let  $c$  depend on  $\epsilon$  (nor  $N$  or  $n$ )!

**the Cauchy Theorem  $\Rightarrow$  Bolzano-Weierstrass**

## **the Cauchy Theorem $\Rightarrow$ Bolzano-Weierstrass**

Take a bounded sequence  $(a_n)$ . We want to find a Cauchy subsequence, which will therefore be convergent.

## the Cauchy Theorem $\Rightarrow$ Bolzano-Weierstrass

Take a bounded sequence  $(a_n)$ . We want to find a Cauchy subsequence, which will therefore be convergent.

Since  $a_n \in [-R, R] \forall n$ , then either

- 1  $a_n \in [-R, 0]$  for infinitely many  $n$  or
- 2  $a_n \in [0, R]$  for infinitely many  $n$ , (or both).

## the Cauchy Theorem $\Rightarrow$ Bolzano-Weierstrass

Take a bounded sequence  $(a_n)$ . We want to find a Cauchy subsequence, which will therefore be convergent.

Since  $a_n \in [-R, R] \forall n$ , then either

- 1  $a_n \in [-R, 0]$  for infinitely many  $n$  or
- 2  $a_n \in [0, R]$  for infinitely many  $n$ , (or both).

Pick one of these intervals with infinite number of  $a_n$  s; call it  $[A_1, B_1]$  of length  $R$ .



## the Cauchy Theorem $\Rightarrow$ Bolzano-Weierstrass

Take a bounded sequence  $(a_n)$ . We want to find a Cauchy subsequence, which will therefore be convergent.

Since  $a_n \in [-R, R] \forall n$ , then either

- 1  $a_n \in [-R, 0]$  for infinitely many  $n$  or
- 2  $a_n \in [0, R]$  for infinitely many  $n$ , (or both).

Pick one of these intervals with infinite number of  $a_n$  s; call it  $[A_1, B_1]$  of length  $R$ .

Now subdivide again; call  $[A_2, B_2]$  one of the intervals  $[A_1, \frac{A_1+B_1}{2}]$  or  $[\frac{A_1+B_1}{2}, B_1]$  contains  $a_n$  for infinitely many  $n$ , with length  $R/2$ .

## the Cauchy Theorem $\Rightarrow$ Bolzano-Weierstrass

Take a bounded sequence  $(a_n)$ . We want to find a Cauchy subsequence, which will therefore be convergent.

Since  $a_n \in [-R, R] \forall n$ , then either

- 1  $a_n \in [-R, 0]$  for infinitely many  $n$  or
- 2  $a_n \in [0, R]$  for infinitely many  $n$ , (or both).

Pick one of these intervals with infinite number of  $a_n$  s; call it  $[A_1, B_1]$  of length  $R$ .

Now subdivide again; call  $[A_2, B_2]$  one of the intervals  $[A_1, \frac{A_1+B_1}{2}]$  or  $[\frac{A_1+B_1}{2}, B_1]$  contains  $a_n$  for infinitely many  $n$ , with length  $R/2$ . Continue this procedure inductively....

## the Cauchy Theorem $\Rightarrow$ Bolzano-Weierstrass

Take a bounded sequence  $(a_n)$ . We want to find a Cauchy subsequence, which will therefore be convergent.

Since  $a_n \in [-R, R] \forall n$ , then either

- 1  $a_n \in [-R, 0]$  for infinitely many  $n$  or
- 2  $a_n \in [0, R]$  for infinitely many  $n$ , (or both).

Pick one of these intervals with infinite number of  $a_n$  s; call it  $[A_1, B_1]$  of length  $R$ .

Now subdivide again; call  $[A_2, B_2]$  one of the intervals  $[A_1, \frac{A_1+B_1}{2}]$  or  $[\frac{A_1+B_1}{2}, B_1]$  contains  $a_n$  for infinitely many  $n$ , with length  $R/2$ . Continue this procedure inductively....

## **the Cauchy Theorem $\Rightarrow$ Bolzano-Weierstrass** continued

We get a sequence of intervals  $[A_n, B_n]$  of length  $2^{1-n}R$  which are nested – i.e.  $[A_{k+1}, B_{k+1}] \subseteq [A_k, B_k]$  – with each containing  $a_n$  for infinitely many  $n$ .